



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### **Usage guidelines**

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>









**INTEGRAL CALCULUS.**

**Cambridge:**

PRINTED BY C. J. CLAY, M.A. AT THE UNIVERSITY PRESS,

**FOR MACMILLAN AND Co.**

**LONDON: BELL AND DALDY.**

**OXFORD: J. H. AND JAMES PARKER.**

**EDINBURGH: EDMONSTON AND DOUGLAS.**

**DUBLIN: WILLIAM ROBERTSON.**


**GLASGOW: JAMES MACLEHOSE.**

A TREATISE ON THE  
INTEGRAL CALCULUS

AND ITS

APPLICATIONS

WITH NUMEROUS EXAMPLES.



BY I. TODHUNTER, M.A.

FELLOW AND ASSISTANT TUTOR OF ST JOHN'S COLLEGE,  
CAMBRIDGE.

Cambridge:  
MACMILLAN AND CO.  
1857.

182. 6. 6.





## P R E F A C E.

IN writing the present treatise on the INTEGRAL CALCULUS, the object has been to produce a work at once elementary and complete—adapted for the use of beginners, and sufficient for the wants of advanced students. In the selection of the propositions, and in the mode of establishing them, I have endeavoured to exhibit fully and clearly the principles of the subject, and to illustrate all their most important results. The process of *summation* has been repeatedly brought forward, with the view of securing the attention of the student to the notions which form the true foundation of the Integral Calculus itself, as well as of its most valuable applications. Considerable space has been devoted to the investigations of the lengths and areas of curves and of the volumes of solids, and an attempt has been made to explain those difficulties which usually perplex beginners—especially with reference to the *limits* of integrations.

The transformation of multiple integrals is one of the most interesting parts of the subject, and the experience of teachers shews that the usual modes of treating it are not free from obscurity. I have therefore adopted a method different from those of previous elementary writers, and have endeavoured to

render it easily intelligible by full detail, and by the solution of several problems.

In order that the student may find in the volume all that he requires, a large collection of examples for exercise has been appended to the different chapters. These examples have been selected from the College and University Examination Papers, and have been carefully verified, so that it is hoped that few errors will be found among them. The short introduction to the Integral Calculus which is given at the end of my treatise on the Differential Calculus, has been incorporated in the present work so as to render it complete in itself. The student has been occasionally referred to the Differential Calculus for such information as he would require before commencing the study of the Integral Calculus.

I. TODHUNTER.

ST JOHN'S COLLEGE,  
*March 1857.*

## CONTENTS.

CHAP.		PAGE
I.	Meaning of Integration. Examples . . . . .	1
II.	Rational Fractions . . . . .	22
III.	Formulæ of Reduction . . . . .	38
IV.	Miscellaneous Remarks . . . . .	48
V.	Double Integration . . . . .	64
VI.	Lengths of Curves . . . . .	71
VII.	Areas of Plane Curves and of Surfaces . . . . .	103
VIII.	Volumes of Solids . . . . .	142
IX.	Differentiation of an Integral with respect to any quantity which it may involve . . . . .	165
X.	Elliptic Integrals . . . . .	174
XI.	Change of the Variables in a Multiple Integral . . . . .	180
XII.	Definite Integrals . . . . .	212
XIII.	Expansion of Functions in Trigonometrical Series . . . . .	250
XIV.	Application of the Integral Calculus to Questions of Mean Value and Probability . . . . .	262



# INTEGRAL CALCULUS.

## CHAPTER I.

### MEANING OF INTEGRATION. EXAMPLES.

1. IN the Differential Calculus we have a system of rules by means of which we deduce from any given function a second function called the differential coefficient of the former; in the Integral Calculus we have to return from the differential coefficient to the function from which it was deduced. We do not say that this is the *object* of the Integral Calculus, for the fundamental problem of the subject is to effect the summation of a certain infinite series of indefinitely small terms; but for the solution of this problem we must generally know the function of which a given function is the differential coefficient. This we now proceed to shew.

2. Let  $\phi(x)$  denote any function of  $x$  which remains finite and continuous for all values of  $x$  comprised between two fixed values  $a$  and  $b$ . Let  $x_1, x_2, \dots, x_{n-1}$  be a series of values lying between  $a$  and  $b$ , so that  $a, x_1, x_2, \dots, x_{n-1}, b$  are in order of magnitude ascending or descending. We propose then to find the limit of the series

$$(x_1 - a)\phi(a) + (x_2 - x_1)\phi(x_1) + (x_3 - x_2)\phi(x_2) \dots \dots \\ + (b - x_{n-1})\phi(x_{n-1}),$$

when  $x_1 - a, x_2 - x_1, \dots, b - x_{n-1}$  are all diminished without limit, and consequently  $n$  increased without limit.

Put  $x_1 - a = h_1, x_2 - x_1 = h_2, \dots, b - x_{n-1} = h_n$ ; thus the series may be written

$$h_1 \phi(a) + h_2 \phi(x_1) \dots + h_{n-1} \phi(x_{n-2}) + h_n \phi(x_{n-1}),$$

and may be denoted by  $\Sigma h\phi(x)$ , for it is the sum of a number of terms of which  $h\phi(x)$  may be taken as the type. Since each of the terms of which  $h$  is the type may be considered as the difference between two values successively ascribed to the variable  $x$ , we may also use the symbol  $\phi(x) \Delta x$  as the type of the terms to be summed, and  $\Sigma\phi(x) \Delta x$  for the sum.

We may shew at once that  $\Sigma\phi(x) \Delta x$  can never exceed a certain finite quantity. For let  $A$  denote the numerically greatest value which  $\phi(x)$  can have when  $x$  lies between  $a$  and  $b$ ; then  $\Sigma\phi(x) \Delta x$  is numerically less than  $(h_1 + h_2 + \dots + h_n) A$ , that is less than  $(b - a) A$ .

We now proceed to determine the limit of  $\Sigma\phi(x) \Delta x$ . Let  $\psi(x)$  be such a function of  $x$  that  $\phi(x)$  is the differential coefficient of it with respect to  $x$ . Then we know that the limit of  $\frac{\psi(x+h) - \psi(x)}{h}$  when  $h$  is indefinitely diminished is  $\phi(x)$ . Hence we may put

$$\psi(x_1) - \psi(a) = h_1\{\phi(a) + \rho_1\},$$

$$\psi(x_2) - \psi(x_1) = h_2\{\phi(x_1) + \rho_2\},$$

.....

$$\psi(x_{n-1}) - \psi(x_{n-2}) = h_{n-1}\{\phi(x_{n-2}) + \rho_{n-1}\},$$

$$\psi(b) - \psi(x_{n-1}) = h_n\{\phi(x_{n-1}) + \rho_n\},$$

where  $\rho_1, \rho_2, \dots, \rho_n$  ultimately vanish. From these equations we have by addition

$$\psi(b) - \psi(a) = \Sigma\phi(x) \Delta x + \Sigma h\rho.$$

Now  $\Sigma h\rho$  is less than  $(b - a) \rho'$  where  $\rho'$  denotes the greatest of the quantities  $\rho_1, \rho_2, \dots, \rho_n$ ; hence  $\Sigma h\rho$  ultimately vanishes, and we obtain this result—the limit of  $\Sigma\phi(x) \Delta x$  when each of the quantities of which  $\Delta x$  is the type diminishes indefinitely is  $\psi(b) - \psi(a)$ .

3. The notation used to express the preceding result is

$$\int_a^b \phi(x) dx = \psi(b) - \psi(a);$$

the symbol  $\int$  is an abbreviation of the word "sum," and  $dx$  represents the  $\Delta x$  of  $\Sigma\phi(x) \Delta x$ .

4. Suppose that  $h_1, h_2, \dots, h_n$  are all equal; then each of them is equal to  $\frac{b-a}{n}$ , and  $x_r$  is equal to  $a + \frac{r}{n}(b-a)$ .

Hence  $\int_a^b \phi(x) dx$  is equivalent to the following direction—“divide  $b-a$  into  $n$  equal parts, each part being  $h$ ; in  $\phi(x)$  substitute for  $x$  successively

$$a, a+h, a+2h, \dots, a+(n-1)h;$$

add these values together, multiply the sum by  $h$  and then diminish  $h$  without limit.” If these operations are performed we shall have as the result  $\psi(b) - \psi(a)$ .

5. A single term such as  $\phi(x) \Delta x$  is frequently called an *element*. It may be observed that the *limit* of  $\Sigma \phi(x) \Delta x$  will not be altered in value if we omit a *finite* number of its elements, or add a finite number of similar elements; for in the limit each element is indefinitely small, and a *finite* number of indefinitely small quantities ultimately vanishes.

6. The above process is called *Integration*; the quantity  $\int_a^b \phi(x) dx$  is called a *definite integral*, and  $a$  and  $b$  are called the *limits of the integral*. Since the value of this definite integral is  $\psi(b) - \psi(a)$  we must, when a function  $\phi(x)$  is to be integrated between assigned limits, first ascertain the function  $\psi(x)$  of which  $\phi(x)$  is the differential coefficient. To express the connexion between  $\phi(x)$  and  $\psi(x)$  we have

$$\phi(x) = \frac{d\psi(x)}{dx},$$


and this is also denoted by the equation

$$\int \phi(x) dx = \psi(x).$$

In such an equation as the last, where we have no limits assigned, we merely assert that  $\psi(x)$  is the function from which  $\phi(x)$  can be obtained by differentiation;  $\psi(x)$  is here called the *indefinite integral* of  $\phi(x)$ .



7. The problem of finding the areas of curves was one of those which gave rise to the Integral Calculus, and furnishes an illustration of the preceding articles.



Let  $DPE$  be a curve of which the equation is  $y = \phi(x)$ , and suppose it required to find the area included between this curve, the axis of  $x$ , and the ordinates corresponding to the abscissæ  $a$  and  $b$ . Let  $OA = a$ ,  $OB = b$ ; divide the space  $AB$  into  $n$  equal intervals and draw ordinates at the points of division. Suppose  $OM = a + (r-1)h$ , then the area of the parallelogram  $PMNp$  is

$$h\phi\{a + (r-1)h\}.$$

The sum found by assigning to  $r$  in this expression all values from 1 to  $n$  differs from the required area of the curve by the sum of all the portions similar to the triangle  $PQp$ , and as this last sum is obviously less than the greatest of the figures of which  $PMNQ$  is one, we can, by sufficiently diminishing  $h$ , obtain a result differing as little as we please from the required area. Therefore the area of the curve is the limit of the series

$$h[\phi(a) + \phi(a+h) + \phi(a+2h) \dots \phi\{a + (n-1)h\}],$$

and is equal to  $\psi(b) - \psi(a)$ .

8. If  $\psi(x)$  be the function from which  $\phi(x)$  springs by differentiation, we denote this by the equation

$$\int \phi(x) dx = \psi(x),$$

and we now proceed to methods of finding  $\psi(x)$  when  $\phi(x)$  is

given. We have shewn, *Dif. Cal.* Art. 102, that if two functions have the same differential coefficient with respect to a variable they can only differ by some constant quantity; hence if  $\psi(x)$  be a function having  $\phi(x)$  for its differential coefficient, then  $\psi(x) + C$ , where  $C$  is any quantity independent of  $x$ , is the only form that can have the same differential coefficient. Hence, hereafter, when we assert that any function is the integral of a proposed function, we may if we please add to such integral any constant quantity.

Integration then will for some time appear to be merely the *inverse* of differentiation, and we might have so defined it; we have however preferred to introduce at the beginning the notion of *summation* because it occurs in many of the most important applications of the subject.

### 9. Immediate integration.

When a function is recognized to be the differential coefficient of another function we know of course the integral of the first. The following list gives the integrals of the different simple functions:

$$\int x^m dx = \frac{x^{m+1}}{m+1},$$

$$\int \frac{dx}{x} = \log x,$$

$$\int a^x dx = \frac{a^x}{\log_a a},$$

$$\int e^x dx = e^x,$$

$$\int \sin x dx = -\cos x,$$

$$\int \cos x dx = \sin x,$$

$$\int \frac{dx}{\cos^2 x} = \tan x,$$

$$\int \frac{dx}{\sin^2 x} = -\cot x,$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a},$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a}.$$

### 10. *Integration by substitution.*

The process of integration is sometimes facilitated by substituting for the variable some function of a new variable. Suppose  $\phi(x)$  the function to be integrated, and  $a$  and  $b$  the limits of the integral. It is evident that we may suppose  $x$  to be a function of a new variable  $z$ , provided that the function chosen is capable of assuming all the values of  $x$  required in the integration. Put then  $x = f(z)$ , and let  $\alpha'$  and  $\beta'$  be the values of  $z$ , which make  $f(z)$  or  $x$  equal to  $a$  and  $b$  respectively; thus  $a = f(\alpha')$  and  $b = f(\beta')$ . Now suppose that  $\psi(x)$  is the function of which  $\phi(x)$  is the differential coefficient, that is  $\phi(x) = \frac{d\psi(x)}{dx}$ ; then

$$\begin{aligned} \int_a^b \phi(x) dx &= \psi(b) - \psi(a) \\ &= \psi\{f(\beta')\} - \psi\{f(\alpha')\}. \end{aligned}$$

But by the principles of the Differential Calculus,

$$\frac{d\psi\{f(z)\}}{dz} = \phi\{f(z)\} f'(z);$$

therefore 
$$\begin{aligned} \psi\{f(\beta')\} - \psi\{f(\alpha')\} &= \int_{\alpha'}^{\beta'} \phi\{f(z)\} f'(z) dz \\ &= \int_{\alpha'}^{\beta'} \phi(x) \frac{dx}{dz} dz; \end{aligned}$$

thus 
$$\int_a^b \phi(x) dx = \int_{\alpha'}^{\beta'} \phi(x) \frac{dx}{dz} dz.$$

This result we may write simply thus

$$\int \phi(x) dx = \int \phi(x) \frac{dx}{dz} dz,$$

provided we remember that when the former integral is taken

between certain limits  $a$  and  $b$ , the latter must be taken between corresponding limits  $a'$  and  $b'$ .

11. As an example of the preceding article let  $\int \frac{dx}{x \sqrt{2ax - a^2}}$  be required. Assume  $x = \frac{a}{1-z}$ , thus

$$\frac{dx}{dz} = \frac{a}{(1-z)^2}, \text{ and } \int \frac{dx}{x \sqrt{2ax - a^2}} = \int \frac{1}{x \sqrt{2ax - a^2}} \frac{dx}{dz} dz$$

$$= \int \frac{dz}{a \sqrt{2(1-z) - (1-z)^2}} = \frac{1}{a} \int \frac{dz}{\sqrt{1-z^2}}$$

$$= \frac{1}{a} \sin^{-1} z = \frac{1}{a} \sin^{-1} \frac{x-a}{x}.$$

Here we have found the proposed integral by substituting for  $x$  in the manner indicated in the preceding article. This process will often simplify a proposed integral, but no rules can be given to guide the student as to the best assumption to make; this point must be left to observation and practice.

12. *Integration by parts.*

From the equation

$$\frac{d(uv)}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

we deduce by integrating both members,

$$uv = \int u \frac{dv}{dx} dx + \int v \frac{du}{dx} dx,$$

therefore 
$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx.$$

The use of this formula is called "integration by parts."

For example, consider  $\int x \cos ax dx$ . Since

$$\cos ax = \frac{1}{a} \frac{d \sin ax}{dx},$$

we may write the proposed expression in the form

$$\int \frac{x}{a} \frac{d. \sin ax}{dx} dx,$$

and this, by the formula, is equal to

$$\begin{aligned} & \frac{x \sin ax}{a} - \int \frac{\sin ax}{a} \frac{dx}{dx} \cdot dx \\ &= \frac{x \sin ax}{a} - \int \frac{\sin ax}{a} dx \\ &= \frac{x \sin ax}{a} + \frac{\cos ax}{a^2}. \end{aligned}$$

$$\begin{aligned} \text{Again, } \int x^2 \cos ax dx &= \int \frac{x^2}{a} \frac{d. \sin ax}{dx} dx \\ &= \frac{x^2 \sin ax}{a} - \int \frac{2x}{a} \sin ax dx \\ &= \frac{x^2 \sin ax}{a} + \int \frac{2x}{a^2} \frac{d. \cos ax}{dx} dx \\ &= \frac{x^2 \sin ax}{a} + \frac{2x \cos ax}{a^2} - \int \frac{2 \cos ax}{a^2} dx \\ &= \frac{x^2 \sin ax}{a} + \frac{2x \cos ax}{a^2} - \frac{2 \sin ax}{a^3}. \end{aligned}$$

$$\begin{aligned} \text{Again } \int e^{cx} \sin ax dx &= \int \frac{\sin ax}{c} \frac{d. e^{cx}}{dx} dx \\ &= \frac{\sin ax}{c} e^{cx} - \int \frac{ae^{cx} \cos ax}{c} dx \\ &= \frac{\sin ax}{c} e^{cx} - \int \frac{a \cos ax}{c^2} \frac{d. e^{cx}}{dx} dx \\ &= \frac{\sin ax}{c} e^{cx} - \frac{a \cos ax}{c^2} e^{cx} - \int \frac{a^2 \sin ax}{c^2} e^{cx} dx. \end{aligned}$$

By transposing,

$$\left(1 + \frac{a^2}{c^2}\right) \int e^{cx} \sin ax dx = \frac{e^{cx}}{c} \left(\sin ax - \frac{a}{c} \cos ax\right),$$

therefore 
$$\int e^{ax} \sin ax \, dx = \frac{e^{ax} (c \sin ax - a \cos ax)}{a^2 + c^2}.$$

Similarly we may shew that,

$$\int e^{ax} \cos ax \, dx = \frac{e^{ax} (c \cos ax + a \sin ax)}{a^2 + c^2}.$$

13. The differential coefficient of any function can always be found by the use of the rules given in the former part of the book, but it is not so with the integral of any assigned function. We know, for example, that if  $m$  be any number, positive or negative, except  $-1$ , then  $\int x^m \, dx = \frac{x^{m+1}}{m+1}$ , but when  $m = -1$  this is not true; in this case we have  $\int \frac{dx}{x} = \log x$ . If however we had not previously defined the term *logarithm*, and investigated the properties of a *logarithm*, we should have been unable to state what function would give  $\frac{1}{x}$  as its differential coefficient. Thus we may find ourselves limited in our powers of integration from our not having given a name to every particular function and investigated its properties.

In order to effect any proposed integration, it will often be necessary to use artifices which can only be suggested by practice.

14. We add a few miscellaneous examples. It should be noticed, that  $\phi_1(x)$  and  $\phi_2(x)$  being any functions of  $x$ ,

$$\int \{\phi_1(x) + \phi_2(x)\} \, dx = \int \phi_1(x) \, dx + \int \phi_2(x) \, dx,$$

or at least the two expressions which we assert to be equal can only differ by a constant; for if we differentiate both we arrive at the same result, namely  $\phi_1(x) + \phi_2(x)$ .

Ex. (1).  $\int \sqrt{a^2 - x^2} \, dx.$

$$\int \sqrt{a^2 - x^2} \, dx = x \sqrt{a^2 - x^2} + \int \frac{x^2 dx}{\sqrt{a^2 - x^2}}, \text{ by Art. 12,}$$

$$\text{and } \int \sqrt{a^2 - x^2} dx = \int \frac{a^2 - x^2}{\sqrt{a^2 - x^2}} dx = \int \frac{a^2 dx}{\sqrt{a^2 - x^2}} - \int \frac{x^2 dx}{\sqrt{a^2 - x^2}};$$

therefore, by addition,

$$2 \int \sqrt{a^2 - x^2} dx = x \sqrt{a^2 - x^2} + a^2 \int \frac{dx}{\sqrt{a^2 - x^2}},$$

$$\text{therefore } \int \sqrt{a^2 - x^2} dx = \frac{x \sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}. \text{ Art. 9.}$$

$$\text{Ex. (2). } \int \frac{dx}{\sqrt{x^2 + a^2}}.$$

Assume  $\sqrt{x^2 + a^2} = z - x,$   
therefore  $a^2 = z^2 - 2zx,$

$$\frac{dx}{dz} = \frac{z - x}{z}.$$

$$\text{Hence } \int \frac{dx}{\sqrt{x^2 + a^2}} = \int \frac{1}{\sqrt{x^2 + a^2}} \frac{dx}{dz} dz = \int \frac{dz}{z} = \log z \\ = \log \{x + \sqrt{x^2 + a^2}\}.$$

$$\text{Ex. (3). } \int \frac{dx}{\sqrt{x^2 - a^2}}.$$

As in Ex. (2), we may shew that the result is

$$\log \{x + \sqrt{x^2 - a^2}\}.$$

$$\text{Ex. (4). } \int \sqrt{x^2 + a^2} dx.$$

$$\int \sqrt{x^2 + a^2} dx = x \sqrt{x^2 + a^2} - \int \frac{x^2 dx}{\sqrt{x^2 + a^2}} \text{ by Art. 12.}$$

$$\text{Also } \int \sqrt{x^2 + a^2} dx = \int \frac{x^2 + a^2}{\sqrt{x^2 + a^2}} dx = \int \frac{x^2 dx}{\sqrt{x^2 + a^2}} + a^2 \int \frac{dx}{\sqrt{x^2 + a^2}};$$

therefore, by addition,

$$2 \int \sqrt{x^2 + a^2} dx = x \sqrt{x^2 + a^2} + a^2 \int \frac{dx}{\sqrt{x^2 + a^2}};$$

$$\text{therefore } \int \sqrt{x^2 + a^2} dx = \frac{x \sqrt{x^2 + a^2}}{2} + \frac{a^2}{2} \log \{x + \sqrt{x^2 + a^2}\}.$$

$$\begin{aligned} \text{Ex. (5.) } \int \frac{dx}{\sqrt{(a+bx+cx^2)}} & \\ \int \frac{dx}{\sqrt{(a+bx+cx^2)}} &= \frac{1}{\sqrt{c}} \int \frac{dx}{\sqrt{\left(\frac{a}{c} + \frac{bx}{c} + x^2\right)}} \\ &= \frac{1}{\sqrt{c}} \int \frac{dx}{\sqrt{\left\{\left(x + \frac{b}{2c}\right)^2 + \frac{4ac-b^2}{4c^2}\right\}}}. \end{aligned}$$

Putting  $x + \frac{b}{2c} = z$ , our integral becomes, by (2) and (3),

$$\frac{1}{\sqrt{c}} \log \{2cx + b + 2\sqrt{c} \sqrt{(a+bx+cx^2)}\},$$

where we omit the constant quantity  $\frac{1}{\sqrt{c}} \log 2c$ .

$$\begin{aligned} \text{Ex. (6.) } \int \frac{dx}{\sqrt{(a+bx-cx^2)}} & \\ \int \frac{dx}{\sqrt{(a+bx-cx^2)}} &= \frac{1}{\sqrt{c}} \int \frac{dx}{\sqrt{\left(\frac{a}{c} + \frac{bx}{c} - x^2\right)}} \\ &= \frac{1}{\sqrt{c}} \int \frac{dx}{\sqrt{\left\{\frac{4ac+b^2}{4c^2} - \left(x - \frac{b}{2c}\right)^2\right\}}}. \end{aligned}$$

Put  $h^2$  for  $\frac{4ac+b^2}{4c^2}$  and  $z$  for  $x - \frac{b}{2c}$ , thus the integral becomes  $\frac{1}{\sqrt{c}} \int \frac{dz}{\sqrt{(h^2-z^2)}}$ , which gives  $\frac{1}{\sqrt{c}} \sin^{-1} \frac{z}{h}$ , or

$$\frac{1}{\sqrt{c}} \sin^{-1} \frac{2cx-b}{\sqrt{(4ac+b^2)}}.$$

$$\text{Ex. (7.) } \int \frac{dx}{x \sqrt{(x^2-a^2)}}.$$

Put  $x = \frac{1}{y}$ , then  $\int \frac{dx}{x \sqrt{(x^2-a^2)}} = \int \frac{1}{x \sqrt{(x^2-a^2)}} \frac{dx}{dy} dy$



$$\begin{aligned}
 &= -\int \frac{dy}{\sqrt{(1-a^2y^2)}} = -\frac{1}{a} \int \frac{dy}{\sqrt{\left(\frac{1}{a^2}-y^2\right)}} = -\frac{1}{a} \sin^{-1} ay \\
 &= -\frac{1}{a} \sin^{-1} \frac{a}{x}.
 \end{aligned}$$

Since  $\sin^{-1} \frac{a}{x} + \cos^{-1} \frac{a}{x} = \frac{\pi}{2}$ , a constant, we may also write our last result thus

$$\int \frac{dx}{x\sqrt{(x^2-a^2)}} = \frac{1}{a} \cos^{-1} \frac{a}{x}.$$

Ex. (8).  $\int \frac{dx}{x\sqrt{(a^2 \pm x^2)}}.$

By putting  $x = \frac{1}{y}$ , as in Ex. (7), we deduce for the required result

$$\frac{1}{a} \log \frac{x}{a + \sqrt{(a^2 \pm x^2)}}.$$

Ex. (9).  $\int \frac{dx}{(x-a)^m}$  and  $\int \frac{dx}{x-a}.$

$$\int \frac{dx}{(x-a)^m} = -\frac{1}{m-1} \frac{1}{(x-a)^{m-1}},$$

$$\int \frac{dx}{x-a} = \log(x-a).$$

These are obvious if we differentiate the right-hand members.

Ex. (10).  $\int \frac{dx}{x^2-a^2}.$

$$\begin{aligned}
 \int \frac{dx}{x^2-a^2} &= \frac{1}{2a} \int \left( \frac{1}{x-a} - \frac{1}{x+a} \right) dx \\
 &= \frac{1}{2a} \int \frac{dx}{x-a} - \frac{1}{2a} \int \frac{dx}{x+a}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2a} \log(x-a) - \frac{1}{2a} \log(x+a) \\
 &= \frac{1}{2a} \log \frac{x-a}{x+a}.
 \end{aligned}$$

This supposes  $\frac{x-a}{x+a}$  positive; if  $\frac{x-a}{x+a}$  be negative, we must write

$$\int \frac{dx}{x^2-a^2} = \frac{1}{2a} \log \frac{a-x}{a+x}.$$

Ex. (11).  $\int \frac{dx}{a+bx+cx^2}$ .

$$\int \frac{dx}{a+bx+cx^2} = \frac{1}{c} \int \frac{dx}{\left(x + \frac{b}{2c}\right)^2 + \frac{4ac-b^2}{4c^2}}.$$

If  $\frac{4ac-b^2}{4c^2}$  be negative, we obtain the integral by Ex. (10), namely

$$\frac{1}{\sqrt{b^2-4ac}} \log \frac{2cx+b-\sqrt{b^2-4ac}}{2cx+b+\sqrt{b^2-4ac}}.$$

If  $\frac{4ac-b^2}{4c^2}$  be positive, then by Art. 9, the integral is

$$\frac{2}{\sqrt{4ac-b^2}} \tan^{-1} \frac{2cx+b}{\sqrt{4ac-b^2}}.$$

Ex. (12).  $\int \frac{Ax+B}{a+bx+cx^2} dx$ .

$$\begin{aligned}
 \int \frac{Ax+B}{a+bx+cx^2} dx &= \int \frac{Ax + \frac{Ab}{2c} + B - \frac{Ab}{2c}}{a+bx+cx^2} dx \\
 &= \frac{A}{2c} \int \frac{2cx+b}{a+bx+cx^2} dx + \left(B - \frac{Ab}{2c}\right) \int \frac{dx}{a+bx+cx^2}.
 \end{aligned}$$

The former integral is  $\frac{A}{2c} \log(a+bx+cx^2)$ , and the latter has been found in Ex. (11).

$$\begin{aligned} \text{Ex. (13). } \int \frac{dx}{\cos x} & \\ \int \frac{dx}{\cos x} &= \int \frac{\cos x \, dx}{\cos^2 x} = \int \frac{dz}{1-z^2}, \text{ if } z = \sin x, \\ &= \frac{1}{2} \log \frac{1+z}{1-z}, \text{ by Ex. (10),} \\ &= \frac{1}{2} \log \frac{1+\sin x}{1-\sin x} = \log \cot \left( \frac{\pi}{4} - \frac{x}{2} \right). \end{aligned}$$

$$\text{Similarly } \int \frac{dx}{\sin x} = \log \tan \frac{x}{2}.$$

$$\begin{aligned} \text{Ex. (14). } \int \frac{dx}{a+b \cos x} & \\ \int \frac{dx}{a+b \cos x} &= \int \frac{dx}{a \left( \sin^2 \frac{x}{2} + \cos^2 \frac{x}{2} \right) + b \left( \cos^2 \frac{x}{2} - \sin^2 \frac{x}{2} \right)} \\ &= \int \frac{\sec^2 \frac{x}{2} \, dx}{a+b + (a-b) \tan^2 \frac{x}{2}} \\ &= 2 \int \frac{dz}{a+b + (a-b) z^2}, \text{ if } z = \tan \frac{x}{2}. \end{aligned}$$

Hence, if  $a$  be greater than  $b$ , the integral is

$$\frac{2}{\sqrt{(a^2-b^2)}} \tan^{-1} \frac{z \sqrt{(a-b)}}{\sqrt{(a+b)}} \text{ or } \frac{2}{\sqrt{(a^2-b^2)}} \tan^{-1} \frac{\tan \frac{x}{2} \cdot \sqrt{(a-b)}}{\sqrt{(a+b)}};$$

and if  $a$  be less than  $b$ ,

$$\frac{1}{\sqrt{(b^2-a^2)}} \log \frac{z \sqrt{(b-a)} + \sqrt{(b+a)}}{z \sqrt{(b-a)} - \sqrt{(b+a)}},$$

$$\text{or } \frac{1}{\sqrt{(b^2-a^2)}} \log \frac{\sqrt{(b-a)} \tan \frac{x}{2} + \sqrt{(b+a)}}{\sqrt{(b-a)} \tan \frac{x}{2} - \sqrt{(b+a)}}.$$

In any of these examples, since we have found the *indefinite* integral, we can immediately ascertain the definite integral between any assigned limits. For example, since

$$\int \frac{dx}{\sqrt{(x^2 + a^2)}} = \log \{x + \sqrt{(x^2 + a^2)}\},$$

therefore

$$\begin{aligned} \int_a^{2a} \frac{dx}{\sqrt{(x^2 + a^2)}} &= \log [2a + \sqrt{\{(2a)^2 + a^2\}}] - \log \{a + \sqrt{(a^2 + a^2)}\} \\ &= \log \frac{2 + \sqrt{5}}{1 + \sqrt{2}}. \end{aligned}$$

15. The integral  $\int x^{m-1} (a + bx^n)^{\frac{p}{q}} dx$  can be found immediately in two cases. For assume

$$a + bx^n = t^q;$$

therefore  $x = \left(\frac{t^q - a}{b}\right)^{\frac{1}{n}},$

$$\frac{dx}{dt} = \frac{qt^{q-1}}{nb} \left(\frac{t^q - a}{b}\right)^{\frac{1}{n}-1}.$$

$$\begin{aligned} \text{Hence } \int x^{m-1} (a + bx^n)^{\frac{p}{q}} dx &= \int x^{m-1} (a + bx^n)^{\frac{p}{q}} \frac{dx}{dt} dt \\ &= \frac{q}{nb} \int t^{p+q-1} \left(\frac{t^q - a}{b}\right)^{\frac{m}{n}-1} dt. \end{aligned}$$

If  $\frac{m}{n}$  be a *positive integer* we can expand  $(t^q - a)^{\frac{m}{n}-1}$  in a finite series of powers of  $t$ , and each term of the product of this series by  $t^{p+q-1}$  will be immediately integrable.

$$\text{Again, } \int x^{m-1} (a + bx^n)^{\frac{p}{q}} dx = \int x^{\frac{pm}{q} + m-1} (ax^{-n} + b)^{\frac{p}{q}} dx;$$

and by the former case, if we put  $ax^{-n} + b = t^q$ , this is immediately integrable if

$$\frac{\frac{pn}{q} + m}{-n}$$

be a positive integer; that is, if  $\frac{p}{q} + \frac{m}{n}$  be a *negative integer*.

In the first case, if  $\frac{m}{n}$  were a *negative integer* the integral might still be found, as we shall see in the next chapter, and similarly, in the second case, if  $\frac{m}{n} + \frac{p}{q}$  were a *positive integer*: but as in these cases some further reductions are necessary, we do not say that the expressions are *immediately* integrable.

Ex. (1).  $\int x^2(a+x)^{\frac{1}{2}} dx.$

Here  $\frac{m}{n} = 3$ : assume  $a+x = t^2$ ; the integral becomes

$$2 \int (t^2 - a)^{\frac{1}{2}} t^2 dt \text{ or } 2 \int (t^6 - 2at^4 + a^2t^2) dt,$$

which gives

$$2 \left\{ \frac{t^7}{7} - \frac{2at^5}{5} + \frac{a^2t^3}{3} \right\};$$

thus  $\int x^2(a+x)^{\frac{1}{2}} dx = 2(a+x)^{\frac{3}{2}} \left\{ \frac{(a+x)^2}{7} - \frac{2a}{5}(a+x) + \frac{a^2}{3} \right\}.$

Ex. (2).  $\int \frac{dx}{x^2(1+x^2)^{\frac{1}{2}}}.$

Here  $m = -1$ ,  $n = 2$ ,  $\frac{p}{q} = -\frac{1}{2}$ ;

therefore  $\frac{m}{n} + \frac{p}{q} = -1.$

Assume  $x^{-2} + 1 = t^2$ ;

therefore  $x^2 = \frac{1}{t^2 - 1}.$

$$\frac{dx}{dt} = -\frac{t}{(t^2 - 1)^{\frac{3}{2}}}.$$

Also 
$$\int \frac{dx}{x^2(1+x^2)^{\frac{3}{2}}} = \int \frac{\frac{dx}{dt}}{x^2(x^{-2}+1)^{\frac{3}{2}}} dt.$$

Substitute for  $x$  and  $\frac{dx}{dt}$  their values, and this becomes  $-\int dt$ ,

which  $= -t$  or  $-\frac{\sqrt{(x^2+1)}}{x}$ .

Ex. (3). 
$$\int \frac{dx}{(a^2+x^2)^{\frac{3}{2}}}.$$

Here  $m = 1, n = 2, \frac{p}{q} = -\frac{3}{2},$

therefore  $\frac{m}{n} + \frac{p}{q} = -1.$

Assume  $a^2x^{-2} + 1 = t^2,$

therefore  $x^2 = \frac{a^2}{t^2-1},$

$$\frac{dx}{dt} = -\frac{at}{(t^2-1)^{\frac{3}{2}}},$$

$$\begin{aligned} \int \frac{dx}{(a^2+x^2)^{\frac{3}{2}}} &= \int \frac{\frac{dx}{dt}}{x^2(a^2x^{-2}+1)^{\frac{3}{2}}} dt = -\frac{1}{a^2} \int \frac{dt}{t^2} = \frac{1}{a^2 t} \\ &= \frac{x}{a^2 \sqrt{(a^2+x^2)}}. \end{aligned}$$

EXAMPLES.

1.  $\int \sqrt{(2ax-x^2)} dx = \frac{x-a}{2} \sqrt{(2ax-x^2)} + \frac{a^2}{2} \sin^{-1} \frac{x-a}{a}.$

2.  $\int \log x dx = x(\log x - 1).$

$$3. \int x^n \log x \, dx = \frac{x^{n+1}}{n+1} \left\{ \log x - \frac{1}{n+1} \right\}.$$

$$4. \int \theta \sin \theta \, d\theta = -\theta \cos \theta + \sin \theta.$$

$$5. \int \frac{dx}{e^x + e^{-x}} = \tan^{-1}(e^x).$$

$$6. \int \sqrt{\left(\frac{m+x}{x}\right)} dx = \sqrt{(mx+x^2)} + m \log \{\sqrt{x} + \sqrt{(m+x)}\}.$$

This may be found by putting  $x = z^2$ .

$$7. \int x \tan^{-1} x \, dx = \frac{1+x^2}{2} \tan^{-1} x - \frac{1}{2} x.$$

$$8. \int (1 - \cos x)^2 dx = \frac{3x}{2} - 2 \sin x + \frac{\sin 2x}{4}.$$

$$9. \int \frac{x dx}{(1-x)^3} = -\frac{1}{1-x} + \frac{1}{2(1-x)^2}.$$

$$10. \int \frac{x^2 dx}{a^3 - x^3} = \frac{1}{6a^3} \log \frac{a^3 + x^3}{a^3 - x^3}.$$

$$11. \int \frac{dx}{\sqrt{(1-3x-x^2)}} = \sin^{-1} \frac{3+2x}{\sqrt{13}}.$$

$$12. \int \frac{x dx}{\sqrt{(2ax-x^2)}} = -\sqrt{(2ax-x^2)} + a \operatorname{vers}^{-1} \frac{x}{a}.$$

$$13. \int \frac{dx}{\sqrt{(x^2-6x+13)}} = \log \{x-3 + \sqrt{(x^2-6x+13)}\}.$$

$$14. \int \frac{1 + \cos x}{x + \sin x} dx = \log (x + \sin x).$$

$$15. \int \frac{x + \sin x}{1 + \cos x} dx = x \tan \frac{x}{2}.$$

$$16. \int \frac{dx}{x (\log x)^n} = -\frac{1}{(n-1) (\log x)^{n-1}}.$$

17.  $\int \frac{\log(\log x)}{x} dx = \log x \cdot \log(\log x) - \log x.$
18.  $\int \frac{dx}{x + \sqrt{x^2 - 1}} = \frac{x^2}{2} - \frac{x\sqrt{x^2 - 1}}{2} + \frac{1}{2} \log \{x + \sqrt{x^2 - 1}\}.$
19.  $\int \frac{x^3 dx}{\sqrt{x-1}} = 2\sqrt{x-1} \left\{ \frac{(x-1)^3}{7} + \frac{2}{3}(x-1)^2 + x \right\}.$
20.  $\int e^{ax} \sin mx \cos nx dx = \frac{e^{ax}}{2} \frac{a \sin(m+n)x - (m+n) \cos(m+n)x}{a^2 + (m+n)^2}$   
 $+ \frac{e^{ax}}{2} \frac{a \sin(m-n)x - (m-n) \cos(m-n)x}{a^2 + (m-n)^2}.$
21.  $\int e^{-x} \cos^3 x dx = \frac{1}{4} \int e^{-x} (\cos 3x + 3 \cos x) dx$   
 $= \frac{e^{-x}}{40} (3 \sin 3x - \cos 3x) + \frac{3e^{-x}}{8} (\sin x - \cos x).$
22.  $\int_0^a \sqrt{a^2 - x^2} dx = \frac{\pi a^2}{4}.$
23.  $\int_0^{2a} \sqrt{2ax - x^2} dx = \frac{\pi a^2}{2}.$
24.  $\int_0^{2a} \text{vers}^{-1} \frac{x}{a} dx = \pi a.$
- Proceed thus—let  $\text{vers}^{-1} \frac{x}{a} = \theta$ , therefore  $x = a(1 - \cos \theta)$ ,  
 and the integral becomes  $\int_0^\pi a\theta \sin \theta d\theta.$
25.  $\int_0^{2a} x \text{vers}^{-1} \frac{x}{a} dx = \frac{5\pi a^3}{4}.$
26.  $\int_0^{2a} x^2 \text{vers}^{-1} \frac{x}{a} dx = \frac{11\pi a^3}{6}.$
27.  $\int_0^{2\pi} \sin^2 \theta \cos^2 \theta d\theta = \frac{2}{15}.$



$$28. \int \frac{dx}{\sin x + \cos x} = \frac{1}{\sqrt{2}} \log \tan \left( \frac{x}{2} + \frac{\pi}{8} \right).$$

$$29. \int \frac{dx}{x \sqrt{(a + bx + cx^2)}}.$$

Put  $x = \frac{1}{y}$  and this becomes a known form.

$$30. \int \frac{\sqrt{(1-x^2)} \sin^{-1} x dx}{x^4} = -\frac{\sin^{-1} x (1-x^2)^{\frac{3}{2}}}{3x^3} - \frac{1}{6x^2} - \frac{\log x}{3}.$$

This may be obtained by putting  $\sin^{-1} x = \theta$ .

$$31. \int \frac{\sin^{-1} x}{(1-x^2)^{\frac{3}{2}}} dx = \theta \tan \theta + \log \cos \theta, \text{ where } \sin \theta = x.$$

$$32. \int \frac{dx}{(a^2 - x^2)^{\frac{3}{2}}} = \frac{1}{a^4} \left( \cot \theta + \frac{\cot^3 \theta}{3} \right), \text{ where } x = a \cos \theta.$$

$$33. \int \frac{\sin^2 x dx}{a + b \cos^2 x} = \left( \frac{a+b}{ab^2} \right)^{\frac{1}{2}} \tan^{-1} \frac{\sqrt{a} \tan x}{\sqrt{a+b}} - \frac{x}{b}.$$

$$34. \int x^3 \sqrt{(a + bx^2)} dx = \left( \frac{a + bx^2}{5b^2} - \frac{a}{3b^2} \right) (a + bx^2)^{\frac{3}{2}}.$$

$$35. \int \frac{dx}{x^4 \sqrt{(1+x^2)}} = \frac{(2x^2-1) \sqrt{(1+x^2)}}{3x^3}.$$

$$36. \int \tan^{2n} \theta d\theta = \frac{x^{2n-1}}{2n-1} - \frac{x^{2n-3}}{2n-3} + \dots - (-1)^n x + (-1)^n \theta,$$

$x$  being  $= \tan \theta$ .

37. Shew that  $\int_0^\pi \sin mx \sin nx dx$  and  $\int_0^\pi \cos mx \cos nx dx$  are zero if  $m$  and  $n$  are *unequal integers*, and  $= \frac{\pi}{2}$  if  $m$  and  $n$  are *equal integers*.

$$38. \int \left\{ \log \left( \frac{x}{a} \right) \right\}^3 dx = x \left\{ \log \left( \frac{x}{a} \right) \right\}^3 - 3x \left\{ \log \frac{x}{a} \right\}^2 + 6x \log \frac{x}{a} - 6x.$$

$$39. \int \frac{\cot^{-1} x}{x^2(1+x^2)} dx = \frac{\theta^2}{2} - \theta \tan \theta - \log \cos \theta, \text{ where } \cot \theta = x.$$

$$40. \int \frac{2a+x}{a+x} \sqrt{\frac{a-x}{a+x}} dx = \sqrt{a^2-x^2} - \frac{2a\sqrt{a-x}}{\sqrt{a+x}}.$$

$$41. \int \frac{\text{vers}^{-1} \frac{x}{a}}{\sqrt{2ax-x^2}} dx = \frac{1}{2} \left( \text{vers}^{-1} \frac{x}{a} \right)^2.$$

$$42. \int_0^{\frac{\pi}{2}} \frac{dx}{1+c \cos x} = \frac{1}{\sqrt{1-c^2}} \cos^{-1} c, \text{ if } c \text{ is } < 1.$$

$$43. \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-\theta} \cos^3 \theta d\theta = \frac{3}{10} (e^{\frac{\pi}{2}} + e^{-\frac{\pi}{2}}).$$

$$44. \int \frac{(x^3-1) dx}{x \sqrt{1+3x^2+x^4}}. \text{ Assume } z = x + \frac{1}{x}.$$

$$45. \int \frac{(a+bx^n)^{\frac{3}{2}} dx}{x}. \text{ Assume } a+bx^n = z^4.$$

## CHAPTER II.

## RATIONAL FRACTIONS.

16. We proceed to the integration of such expressions as

$$\frac{A' + B'x + C'x^2 \dots + M'x^m}{A + Bx + Cx^2 \dots + Nx^n},$$

where  $A, B, \dots A', B', \dots$  are constants, so that both numerator and denominator are finite rational functions of  $x$ . If  $m$  be equal to  $n$ , or greater than  $n$ , we may by division reduce the preceding to the form of an integral function of  $x$ , and a fraction in which the numerator is of lower dimensions in  $x$  than the denominator. As the integral function of  $x$  can be integrated immediately, we may confine ourselves to the case of a fraction having its numerator at least one dimension lower than its denominator. In order to effect the integration we resolve the fraction into a series of more simple fractions called *partial fractions*, the possibility of which we proceed to demonstrate.

Let  $\frac{U}{V}$  be a rational fraction which is to be resolved into a series of partial fractions; suppose  $V$  a function of  $x$  of the  $n^{\text{th}}$  degree, and  $U$  a function of  $x$  of the  $(n-1)^{\text{th}}$  degree at most; we may without loss of generality suppose the coefficient of  $x^n$  in  $V$  to be unity. Suppose

$$V = (x-a)(x-b)^r(x^2 - 2ax + a^2 + \beta^2)(x^2 - 2\gamma x + \gamma^2 + \delta^2)^s,$$

so that the equation  $V=0$  has

- (1) one real root =  $a$ ,
- (2)  $r$  equal real roots, each =  $b$ ,
- (3) a pair of imaginary roots  $\alpha \pm \beta \sqrt{-1}$ ,
- (4)  $s$  pairs of imaginary roots, each being  $\gamma \pm \delta \sqrt{-1}$ .

By the theory of equations  $V$  must be the product of factors of the form we have supposed, the factors being more or fewer in number. Since  $V$  is of the  $n^{\text{th}}$  degree we have

$$1 + r + 2 + 2s = n.$$

Assume

$$\begin{aligned} \frac{U}{V} = & \frac{A}{x-a} + \frac{B_1}{(x-b)^r} + \frac{B_2}{(x-b)^{r-1}} + \frac{B_3}{(x-b)^{r-2}} \dots + \frac{B_r}{x-b} \\ & + \frac{Cx + D}{x^2 - 2ax + a^2 + \beta^2} \\ & + \frac{E_1x + F_1}{(x^2 - 2\gamma x + \gamma^2 + \delta^2)^s} + \frac{E_2x + F_2}{(x^2 - 2\gamma x + \gamma^2 + \delta^2)^{s-1}} \dots + \frac{E_sx + F_s}{x^2 - 2\gamma x + \gamma^2 + \delta^2}, \end{aligned}$$

where  $A, B_1, B_2, \dots, C, D, E_1, \dots$  are constants which, in order to justify our assumption, we must shew can be so determined as to make the second member of the above equation *identically* equal to the first. If we bring all the partial fractions to a common denominator and add them together, we have  $V$  for that common denominator, and for the numerator a function of  $x$  of the  $(n-1)^{\text{th}}$  degree. If we equate the coefficients of the different powers of  $x$  in this numerator with the corresponding coefficients in  $U$ , we shall have  $n$  equations of the first degree to determine the  $n$  quantities  $A, B_1, B_2, \dots$  and with these values of  $A, B_1, B_2, \dots$  the second member of the above equation becomes *identically equal* to the first, and thus  $\frac{U}{V}$  is decomposed into a series of partial fractions.

If  $V$  involves other single factors like  $x-a$ , each such factor will give rise to a fraction like  $\frac{A}{x-a}$ , and any repeated factor like  $(x-b)^r$  will give rise to a series of partial fractions of the form  $\frac{B_1}{(x-b)^r}, \frac{B_2}{(x-b)^{r-1}}, \&c.$  In like manner other factors of the form  $x^2 - 2ax + a^2 + \beta^2$  or  $(x^2 - 2\gamma x + \gamma^2 + \delta^2)^s$  will give rise to a fraction or a series of fractions respectively of the forms indicated above.

17. The demonstration given in Art. 16 is not very satisfactory, since we have not proved that the  $n$  equations of the

first degree which we use to determine  $A, B_1, B_2, \dots$  are *independent* and *consistent*.

A method of greater rigour has been given in a treatise on the Integral Calculus by Mr Homersham Cox, which we will here briefly indicate. Suppose  $F(x)$  to contain the factor  $x - a$  repeated  $n$  times; we have, if

$$F(x) = (x - a)^n \psi(x),$$

$$\frac{\phi(x)}{F(x)} = \frac{\phi(x)}{(x - a)^n \psi(x)} = \frac{\phi(x) - \frac{\phi(a)}{\psi(a)} \psi(x)}{(x - a)^n \psi(x)} + \frac{\frac{\phi(a)}{\psi(a)}}{(x - a)^n}.$$

Now  $\phi(x) - \frac{\phi(a)}{\psi(a)} \psi(x)$  vanishes when  $x = a$ , and is therefore divisible by  $x - a$ ; suppose the quotient denoted by  $\chi(x)$ , then

$$\frac{\phi(x)}{F(x)} = \frac{\chi(x)}{(x - a)^{n-1} \psi(x)} + \frac{\phi(a)}{\psi(a)} \frac{1}{(x - a)^n}.$$

The process may now be repeated on  $\frac{\chi(x)}{(x - a)^{n-1} \psi(x)}$ , and thus by successive operations the decomposition of  $\frac{\phi(x)}{F(x)}$  completely effected. In this proof  $a$  may be either a real root or an imaginary root of the equation  $F(x) = 0$ ; if  $a = \alpha + \beta \sqrt{-1}$ , then  $\alpha - \beta \sqrt{-1}$  will also be a root of  $F(x) = 0$ ; let  $b$  denote this root, then if we add the two partial fractions

$$\frac{\phi(a)}{\psi(a)} \frac{1}{(x - a)^n} \text{ and } \frac{\phi(b)}{\psi(b)} \frac{1}{(x - b)^n},$$

we shall obtain a result free from  $\sqrt{-1}$ .

18. With respect to the integration of these partial fractions we refer to Examples (9—12) of Art. 14 for all the forms except  $\frac{Lx + M}{(x^2 - 2\gamma x + \gamma^2 + \delta^2)^m}$ , and this will be given hereafter.

Having proved that a rational fraction can be decomposed in the manner assumed in Art. 16, we may make use of

different algebraical artifices in order to diminish the labour of determining  $A, B_1, B_2, \&c.$  The most useful consideration is, that since the numerator of the proposed fraction is *identically* equal to the numerator formed by adding together the partial fractions, if we assign *any* value to the variable  $x$  the equality still subsists.

19. *To determine the partial fraction corresponding to a single factor of the first degree.*

Suppose  $\frac{\phi(x)}{F(x)}$  represents a fraction to be decomposed, and let  $F(x)$  contain the factor  $x - a$  once; assume

$$\frac{\phi(x)}{F(x)} = \frac{A}{x - a} + \frac{\chi(x)}{\psi(x)} \dots\dots\dots(1),$$

where  $A$  is a constant, and  $\frac{\chi(x)}{\psi(x)}$  represents the sum of all the partial fractions exclusive of  $\frac{A}{x - a}$ , and  $F(x) = (x - a)\psi(x)$ .

From (1)

$$\phi(x) = A\psi(x) + (x - a)\chi(x) \dots\dots\dots(2).$$

In (2), which holds for any value of  $x$ , make  $x = a$ , then

$$\phi(a) = A\psi(a),$$

therefore 
$$A = \frac{\phi(a)}{\psi(a)}.$$

Since  $F'(x) = \psi(x) + (x - a)\psi'(x)$ , we have

$$F'(a) = \psi(a),$$

therefore 
$$A = \frac{\phi(a)}{F'(a)}.$$

20. *To determine the partial fractions corresponding to a factor of the first degree which is repeated.*

Suppose  $F(x)$  contains a factor  $x - a$  repeated  $n$  times, and let  $F(x) = (x - a)^n \psi(x)$ . Assume

$$\frac{\phi(x)}{F(x)} = \frac{A_1}{(x-a)^n} + \frac{A_2}{(x-a)^{n-1}} + \frac{A_3}{(x-a)^{n-2}} \dots + \frac{A_n}{x-a} + \frac{\chi(x)}{\psi(x)},$$

where  $\frac{\chi(x)}{\psi(x)}$  denotes the sum of the partial fractions arising from the other factors of  $F(x)$ . Multiply both sides of the equation by  $(x-a)^n$  and put  $f(x)$  for  $\frac{\phi(x)}{F(x)}(x-a)^n$ ; thus

$$f(x) = A_1 + A_2(x-a) + A_3(x-a)^2 \dots + A_n(x-a)^{n-1} + \frac{\chi(x)}{\psi(x)}(x-a)^n.$$

Differentiate successively both members of this identity and put  $x = a$  after differentiation; then

$$\begin{aligned} f(a) &= A_1, \\ f'(a) &= A_2, \\ f''(a) &= 1.2A_3, \\ f'''(a) &= \underline{3}A_4, \\ &\dots\dots\dots \\ f^{n-1}(a) &= \underline{n-1}A_n. \end{aligned}$$

Thus  $A_1, A_2, \dots, A_n$  are determined.

21. To determine the partial fractions corresponding to a pair of imaginary roots which do not recur.

Let  $\frac{\phi(x)}{F(x)}$  denote the fraction to be decomposed; and  $\alpha \pm \beta \sqrt{-1}$  a pair of imaginary roots; then if we denote these roots by  $a$  and  $b$  and proceed as in Art. 19, we have for the partial fractions

$$\frac{\phi(a)}{F''(a)} \frac{1}{x-a} \quad \text{and} \quad \frac{\phi(b)}{F''(b)} \frac{1}{x-b}.$$

Suppose  $\frac{\phi(a)}{F''(a)} = A - B \sqrt{-1}$ ; then since  $\frac{\phi(b)}{F''(b)}$  may be obtained from  $\frac{\phi(a)}{F''(a)}$  by changing the sign of  $\sqrt{-1}$ , we must have  $\frac{\phi(b)}{F''(b)} = A + B \sqrt{-1}$ . Hence the fractions are

$$\frac{A - B\sqrt{-1}}{x - \alpha - \beta\sqrt{-1}} \text{ and } \frac{A + B\sqrt{-1}}{x - \alpha + \beta\sqrt{-1}};$$

and their sum is

$$\frac{2A(x - \alpha) + 2B\beta}{(x - \alpha)^2 + \beta^2}.$$

22. Or we may proceed thus. Suppose  $x^2 - px + q$  to denote the quadratic factor which gives rise to the pair of imaginary roots  $\alpha \pm \beta\sqrt{-1}$ ; then assume

$$\frac{\phi(x)}{F(x)} = \frac{Lx + M}{x^2 - px + q} + \frac{\chi(x)}{\psi(x)},$$

so that  $F(x) = (x^2 - px + q)\psi(x)$ . Multiply by  $F(x)$ ; thus

$$\phi(x) = (Lx + M)\psi(x) + (x^2 - px + q)\chi(x) \dots \dots (1).$$

Now ascribe to  $x$  either of the values which makes  $x^2 - px + q$  vanish; then (1) reduces to

$$\phi(x) = (Lx + M)\psi(x) \dots \dots \dots (2).$$

Now by the repeated substitution of  $px - q$  for  $x^2$  in both members of (2), we shall at last have  $x$  occurring in the first power only, so that the equation takes the form

$$Px + Q = P'x + Q'.$$

Now put for  $x$  its value  $\alpha + \beta\sqrt{-1}$  and equate the coefficients of the impossible parts; thus

$$P = P' \text{ and therefore also } Q = Q'.$$

Here  $P$  and  $Q$  are known quantities, and  $P'$  and  $Q'$  involve the unknown quantities  $L$  and  $M$  to the first power only, so that we have two equations of the first degree for finding  $L$  and  $M$ .

23. *To determine the partial fractions corresponding to a pair of imaginary roots which is repeated.*

We may proceed as in Art. 20. Or we may adopt the following method. Suppose  $x^2 - px + q$  to be the quadratic factor which occurs  $r$  times; assume



$$\frac{\phi(x)}{F(x)} = \frac{L_r x + M_r}{(x^2 - px + q)^r} + \frac{L_{r-1} x + M_{r-1}}{(x^2 - px + q)^{r-1}} + \dots + \frac{L_1 x + M_1}{x^2 - px + q} + \frac{\chi(x)}{\psi(x)},$$

so that  $F(x) = (x^2 - px + q)^r \psi(x)$ . Multiply by  $F(x)$ ; thus

$$\begin{aligned} \phi(x) &= (L_r x + M_r) \psi(x) + (L_{r-1} x + M_{r-1}) (x^2 - px + q) \psi(x) \\ &\quad + \dots + (x^2 - px + q)^r \chi(x) \dots \dots \dots (1). \end{aligned}$$

Now ascribe to  $x$  either of the values which makes  $x^2 - px + q$  vanish; thus the equation reduces to

$$\phi(x) = (L_r x + M_r) \psi(x).$$

Proceed as in Art. 22, and thus find  $L_r$  and  $M_r$ . Then from (1) by transposition we have

$$\phi(x) - (L_r x + M_r) \psi(x) = (L_{r-1} x + M_{r-1}) (x^2 - px + q) \psi(x) + \dots$$

The right hand member has  $x^2 - px + q$  for a factor of every term; hence as the two members are *identical* we can divide by this factor. Let  $\phi_1(x)$  indicate the quotient obtained on the left; then

$$\begin{aligned} \phi_1(x) &= (L_{r-1} x + M_{r-1}) \psi(x) + (L_{r-2} x + M_{r-2}) (x^2 - px + q) \psi(x) \\ &\quad + \dots + (x^2 - px + q)^{r-1} \chi(x) \dots \dots \dots (2). \end{aligned}$$

From (2) we find  $L_{r-1}$  and  $M_{r-1}$  as before; then by transposition and division

$$\phi_2(x) = (L_{r-2} x + M_{r-2}) \psi(x) + (L_{r-3} x + M_{r-3}) (x^2 - px + q) \psi(x) + \dots$$

and so on until all the quantities are determined.

24. Take for example  $\frac{x^2 - 3x - 2}{(x^2 + x + 1)^2 (x + 1)^2}$ . Assume it equal to

$$\frac{L_2 x + M_2}{(x^2 + x + 1)^2} + \frac{L_1 x + M_1}{x^2 + x + 1} + \frac{\chi(x)}{(x + 1)^2};$$

then  $x^2 - 3x - 2 = (L_2 x + M_2) (x + 1)^2$

$$+ (L_1 x + M_1) (x^2 + x + 1) (x + 1) + (x^2 + x + 1)^2 \chi(x) \dots \dots (1).$$

Suppose  $x^2 + x + 1 = 0$ ; thus the equation reduces to

$$\begin{aligned} x^2 - 3x - 2 &= (L_2x + M_2)(x + 1)^2 \\ &= (L_2x + M_2)(x^2 + 2x + 1). \end{aligned}$$

Put  $-x - 1$  for  $x^2$ ; thus

$$\begin{aligned} -4x - 3 &= (L_2x + M_2)x = L_2x^2 + M_2x \\ &= -L_2(x + 1) + M_2x; \end{aligned}$$

therefore  $-4 = -L_2 + M_2$ , and  $-3 = -L_2$ ;

thus  $L_2 = 3$ , and  $M_2 = -1$ .

From (1) by transposition

$$\begin{aligned} x^2 - 3x - 2 - (3x - 1)(x + 1)^2 \\ = (L_1x + M_1)(x^2 + x + 1)(x + 1)^2 + (x^2 + x + 1)^2 \chi(x). \end{aligned}$$

The left hand member is  $-3x^3 - 4x^2 - 4x - 1$ ; divide by  $x^2 + x + 1$ ; thus

$$-(3x + 1) = (L_1x + M_1)(x + 1)^2 + (x^2 + x + 1)\chi(x) \dots (2).$$

Again, suppose  $x^2 + x + 1 = 0$ ; thus

$$\begin{aligned} -3x - 1 &= (L_1x + M_1)(x^2 + 2x + 1) = (L_1x + M_1)x \\ &= -L_1(x + 1) + M_1x; \end{aligned}$$

therefore  $-3 = -L_1 + M_1$ , and  $-1 = -L_1$ ;

thus  $L_1 = 1$  and  $M_1 = -2$ .

Thus the partial fractions corresponding to the quadratic factor are found. The partial fractions corresponding to the factor  $(x + 1)^2$  may then be found by Art. 20. Or we may from (2) by transposition and division by  $x^2 + x + 1$  obtain

$$-(x - 1) = \chi(x).$$

Thus

$$\frac{\chi(x)}{(x + 1)^2} = -\frac{x - 1}{(x + 1)^2} = -\frac{x + 1}{(x + 1)^2} + \frac{2}{(x + 1)^2} = -\frac{1}{x + 1} + \frac{2}{(x + 1)^2};$$

therefore

$$\frac{x^2 - 3x - 2}{(x^2 + x + 1)^2(x + 1)^2} = \frac{3x - 1}{(x^2 + x + 1)^2} + \frac{x - 2}{x^2 + x + 1} + \frac{2}{(x + 1)^2} - \frac{1}{x + 1}.$$

25. Examples. Required the integral of  $\frac{5x^3+1}{x^2-3x+2}$ .

By division we have

$$\frac{5x^3+1}{x^2-3x+2} = 5x + 15 + \frac{35x-29}{x^2-3x+2}.$$

Assume 
$$\frac{35x-29}{x^2-3x+2} = \frac{A}{x-1} + \frac{B}{x-2};$$

therefore 
$$35x-29 = A(x-2) + B(x-1).$$

Make  $x$  successively equal to 1 and 2; then

$$35-29 = -A, \text{ or } A = -6,$$

$$70-29 = B, \text{ or } B = 41;$$

therefore 
$$\frac{5x^3+1}{x^2-3x+2} = 5x + 15 - \frac{6}{x-1} + \frac{41}{x-2};$$

therefore 
$$\int \frac{5x^3+1}{x^2-3x+2} dx = \frac{5x^2}{2} + 15x - 6 \log(x-1) + 41 \log(x-2).$$

Required the integral of  $\frac{9x^2+9x-128}{x^3-5x^2+3x+9}$ .

Since  $x^3-5x^2+3x+9 = (x-3)^2(x+1)$ , we assume

$$\frac{9x^2+9x-128}{x^3-5x^2+3x+9} = \frac{A}{x+1} + \frac{B_1}{(x-3)^2} + \frac{B_2}{x-3};$$

therefore 
$$9x^2+9x-128 = A(x-3)^2 + B_1(x+1) + B_2(x+1)(x-3).$$

Make  $x = 3$  and  $-1$  successively, and we find

$$B_1 = -5, \quad A = -8.$$

Also by equating the coefficients of  $x^2$ , we have

$$9 = A + B_2,$$

therefore 
$$B_2 = 17;$$

therefore

$$\int \frac{9x^2+9x-128}{x^3-5x^2+3x+9} dx = -8 \log(x+1) + \frac{5}{x-3} + 17 \log(x-3).$$

Required the integral of  $\frac{x^2 + 1}{(x - 1)^4(x^2 + 1)}$ .

Assume  $\frac{x^2 + 1}{(x - 1)^4(x^2 + 1)}$

$$= \frac{A_1}{(x - 1)^4} + \frac{A_2}{(x - 1)^3} + \frac{A_3}{(x - 1)^2} + \frac{A_4}{x - 1} + \frac{B}{x + 1} + \frac{Cx + D}{x^2 - x + 1};$$

therefore  $x^2 + 1 = \{A_1 + A_2(x - 1) + A_3(x - 1)^2 + A_4(x - 1)^3\}(x^2 + 1)$   
 $+ \{B(x^2 - x + 1) + (Cx + D)(x + 1)\}(x - 1)^4 \dots (1).$

Put  $x = 1$ , then  $2 = 2A_1 \dots \dots \dots (2);$

therefore  $A_1 = 1.$

From (1) and (2) we have by subtraction,

$$x^2 - 1 = A_1(x^2 - 1) + \{A_2 + A_3(x - 1) + A_4(x - 1)^2\}(x - 1)(x^2 + 1)$$

$$+ \{B(x^2 - x + 1) + (Cx + D)(x + 1)\}(x - 1)^4.$$

Divide by  $x - 1$ , then

$$x + 1 = A_1(x^2 + x + 1) + \{A_2 + A_3(x - 1) + A_4(x - 1)^2\}(x^2 + 1)$$

$$+ \{B(x^2 - x + 1) + (Cx + D)(x + 1)\}(x - 1)^3 \dots (3).$$

Put  $x = 1$ , then  $2 = 3A_1 + 2A_2 \dots \dots \dots (4);$

therefore  $A_2 = -\frac{1}{2}.$

From (3) and (4), by subtraction,

$$x - 1 = A_1(x^2 + x - 2) + A_2(x^2 - 1) + \{A_3 + A_4(x - 1)\}(x - 1)(x^2 + 1)$$

$$+ \{B(x^2 - x + 1) + (Cx + D)(x + 1)\}(x - 1)^3.$$

Divide by  $x - 1$ , then

$$1 = A_1(x + 2) + A_2(x^2 + x + 1) + \{A_3 + A_4(x - 1)\}(x^2 + 1)$$

$$+ \{B(x^2 - x + 1) + (Cx + D)(x + 1)\}(x - 1)^2 \dots (5).$$

Put  $x = 1$ , then  $1 = 3A_1 + 3A_2 + 2A_3 \dots \dots \dots (6)$ ;  
therefore  $A_3 = -\frac{1}{4}$ .

From (5) and (6), by subtraction,  
 $0 = A_1(x-1) + A_2(x^2+x-2) + A_3(x^2-1) + A_4(x-1)(x^2+1)$   
 $+ \{B(x^2-x+1) + (Cx+D)(x+1)\}(x-1)^2$ .

Divide by  $x-1$ , then  
 $0 = A_1 + A_2(x+2) + A_3(x^2+x+1) + A_4(x^2+1)$   
 $+ \{B(x^2-x+1) + (Cx+D)(x+1)\}(x-1) \dots \dots (7)$ .

Put  $x = 1$ , then  $0 = A_1 + 3A_2 + 3A_3 + 2A_4 \dots \dots \dots (8)$ .  
therefore  $A_4 = \frac{5}{8}$ .

From (7) and (8), by subtraction,  
 $0 = A_2(x-1) + A_3(x^2+x-2) + A_4(x^2-1)$   
 $+ \{B(x^2-x+1) + (Cx+D)(x+1)\}(x-1)$ .

Divide by  $x-1$ , then  
 $0 = A_2 + A_3(x+2) + A_4(x^2+x+1)$   
 $+ B(x^2-x+1) + (Cx+D)(x+1) \dots \dots (9)$ .

Put  $x = -1$ , then  
 $0 = A_2 + A_3 + A_4 + 3B \dots \dots \dots (10)$ ;  
therefore  $B = \frac{1}{24}$ .

From (9) and (10), by subtraction,  
 $0 = A_3(x+1) + A_4(x^2+x) + B(x^2-x-2) + (Cx+D)(x+1)$ .

Divide by  $x+1$ , then  
 $0 = A_3 + A_4x + B(x-2) + Cx + D \dots \dots (11)$ .

Put  $x = 0$ , then  
 $A_3 - 2B + D = 0 \dots \dots \dots (12)$ ;  
therefore  $D = \frac{1}{8}$ .

From (11) and (12), by subtraction

$$A_4 + B + C = 0;$$

therefore  $C = -\frac{2}{3}$ ;

$$\text{therefore } \frac{x^2 + 1}{(x-1)^4(x^2+1)} = \frac{1}{(x-1)^4} - \frac{1}{2(x-1)^3} - \frac{1}{4(x-1)^2} \\ + \frac{5}{8(x-1)} + \frac{1}{24(x+1)} - \frac{2x-1}{3(x^2-x+1)};$$

$$\text{therefore } \int \frac{(x^2+1) dx}{(x-1)^4(x^2+1)} = -\frac{1}{3(x-1)^3} + \frac{1}{4(x-1)^2} + \frac{1}{4(x-1)} \\ + \frac{5}{8} \log(x-1) + \frac{1}{24} \log(x+1) - \frac{1}{3} \log(x^2-x+1).$$

26. We will give as additional examples the integration of  $\frac{x^{m-1}}{x^n \pm 1}$ , supposing  $m$  and  $n$  positive integers, and  $m-1$  less than  $n$ .

Required the integral of  $\frac{x^{m-1}}{x^n - 1}$ ,  $n$  being supposed even.

By the theory of equations the real roots of  $x^n - 1 = 0$  are 1 and  $-1$ , and the imaginary roots are found from the expression  $\cos r\theta \pm \sqrt{-1} \sin r\theta$ , where  $\theta = \frac{\pi}{n}$ , and  $r$  takes in succession the values 2, 4, ... up to  $n-2$ . Now by Art. 19 if  $\frac{\phi(x)}{F(x)}$  be the fraction to be decomposed, the partial fraction corresponding to the root  $a$  is  $\frac{\phi(a)}{F'(a)} \frac{1}{x-a}$ . In the present case

$$\frac{\phi(a)}{F'(a)} = \frac{a^{m-1}}{na^{n-1}} = \frac{a^m}{na^n} = \frac{a^m}{n}, \text{ since } a^n = 1.$$

Hence corresponding to the root 1 we have the partial fraction  $\frac{1}{n(x-1)}$ , and corresponding to the root  $-1$  we have the partial fraction  $\frac{(-1)^m}{n(x+1)}$ . And corresponding to the pair of roots

$$\cos r\theta \pm \sqrt{-1} \sin r\theta$$

we have

$$\frac{\{\cos r\theta + \sqrt{(-1) \sin r\theta}\}^m}{n \{x - \cos r\theta - \sqrt{(-1) \sin r\theta}\}} + \frac{\{\cos r\theta - \sqrt{(-1) \sin r\theta}\}^m}{n \{x - \cos r\theta + \sqrt{(-1) \sin r\theta}\}},$$

that is

$$\frac{\cos mr\theta + \sqrt{(-1) \sin mr\theta}}{n \{x - \cos r\theta - \sqrt{(-1) \sin r\theta}\}} + \frac{\cos mr\theta - \sqrt{(-1) \sin mr\theta}}{n \{x - \cos r\theta + \sqrt{(-1) \sin r\theta}\}},$$

that is 
$$\frac{2 \cos mr\theta (x - \cos r\theta) - 2 \sin mr\theta \sin r\theta}{n (x^2 - 2x \cos r\theta + 1)}.$$

Thus 
$$\frac{x^{m-1}}{x^n - 1} = \frac{1}{n(x-1)} + \frac{(-1)^m}{n(x+1)} + \frac{2}{n} \sum \frac{\cos mr\theta (x - \cos r\theta) - \sin mr\theta \sin r\theta}{(x - \cos r\theta)^2 + \sin^2 r\theta},$$

where  $\Sigma$  indicates a sum to be formed by giving to  $r$  all the even integral values from 2 to  $n-2$  inclusive. Hence

$$\int \frac{x^{m-1} dx}{x^n - 1} = \frac{1}{n} \log(x-1) + \frac{(-1)^m}{n} \log(x+1) + \frac{1}{n} \sum \cos mr\theta \log(x^2 - 2x \cos r\theta + 1) - \frac{2}{n} \sum \sin mr\theta \tan^{-1} \frac{x - \cos r\theta}{\sin r\theta}.$$

27. Required the integral of  $\frac{x^{m-1}}{x^n - 1}$ ,  $n$  being supposed odd.

The real root of  $x^n - 1 = 0$  is 1, and the imaginary roots are found from the expression  $\cos r\theta \pm \sqrt{(-1) \sin r\theta}$ , where  $\theta = \frac{\pi}{n}$ , and  $r$  takes in succession the values 2, 4, ... up to  $n-1$ . Hence as before we shall find

$$\int \frac{x^{m-1} dx}{x^n - 1} = \frac{1}{n} \log(x-1) + \frac{1}{n} \sum \cos mr\theta \log(x^2 - 2x \cos r\theta + 1) - \frac{2}{n} \sum \sin mr\theta \tan^{-1} \frac{x - \cos r\theta}{\sin r\theta}.$$

28. Required the integral of  $\frac{x^{m-1}}{x^n+1}$ ,  $n$  being supposed even.

The equation  $x^n + 1 = 0$  has now no real root; the imaginary roots are found from the expression  $\cos r\theta \pm \sqrt{-1} \sin r\theta$ , where  $\theta = \frac{\pi}{n}$ , and  $r$  takes in succession the values 1, 3, ... up to  $n - 1$ . And if  $a$  be a root of  $x^n + 1 = 0$ , we have

$$\frac{\phi(a)}{F'(a)} = \frac{a^{m-1}}{na^{n-1}} = \frac{a^m}{na^n} = -\frac{a^m}{n};$$

thus the sum of the two fractions corresponding to a pair of imaginary roots is

$$-\frac{2 \cos mr\theta (x - \cos r\theta) - \sin mr\theta \sin r\theta}{n (x - \cos r\theta)^2 + \sin^2 r\theta}.$$

Hence

$$\int \frac{x^{m-1} dx}{x^n + 1} = -\frac{1}{n} \sum \cos mr\theta \log (x^2 - 2x \cos r\theta + 1) + \frac{2}{n} \sum \sin mr\theta \tan^{-1} \frac{x - \cos r\theta}{\sin r\theta},$$

where  $\sum$  indicates a sum to be formed by giving to  $r$  all the odd integral values from 1 to  $n - 1$  inclusive.

29. Required the integral of  $\frac{x^{m-1}}{x^n+1}$ ,  $n$  being supposed odd.

The real root of  $x^n + 1 = 0$  is in this case  $-1$ , and the imaginary roots are found from the expression  $\cos r\theta \pm \sqrt{-1} \sin r\theta$ , where  $\theta = \frac{\pi}{n}$  and  $r$  takes in succession the values 1, 3, ... up to  $n - 2$ . Hence we shall obtain

$$\int \frac{x^{m-1} dx}{x^n + 1} = \frac{(-1)^{m-1}}{n} \log (x + 1)$$

$$-\frac{1}{n} \sum \cos mr\theta \log (x^2 - 2x \cos r\theta + 1) + \frac{2}{n} \sum \sin mr\theta \tan^{-1} \frac{x - \cos r\theta}{\sin r\theta}.$$



## EXAMPLES.

1.  $\int \frac{dx}{x^3-1} = \frac{1}{6} \log \frac{(x-1)^2}{x^2+x+1} - \frac{1}{\sqrt{3}} \tan^{-1} \frac{2x+1}{\sqrt{3}}.$
2.  $\int \frac{x^2-1}{x^2-4} dx = x + \log \left( \frac{x-2}{x+2} \right)^{\frac{3}{2}}.$
3.  $\int \frac{x^3 dx}{x^3+7x+12} = \frac{x^3}{2} - 7x + 64 \log(x+4) - 27 \log(x+)$
4.  $\int \frac{dx}{a^4-x^4} = \frac{1}{2a^3} \tan^{-1} \frac{x}{a} + \frac{1}{4a^3} \log \frac{a+x}{a-x}.$
5.  $\int \frac{2x^2-3a^2}{x^4-a^4} dx = \frac{5}{2a} \tan^{-1} \frac{x}{a} - \frac{1}{4a} \log \frac{x-a}{x+a}.$
6.  $\int \frac{dx}{(x^2+1)(x^2+x+1)} = \frac{1}{2} \log \frac{x^2+x+1}{x^2+1} + \frac{1}{\sqrt{3}} \tan^{-1} \frac{2x+}{\sqrt{3}}$
7.  $\int \frac{x^2 dx}{x^4+x^2-2} = \frac{1}{6} \log \frac{x-1}{x+1} + \frac{\sqrt{2}}{3} \tan^{-1} \frac{x}{\sqrt{2}}.$
8.  $\int \frac{x^2-1}{x^4+x^2+1} dx = \frac{1}{2} \log \frac{x^2-x+1}{x^2+x+1}.$
9.  $\int \frac{x^2-3x+3}{(x-1)(x-2)} dx = x + \log \frac{x-2}{x-1}.$
10.  $\int \frac{(3x-1) dx}{x^3-x^2-2x} = \frac{1}{2} \log x + \frac{5}{6} \log(x-2) - \frac{4}{3} \log(x+1).$
11.  $\int \frac{dx}{(x^2+a^2)(x+b)} = \frac{1}{b^2+a^2} \left\{ \log \frac{x+b}{\sqrt{x^2+a^2}} + \frac{b}{a} \tan^{-1} \frac{x}{a} \right\}.$
12.  $\int \frac{dx}{x(1+x+x^2+x^3)} = \log x - \frac{1}{2} \log(1+x) - \frac{1}{4} \log(1+x)$   
 $- \frac{1}{2} \tan^{-1}$

$$13. \int \frac{dx}{(x-1)^2(x^2+1)^2} = -\frac{1}{4(x-1)} - \frac{1}{2} \log(x-1) \\ + \frac{1}{4} \tan^{-1} x - \frac{1}{4(x^2+1)} + \frac{1}{4} \log(x^2+1).$$

$$14. \int \frac{x dx}{(1+x)(1+2x)^2(1+x^2)} = \frac{2}{5} \frac{1}{1+2x} - \frac{1}{2} \log(1+x) \\ - \frac{7}{100} \log(1+x^2) + \frac{16}{25} \log(1+2x) + \frac{1}{50} \tan^{-1} x.$$

$$15. \int \frac{x^2 dx}{x^4+1} = \frac{1}{4\sqrt{2}} \log \frac{x^2 - x\sqrt{2} + 1}{x^2 + x\sqrt{2} + 1} \\ + \frac{1}{2\sqrt{2}} \{ \tan^{-1}(x\sqrt{2} + 1) + \tan^{-1}(x\sqrt{2} - 1) \}.$$

$$16. \int \frac{x^2 dx}{x^6+1} = \frac{1}{12} \log(x^4 - x^2 + 1) - \frac{1}{6} \log(x^2 + 1) \\ + \frac{1}{2\sqrt{3}} \{ \tan^{-1}(2x - \sqrt{3}) - \tan^{-1}(2x + \sqrt{3}) \}.$$

$$17. \int \frac{dy}{\sqrt[3]{1-y^3}}. \quad \text{Assume } 1-y^3 = y^3 z^3.$$

$$18. \int \frac{dx}{(1+x)\sqrt[3]{1+3x+3x^2}}. \quad \text{Assume } y = \frac{x}{1+x}.$$

## CHAPTER III.

## FORMULÆ OF REDUCTION.

30. LET  $a + bx^n$  be denoted by  $X$ ; by integration by parts we have

$$\begin{aligned} \int x^{m-1} X^p dx &= \frac{X^p x^m}{m} - \int \frac{x^m}{m} p X^{p-1} \frac{dX}{dx} dx \\ &= \frac{X^p x^m}{m} - \frac{bnp}{m} \int x^{m+n-1} X^{p-1} dx \dots \dots \dots (1). \end{aligned}$$

The equation (1) is called a *formula of reduction*; by means of it we make the integral of  $x^{m-1} X^p$  depend on that of  $x^{m+n-1} X^{p-1}$ . In the same way the latter integral can be made to depend on that of  $x^{m+2n-1} X^{p-2}$ ; and thus, if  $p$  be an integer we may proceed until we arrive at  $x^{m+n(p-1)} X^{p-p}$ , that is  $x^{m+n(p-1)}$ , which is immediately integrable.

From (1), by transposition,

$$\int x^{m+n-1} X^{p-1} dx = \frac{x^m X^p}{bnp} - \frac{m}{bnp} \int x^{m-1} X^p dx.$$

Change  $m$  into  $m-n$  and  $p$  into  $p+1$ ; thus

$$\int x^{m-1} X^p dx = \frac{x^{m-n} X^{p+1}}{bn(p+1)} - \frac{m-n}{bn(p+1)} \int x^{m-n-1} X^{p+1} dx \dots (2).$$

This formula may be used when we wish to make the integral of  $x^m X^p$  depend upon another in which the exponent of  $x$  is diminished and that of  $X$  increased. For example, if  $m=3$ ,  $n=2$ , and  $p=-\frac{3}{2}$ , we have

$$\int \frac{x^3 dx}{(a+bx^2)^{\frac{3}{2}}} = -\frac{x}{b\sqrt{(a+bx^2)}} + \frac{1}{b} \int \frac{dx}{\sqrt{(a+bx^2)}}.$$

The latter integral has already been determined, and thus the proposed integration is accomplished.

$$\begin{aligned} \text{Since } \int x^{m-1} X^p dx &= \int x^{m-1} X^{p-1} (a + bx^n) dx \\ &= a \int x^{m-1} X^{p-1} dx + b \int x^{m+n-1} X^{p-1} dx, \end{aligned}$$

we have by (1)

$$\frac{x^m X^p}{m} - \frac{bnp}{m} \int x^{m+n-1} X^{p-1} dx = a \int x^{m-1} X^{p-1} dx + b \int x^{m+n-1} X^{p-1} dx,$$

$$\text{therefore } \int x^{m-1} X^{p-1} dx = \frac{x^m X^p}{am} - \frac{b(m+np)}{am} \int x^{m+n-1} X^{p-1} dx.$$

Change  $p$  into  $p + 1$ , and we have

$$\int x^{m-1} X^p dx = \frac{x^m X^{p+1}}{am} - \frac{b(m+np+n)}{am} \int x^{m+n-1} X^p dx \dots \dots (3).$$

Change  $m$  into  $m - n$  and transpose, then

$$\int x^{m-1} X^p dx = \frac{x^{m-n} X^{p+1}}{b(m+np)} - \frac{(m-n)a}{b(m+np)} \int x^{m-n-1} X^p dx \dots \dots (4).$$

In (2), change  $m$  into  $m + n$  and  $p$  into  $p - 1$ , then

$$\int x^{m+n-1} X^{p-1} dx = \frac{x^m X^p}{bnp} - \frac{m}{bnp} \int x^{m-1} X^p dx.$$

$$\text{Also } \int x^{m-1} X^p dx = a \int x^{m-1} X^{p-1} dx + b \int x^{m+n-1} X^{p-1} dx,$$

$$\text{therefore } \int x^{m-1} X^p dx = a \int x^{m-1} X^{p-1} dx + \frac{x^m X^p}{np} - \frac{m}{np} \int x^{m-1} X^p dx,$$

$$\text{therefore } \int x^{m-1} X^p dx = \frac{x^m X^p}{m+np} + \frac{anp}{m+np} \int x^{m-1} X^{p-1} dx \dots \dots (5).$$

Change  $p$  into  $p + 1$  and transpose; thus

$$\int x^{m-1} X^p dx = -\frac{x^m X^{p+1}}{an(p+1)} + \frac{m+np+n}{an(p+1)} \int x^{m-1} X^{p+1} dx \dots \dots (6).$$

31. If an example is proposed to which one of the preceding formulæ is applicable, we may either quote that particular formula or may obtain the required result independently. Thus, suppose we require  $\int \frac{x^{m-1} dx}{\sqrt{(c^2 - x^2)}}$ ; we have

$$\begin{aligned} \int \frac{x^{m-1} dx}{\sqrt{(c^2 - x^2)}} &= - \int \frac{d \sqrt{(c^2 - x^2)}}{dx} x^{m-2} dx \\ &= - \sqrt{(c^2 - x^2)} x^{m-2} + (m-2) \int x^{m-3} \sqrt{(c^2 - x^2)} dx \\ &= - \sqrt{(c^2 - x^2)} x^{m-2} + (m-2) \int \frac{(c^2 - x^2) x^{m-3} dx}{\sqrt{(c^2 - x^2)}}. \end{aligned}$$

By transposition,

$$(1+m-2) \int \frac{x^{m-1} dx}{\sqrt{(c^2 - x^2)}} = -\sqrt{(c^2 - x^2)} x^{m-2} + (m-2) c^2 \int \frac{x^{m-3} dx}{\sqrt{(c^2 - x^2)}},$$

therefore

$$\int \frac{x^{m-1} dx}{\sqrt{(c^2 - x^2)}} = - \frac{x^{m-2} \sqrt{(c^2 - x^2)}}{m-1} + \frac{(m-2) c^2}{m-1} \int \frac{x^{m-3} dx}{\sqrt{(c^2 - x^2)}} \dots\dots(1).$$

This result agrees with the equation (4) of the preceding article if we make  $a = c^2$ ,  $b = -1$ ,  $n = 2$ ,  $p = -\frac{1}{2}$ .

Another example is furnished by  $\int \frac{x^m dx}{\sqrt{(2ax - x^2)}}$ , which may be written  $\int \frac{x^{m-1} dx}{\sqrt{(2a-x)}}$ ; if in equation (4) of the preceding article we make  $b = -1$ ,  $n = 1$ ,  $p = -\frac{1}{2}$ , and change  $a$  and  $m$  into  $2a$  and  $m + \frac{1}{2}$  respectively, we have

$$\int \frac{x^m dx}{\sqrt{(2ax - x^2)}} = - \frac{x^{m-1} \sqrt{(2ax - x^2)}}{m} + \frac{a(2m-1)}{m} \int \frac{x^{m-1} dx}{\sqrt{(2ax - x^2)}} \dots\dots\dots(2),$$

which of course may be found independently.

32. In equation (6) of Art. 30 put  $a = c^2$ ,  $m = 1$ ,  $n = 2$ ,  $b = 1$ , and  $p = -r$ ; thus

$$\int \frac{dx}{(x^2 + c^2)^r} = \frac{x}{2(r-1)c^2(x^2 + c^2)^{r-1}} + \frac{2r-3}{2(r-1)c^2} \int \frac{dx}{(x^2 + c^2)^{r-1}}.$$

This formula will serve to reduce the form

$$\int \frac{(Ax + B) dx}{(x^2 - 2ax + a^2 + \beta^2)^r},$$

which occurs in Art. 18; for this last expression may be written thus

$$\int \frac{A(x - a) dx}{\{(x - a)^2 + \beta^2\}^r} + (Aa + B) \int \frac{dx}{\{(x - a)^2 + \beta^2\}^r},$$

that is

$$-\frac{A}{2(r-1)} \frac{1}{\{(x - a)^2 + \beta^2\}^{r-1}} + (Aa + B) \int \frac{dx}{\{(x - a)^2 + \beta^2\}^r}.$$

By putting  $x - a = x'$ , we have

$$\int \frac{dx}{\{(x - a)^2 + \beta^2\}^r} = \int \frac{dx'}{\{x'^2 + \beta^2\}^r},$$

and thus the above formula becomes applicable.

33. These formulæ of reduction are most useful when the integral has to be taken between certain limits. Suppose  $\phi(x)$ ,  $\chi(x)$ ,  $\psi(x)$ , functions of  $x$ , such that

$$\int \phi(x) dx = \chi(x) + \int \psi(x) dx,$$

then 
$$\int_a^b \phi(x) dx = \chi(b) - \chi(a) + \int_a^b \psi(x) dx,$$

as is obvious from Art. 3.

For example, it may be shewn that

$$\int (c^2 - x^2)^{\frac{n}{2}} dx = \frac{x(c^2 - x^2)^{\frac{n}{2}}}{n+1} + \frac{nc^2}{n+1} \int (c^2 - x^2)^{\frac{n}{2}-1} dx;$$

suppose  $\frac{n}{2}$  a *positive* quantity, then  $x(c^2 - x^2)^{\frac{n}{2}}$  vanishes both when  $x = 0$  and when  $x = c$ . Hence

$$\int_0^c (c^2 - x^2)^{\frac{n}{2}} dx = \frac{nc^2}{n+1} \int_0^c (c^2 - x^2)^{\frac{n}{2}-1} dx.$$

The following is a similar example. By integration by parts

$$\int x^{r-1}(1-x)^{n-1} dx = -\frac{(1-x)^n}{n} x^{r-1} + \frac{r-1}{n} \int x^{r-2}(1-x)^n dx.$$

$$\text{Hence } \int_0^1 x^{r-1}(1-x)^{n-1} dx = \frac{r-1}{n} \int_0^1 x^{r-2}(1-x)^n dx.$$

Thus if  $r$  be an integer we may reduce the integral to  $\int_0^1 (1-x)^{n+r-2} dx$ , that is  $\frac{1}{n+r-1}$ ; hence

$$\int_0^1 x^{r-1}(1-x)^{n-1} dx = \frac{(r-1)(r-2)\dots\dots 3 \cdot 2 \cdot 1}{n(n+1)(n+2)\dots\dots(n+r-1)}.$$

34. The integration of trigonometrical functions is facilitated by formulæ of reduction. Let  $\phi(\sin x, \cos x)$  denote any function of  $\sin x$  and  $\cos x$ ; then if we put  $\sin x = z$ , we have

$$\begin{aligned} \int \phi(\sin x, \cos x) dx &= \int \phi\{z, \sqrt{(1-z^2)}\} \frac{dx}{dz} dz \\ &= \int \phi\{z, \sqrt{(1-z^2)}\} \frac{dz}{\sqrt{(1-z^2)}} \dots\dots\dots (1). \end{aligned}$$

For example, let  $\phi(\sin x, \cos x) = \sin^p x \cos^q x$ ; then

$$\int \sin^p x \cos^q x dx = \int z^p (1-z^2)^{\frac{1}{2}(q-1)} dz \dots\dots\dots (2).$$

If in the six formulæ of Art. 30 we put  $a=1$ ,  $b=-1$ ,  $n=2$ ,  $p=\frac{1}{2}(q-1)$ , we have

$$\begin{aligned} \int z^{m-1}(1-z^2)^{\frac{1}{2}(q-1)} dz \\ &= \frac{z^m(1-z^2)^{\frac{1}{2}(q-1)}}{m} + \frac{q-1}{m} \int z^{m+1}(1-z^2)^{\frac{1}{2}(q-3)} dz \\ &= -\frac{z^{m-2}(1-z^2)^{\frac{1}{2}(q+1)}}{q+1} + \frac{m-2}{q+1} \int z^{m-3}(1-z^2)^{\frac{1}{2}(q+1)} dz \end{aligned}$$

$$\begin{aligned}
 &= \frac{z^m (1-z^2)^{\frac{1}{2}(q+1)}}{m} + \frac{m+q+1}{m} \int z^{m+1} (1-z^2)^{\frac{1}{2}(q-1)} dz \\
 &= -\frac{z^{m-2} (1-z^2)^{\frac{1}{2}(q+1)}}{m+q-1} + \frac{m-2}{m+q-1} \int z^{m-3} (1-z^2)^{\frac{1}{2}(q-1)} dz \\
 &= \frac{z^m (1-z^2)^{\frac{1}{2}(q-1)}}{m+q-1} + \frac{q-1}{m+q-1} \int z^{m-1} (1-z^2)^{\frac{1}{2}(q-3)} dz \\
 &= -\frac{z^m (1-z^2)^{\frac{1}{2}(q+1)}}{q+1} + \frac{m+q+1}{q+1} \int z^{m-1} (1-z^2)^{\frac{1}{2}(q+1)} dz.
 \end{aligned}$$

If we put  $m = p + 1$ , and  $z = \sin x$ , the first of the above equations becomes

$$\int \sin^p x \cos^q x dx = \frac{\sin^{p+1} x \cos^{q-1} x}{p+1} + \frac{q-1}{p+1} \int \sin^{p+2} x \cos^{q-2} x dx,$$

and similarly the other five equations may be expressed.

35. The following is a very important case :

$$\begin{aligned}
 \int \sin^n x dx &= -\int \frac{d \cos x}{dx} \cdot \sin^{n-1} x \cdot dx \\
 &= -\cos x \cdot \sin^{n-1} x + (n-1) \int \cos^2 x \sin^{n-2} x dx \\
 &= -\cos x \cdot \sin^{n-1} x + (n-1) \int (1 - \sin^2 x) \sin^{n-2} x dx.
 \end{aligned}$$

Transposing, we have

$$n \int \sin^n x dx = -\cos x \sin^{n-1} x + (n-1) \int \sin^{n-2} x dx;$$

$$\text{therefore } \int \sin^n x dx = -\frac{\cos x \sin^{n-1} x}{n} + \frac{n-1}{n} \int \sin^{n-2} x dx.$$

From the last equation we deduce

$$\int_0^{\frac{1}{2}\pi} \sin^n x dx = \frac{n-1}{n} \int_0^{\frac{1}{2}\pi} \sin^{n-2} x dx.$$



Similarly 
$$\int_0^{i\pi} \sin^{n-2} x dx = \frac{n-3}{n-2} \int_0^{i\pi} \sin^{n-4} x dx.$$

Proceeding thus we shall arrive, if  $n$  be an *even* integer, at  $\int_0^{i\pi} dx$  or  $\frac{1}{2}\pi$ ; if  $n$  be an *odd* integer we shall arrive at

$\int_0^{i\pi} \sin x dx$ , which is unity. Hence, if  $n$  be an integer,

$$\int_0^{i\pi} \sin^n x dx = \frac{(n-1)(n-3)(n-5)\dots\dots 1}{n(n-2)(n-4)\dots\dots 2} \frac{\pi}{2} \quad (n \text{ even}),$$

$$\int_0^{i\pi} \sin^n x dx = \frac{(n-1)(n-3)(n-5)\dots\dots 2}{n(n-2)(n-4)\dots\dots 3} \quad (n \text{ odd}).$$

These two results hold if we change  $\sin x$  into  $\cos x$ , as will be found on investigation.

36. From the preceding results we may deduce an important theorem, called Wallis's Formula.

Suppose  $n$  even; then

$$\int_0^{i\pi} \sin^n x dx = \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots\dots \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} \dots\dots (1),$$

$$\int_0^{i\pi} \sin^{n-1} x dx = \frac{n-2}{n-1} \cdot \frac{n-4}{n-3} \cdot \frac{n-6}{n-5} \dots\dots \frac{2}{3} \dots\dots (2).$$

Now it is obvious that  $\int_0^{i\pi} \sin^{n-1} x dx$  is less than  $\int_0^{i\pi} \sin^{n-2} x dx$  and greater than  $\int_0^{i\pi} \sin^n x dx$ ; because each element of the first integral is less than the corresponding element of the second integral and greater than the corresponding element of the third integral.

Thus  $\frac{\int_0^{\pi} \sin^n x dx}{\int_0^{\pi} \sin^{n-1} x dx}$  is less than 1 and greater than  $\frac{n-1}{n}$ .

Hence the ratio of the right hand member of (1) to the right hand member of (2) is less than unity and greater than  $\frac{n-1}{n}$ ; thus

$$\frac{\pi}{2} > \frac{2 \cdot 2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \dots (n-2)(n-2)}{1 \cdot 3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \dots (n-3)(n-1)},$$

and  $< \frac{2 \cdot 2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \dots (n-2)(n-2)}{1 \cdot 3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \dots (n-3)(n-1)} \frac{n}{n-1}$ .

EXAMPLES.

1.  $\int (a^2 + x^2)^{\frac{n}{2}} dx = \frac{x(a^2 + x^2)^{\frac{n}{2}}}{n+1} + \frac{na^2}{n+1} \int (a^2 + x^2)^{\frac{n}{2}-1} dx.$
2.  $\int x^m \sqrt{(2ax - x^2)} dx = -\frac{x^{m+1}(2ax - x^2)^{\frac{1}{2}}}{m+2} + \frac{a(2m+1)}{m+2} \int x^{m-1} \sqrt{(2ax - x^2)} dx.$
3.  $\int x \sqrt{(2ax - x^2)} dx = -\frac{1}{3}(2ax - x^2)^{\frac{3}{2}} + a \int \sqrt{(2ax - x^2)} dx.$
4.  $\int_0^{2a} x \sqrt{(2ax - x^2)} dx = \frac{\pi a^3}{2}.$
5.  $\int x^2 \sqrt{(2ax - x^2)} dx = -\frac{x}{4}(2ax - x^2)^{\frac{3}{2}} + \frac{5a}{4} \int x \sqrt{(2ax - x^2)} dx.$
6.  $\int_0^{2a} x^2 \sqrt{(2ax - x^2)} dx = \frac{5a^4 \pi}{8}.$

$$7. \int_0^{2a} x^3 \sqrt{2ax - x^2} dx = \frac{7\pi a^5}{8}.$$

$$8. \int x^n (\log x)^m dx = \frac{x^{n+1} (\log x)^m}{n+1} - \frac{m}{n+1} \int x^n (\log x)^{m-1} dx.$$

$$9. \int x^n (\log x)^2 dx = \frac{x^{n+1}}{n+1} \left\{ (\log x)^2 - \frac{2}{n+1} \log x + \frac{2}{(n+1)^2} \right\}$$

$$10. \int_0^{\frac{\pi}{4}} \sec^4 \theta d\theta = \frac{5}{8}.$$

$$11. \int_0^a \frac{x^3 \sqrt{a-x}}{\sqrt{a+x}} dx = \left( \frac{\pi}{4} - \frac{3}{8} \right) a^3.$$

$$12. \int \sin^3 \theta \cos^3 \theta d\theta = -\frac{1}{4} \cos^4 \theta + \frac{1}{8} \cos^6 \theta.$$

$$13. \int \frac{d\theta}{\sin^4 \theta \cos^4 \theta} = 3 (\tan \theta - \cot \theta) + \frac{1}{3} (\tan^3 \theta - \cot^3 \theta).$$

$$14. \int \frac{\sin^3 \theta d\theta}{\cos^3 \theta} = \frac{\sin \theta}{2 \cos^2 \theta} + \frac{1}{4} \log \frac{1 - \sin \theta}{1 + \sin \theta}.$$

$$15. \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} (\cos 2\theta)^{\frac{3}{2}} \cos \theta d\theta = \frac{3\pi\sqrt{2}}{16}.$$

Assume  $\sqrt{2} \sin \theta = \sin$

$$16. \int_0^a \sqrt{a^2 - x^2} \cos^{-1} \frac{x}{a} dx = \left( 1 + \frac{\pi^2}{4} \right) \frac{a^2}{4}.$$

$$17. \int_0^{2a} \left( \text{vers}^{-1} \frac{x}{a} \right)^2 dx = (\pi^2 - 4) a.$$

$$18. \int_0^{\frac{\pi}{2}} \frac{\sin^3 x dx}{1 + c \cos x} = \frac{c^3 - 1}{c^3} \log(1 + c) + \frac{2 - c}{2c^3}.$$

19. If  $\phi(n) = \int (1 + c \cos x)^{-n} dx$ , shew that

$$(n-1)(1-c^2)\phi(n) = -c \sin x (1+c \cos x)^{-n+1} \\ + (2n-3)\phi(n-1) - (n-2)\phi(n-2).$$

$$20. \int_0^{2a} \sqrt{(2ax-x^2)} \operatorname{vers}^{-1} \frac{x}{a} dx = \frac{\pi^2 a^2}{4}.$$

$$21. \int_0^{2a} x \sqrt{(2ax-x^2)} \operatorname{vers}^{-1} \frac{x}{a} dx = \frac{4a^3}{9} + \frac{\pi^2 a^3}{4}.$$

$$22. \int_0^{\frac{1}{2}\pi} (\tan x)^7 dx = \frac{1}{12} - \frac{1}{2} \log 2.$$

$$23. \int_0^{\frac{\pi}{2}} \frac{dx}{\sqrt{(1-c^2 \sin^2 x)}} = \frac{\pi}{2} \left\{ 1 + \left(\frac{1}{2}\right)^2 c^2 + \left(\frac{1.3}{2.4}\right)^2 c^4 + \left(\frac{1.3.5}{2.4.6}\right)^2 c^6 + \dots \right\}$$

$c$  being  $< 1$ .

## CHAPTER IV.

## MISCELLANEOUS REMARKS.

37. WE have at the beginning of this book defined the *integral* of  $\phi(x)$  between assigned limits  $a$  and  $b$  as the limit of a certain sum  $\Sigma \phi(x) \Delta x$ , and have denoted this limit by  $\int_a^b \phi(x) dx$ . We have shewn that this limit is known as soon as we know the function  $\psi(x)$  of which  $\phi(x)$  is the differential coefficient. In the pages immediately following we gave methods for finding  $\psi(x)$  in different cases. We shall now add some miscellaneous remarks and theorems, some of which will recall the attention of the student to the process of summation which we placed at the foundation of the subject.

38. Suppose we wish to find the integral of  $\sin x$  between limits  $a$  and  $b$  *immediately from the definition*. By Art. 4 we have to find the limit when  $n$  is infinite of

$$h [\sin a + \sin (a + h) + \sin (a + 2h) \dots + \sin \{a + (n - 1) h\}]$$

where  $h = \frac{1}{n}(b - a)$ .

It is known from Trigonometry that this series

$$= \frac{h \sin \left( a + \frac{n-1}{2} h \right) \sin \frac{nh}{2}}{\sin \frac{h}{2}} = \frac{h \sin \left( a + \frac{b-a}{2} - \frac{h}{2} \right) \sin \frac{b-a}{2}}{\sin \frac{h}{2}}.$$

The limit of  $\frac{h}{\sin \frac{h}{2}}$  is 2; hence the required integral is

$$2 \sin \frac{b+a}{2} \sin \frac{b-a}{2} = \cos a - \cos b.$$

39. Required the limit when  $n$  is made infinite of the series

$$\frac{n}{n^2} + \frac{n}{1+n^2} + \frac{n}{2^2+n^2} + \frac{n}{3^2+n^2} \dots\dots + \frac{n}{(n-1)^2+n^2}.$$

This series may be written

$$\frac{1}{n} \left\{ \frac{1}{1} + \frac{1}{1 + \left(\frac{1}{n}\right)^2} + \frac{1}{1 + \left(\frac{2}{n}\right)^2} + \frac{1}{1 + \left(\frac{3}{n}\right)^2} \dots\dots + \frac{1}{1 + \left(\frac{n-1}{n}\right)^2} \right\};$$

putting  $h$  for  $\frac{1}{n}$ , we obtain

$$h \left\{ \frac{1}{1} + \frac{1}{1+h^2} + \frac{1}{1+(2h)^2} \dots\dots + \frac{1}{1+(n-1)^2 h^2} \right\}.$$

Comparing this with Art. 4 we see that the required limit is what we denote by  $\int_0^1 \frac{dx}{1+x^2}$ . Now  $\int \frac{dx}{1+x^2} = \tan^{-1} x$ ; hence  $\frac{\pi}{4}$  is the required limit.

40. We define  $\int_a^b \phi(x) dx$  as the limit when  $n$  is infinite of

$$h_1 \phi(a) + h_2 \phi(x_1) \dots\dots + h_n \phi(x_{n-1}).$$

Now let  $A$  and  $B$  be the greatest and least values which  $\phi(x)$  takes between the limits  $a$  and  $b$ ; then the series is less than

$$(h_1 + h_2 + \dots\dots + h_n) A,$$

and is greater than

$$(h_1 + h_2 + \dots + h_n) B;$$

that is, the series lies between

$$(b-a) A \text{ and } (b-a) B.$$

The limit must therefore be equal to  $(b-a) C$  where  $C$  is some quantity lying between  $A$  and  $B$ ; but since  $\phi(x)$  is supposed continuous, it must, while  $x$  ranges from  $a$  to  $b$ , pass through every value between  $A$  and  $B$ , and must therefore be equal to  $C$  when  $x$  has some value between  $a$  and  $b$ . Thus  $C = \phi\{a + \theta(b-a)\}$  where  $\theta$  is some proper fraction, and

$$\int_a^b \phi(x) dx = (b-a) \phi\{a + \theta(b-a)\}.$$

41. The truth of the equation

$$\int_a^b \phi(x) dx = \int_a^c \phi(x) dx + \int_c^b \phi(x) dx \dots \dots \dots (1)$$

will appear immediately; for suppose  $\psi(x)$  to be the integral of  $\phi(x)$ , then we have on the left-hand side

$$\psi(b) - \psi(a),$$

and on the right hand

$$\psi(c) - \psi(a) + \psi(b) - \psi(c).$$

In like manner

$$\int_a^b \phi(x) dx = - \int_b^a \phi(x) dx \dots \dots \dots (2)$$

is obviously true. We may shew also that

$$\int_0^a \phi(x) dx = \int_0^a \phi(a-x) dx \dots \dots \dots (3)$$

For putting  $a-x=z$  we have

$$\int \phi(a-x) dx = - \int \phi(z) dz,$$

therefore 
$$\int_0^a \phi(a-x) dx = - \int_a^0 \phi(z) dz$$

$$= \int_0^a \phi(z) dz, \text{ by (2).}$$

Of course  $\int_0^a \phi(z) dz = \int_0^a \phi(x) dx$ , since it is indifferent whether we use the symbol  $x$  or  $z$  in obtaining a result which does not involve  $x$  or  $z$ .

We have from (1)

$$\int_0^{2a} \phi(x) dx = \int_0^a \phi(x) dx + \int_a^{2a} \phi(x) dx.$$

The second integral, by changing  $x$  into  $2a - x'$ , will be found equal to

$$\int_0^a \phi(2a - x') dx' \text{ or } \int_0^a \phi(2a - x) dx.$$

Hence

$$\int_0^{2a} \phi(x) dx = \int_0^a \{\phi(x) + \phi(2a - x)\} dx.$$

Hence, if  $\phi(x) = \phi(2a - x)$  for all values of  $x$  comprised between 0 and  $a$ , we have

$$\int_0^{2a} \phi(x) dx = 2 \int_0^a \phi(x) dx \dots \dots \dots (4),$$

and if  $\phi(2a - x) = -\phi(x)$ , we have

$$\int_0^{2a} \phi(x) dx = 0 \dots \dots \dots (5).$$

For example,

$$\int_0^\pi \sin^3 \theta d\theta = 2 \int_0^{\frac{1}{2}\pi} \sin^3 \theta d\theta \dots \dots \text{by (4),}$$

and 
$$\int_0^\pi \cos^3 \theta d\theta = 0 \dots \dots \text{by (5).}$$



42. Such equations as those just given should receive careful attention from the student, and he should not leave them until he recognizes their obvious and self-evident truth.

$\int_0^\pi \cos^2 \theta d\theta$  is by definition the limit when  $n$  is infinite of the series

$$h \{ \cos^2 h + \cos^2 2h + \cos^2 3h \dots + \cos^2 (n-1) h \}$$

where  $nh = \pi$ . Now

$$\cos^2 h = -\cos^2 (n-1) h, \quad \cos^2 2h = -\cos^2 (n-2) h, \dots;$$

thus the positive terms of the series just balance the negative terms and leave zero as the result.

In the same way the truth of  $\int_0^\pi \sin^2 \theta d\theta = 2 \int_0^{\frac{\pi}{2}} \sin^2 \theta d\theta$  follows *immediately* from the definition of integration, and the fact that the sine of an angle is equal to the sine of the supplemental angle.

43. Suppose  $b$  greater than  $a$  and  $\phi(x)$  always positive between the limits  $a$  and  $b$  of  $x$ ; then every term in the series  $\Sigma \phi(x) \Delta x$  is positive, and hence the limit  $\int_a^b \phi(x) dx$  must be a positive quantity.

44. The statement of the last article supposes that  $\phi(x)$  is always finite between the limits  $a$  and  $b$ ; it must be remembered that this condition was expressly introduced in the fundamental proposition, Art. 2. If therefore the function to be integrated becomes infinite between the limits of integration, the rules of integration cannot be applied; at least the case must be specially examined.

45. Consider  $\int_0^a \frac{dx}{\sqrt{1-x}}$ ; the value of this integral is  $2 - 2\sqrt{1-a}$ . Here the function to be integrated becomes infinite when  $x=1$ ; but the expression  $2 - 2\sqrt{1-a}$  is finite when  $a=1$ . Hence in this case we may write  $\int_0^1 \frac{dx}{\sqrt{1-x}} = 2$ , provided that we regard this as an abbreviation

tion of the following statement: " $\int_0^a \frac{dx}{\sqrt{1-x}}$  is always finite if  $a$  be any quantity less than unity, and by taking  $a$  sufficiently near to unity, we can make the value of the integral differ as little as we please from 2."

46. Next take  $\int_0^a \frac{dx}{1-x}$ ; the value of this integral is  $-\log(1-a)$  which increases indefinitely as  $a$  approaches to unity. Hence in this case we may write  $\int_0^1 \frac{dx}{1-x} = \infty$  provided that we regard this as an abbreviation of the following statement: " $\int_0^a \frac{dx}{1-x}$  increases indefinitely as  $a$  approaches to unity, and by taking  $a$  sufficiently near to unity we can make the integral greater than any assigned quantity."

47. Next consider  $\int \frac{dx}{(1-x)^2}$ ; the integral here is  $\frac{1}{1-x}$ . If without remarking that the function to be integrated becomes infinite when  $x=1$ , we propose to find the value of the integral between the limits 0 and 2, we obtain  $-1-1$ , that is  $-2$ . But this is obviously false, for in this case every term of the series indicated by  $\sum \phi(x) \Delta x$  is positive, and therefore the limit cannot be negative. In fact  $\int_0^1 \frac{dx}{(1-x)^2}$  and  $\int_1^2 \frac{dx}{(1-x)^2}$  are both infinite. This example shews that the ordinary rules for integrating between assigned limits cannot be used when the function to be integrated becomes infinite between those limits.

48. In the fundamental investigation in Art. 2, of the value of  $\int_a^b \phi(x) dx$ , the limits  $a$  and  $b$  are supposed to be finite as well as the function  $\phi(x)$ . But we shall often find it convenient to suppose one or both of the limits infinite, as we will now indicate by examples.

Consider  $\int \frac{dx}{1+x^2}$ ; the integral is  $\tan^{-1} x$ . Hence  $\int_0^a \frac{dx}{1+x^2} = \tan^{-1} a$ ; the larger  $a$  becomes, the nearer  $\tan^{-1} a$  approaches

to  $\frac{\pi}{2}$ , and by taking  $a$  sufficiently large, we can make  $\tan^{-1} a$  differ as little as we please from  $\frac{\pi}{2}$ ; hence we may write  $\int_0^{\infty} \frac{dx}{1+x^2} = \frac{\pi}{2}$  as an abbreviation of this statement.

Similarly  $\int_0^a \frac{dx}{1+x} = \log(1+a)$ ; and by taking  $a$  large enough we can make  $\log(1+a)$  greater than any assigned quantity. Hence for abbreviation we may write

$$\int_0^{\infty} \frac{dx}{1+x} = \infty.$$

49. Suppose the function  $\phi(x)$  to become infinite *once* between the limits  $a$  and  $b$ , namely, when  $x=c$ . We cannot then apply the ordinary rules of integration to  $\int_a^b \phi(x) dx$ ; but we may apply those rules to

$$\int_a^{c-\mu} \phi(x) dx + \int_{c+\mu}^b \phi(x) dx$$

for any assigned value of  $\mu$  however small. The limit of the last expression when  $\mu$  is diminished indefinitely is called by Cauchy the *principal* value of the integral  $\int_a^b \phi(x) dx$ .

For example, let  $\phi(x) = \frac{1}{c-x}$ ;

$$\text{then } \int_a^{c-\mu} \frac{dx}{c-x} = \log \frac{c-a}{\mu},$$

and  $\int_{c+\mu}^b \frac{dx}{c-x} = -\int_{c+\mu}^b \frac{dx}{x-c} = -\log \frac{b-c}{\mu}$ ;

hence the *principal* value is  $\log \frac{c-a}{\mu} - \log \frac{b-c}{\mu}$ , that is  $\log \frac{c-a}{b-c}$ .

50. The value of  $\int \frac{dx}{\sqrt{(a^2 - x^2)}}$  is  $\sin^{-1} \frac{x}{a}$ ; hence

$$\int_{-a}^a \frac{dx}{\sqrt{(a^2 - x^2)}} = \sin^{-1}(1) - \sin^{-1}(-1).$$

Students are sometimes doubtful respecting the value which is to be assigned to  $\sin^{-1}(1)$  and  $\sin^{-1}(-1)$  in such a result as the above. Suppose we assume  $x = a \sin \theta$ ; thus the integral becomes  $\int d\theta$  or  $\theta$ . Now  $x$  increases from  $-a$  to  $a$ , hence the limits assigned to  $\theta$  must be such as correspond to this range of values of  $x$ . When  $x = -a$  then  $\theta$  may have any value contained in the formula  $(4n - 1) \frac{\pi}{2}$ , where  $n$  is any integer. Suppose we take the value  $(4n - 1) \frac{\pi}{2}$ , where  $n$  is some definite integer, then corresponding to the value  $x = a$  we *must* take  $\theta = (4n - 1) \frac{\pi}{2} + \pi$ ; this will be obvious on examination, because  $x$  is to change from  $-a$  to  $+a$ , so that it *continually increases and only once passes through the value zero*.

Hence 
$$\int_{-a}^a \frac{dx}{\sqrt{(a^2 - x^2)}} = \pi.$$

51. Required  $\int_0^{\frac{1}{2}\pi} \log \sin x \, dx$ .

By equation (3) of Art. 41,

$$\int_0^{\frac{1}{2}\pi} \log \sin x \, dx = \int_0^{\frac{1}{2}\pi} \log \sin \left( \frac{\pi}{2} - x \right) dx = \int_0^{\frac{1}{2}\pi} \log \cos x \, dx.$$

Hence, putting  $y$  for the required integral,

$$\begin{aligned} 2y &= \int_0^{\frac{1}{2}\pi} (\log \sin x + \log \cos x) \, dx \\ &= \int_0^{\frac{1}{2}\pi} \log (\sin x \cos x) \, dx \end{aligned}$$

$$\begin{aligned}
 &= \int_0^{2\pi} \log \frac{\sin 2x}{2} dx \\
 &= \int_0^{2\pi} \{\log \sin 2x - \log 2\} dx \\
 &= \int_0^{2\pi} \log \sin 2x dx - \frac{1}{2} \pi \log 2.
 \end{aligned}$$

But putting  $2x = x'$ , we have

$$\begin{aligned}
 \int_0^{2\pi} \log \sin 2x dx &= \frac{1}{2} \int_0^{2\pi} \log \sin x' dx' \\
 &= \int_0^{\pi} \log \sin x dx, \text{ by equation (4) of Art. 41;}
 \end{aligned}$$

therefore  $2y = y - \frac{\pi}{2} \log 2,$

therefore  $y = \frac{\pi}{2} \log \frac{1}{2}.$

Again,  $\int_0^{\pi} \theta^2 \log \sin \theta d\theta = \int_0^{\pi} (\pi - \theta)^2 \log \sin \theta d\theta,$  by equation (3) of Art. 41; therefore

$$0 = \int_0^{\pi} (\pi^2 - 2\pi\theta) \log \sin \theta d\theta,$$

therefore  $\int_0^{\pi} \theta \log \sin \theta d\theta = \frac{\pi}{2} \int_0^{\pi} \log \sin \theta d\theta$

$$= \frac{\pi^2}{2} \log \frac{1}{2}.$$

Required  $\int_0^1 \frac{\log(1+x)}{1+x^2} dx.$  Put  $x = \tan y,$  and the integral becomes  $\int_0^{\frac{\pi}{4}} \log(1 + \tan y) dy;$  but by equation (3) of Art. 41

$$\int_0^{\frac{\pi}{4}} \log(1 + \tan y) dy = \int_0^{\frac{\pi}{4}} \log \left\{ 1 + \tan \left( \frac{\pi}{4} - y \right) \right\} dy,$$

and  $1 + \tan\left(\frac{\pi}{4} - y\right) = 1 + \frac{1 - \tan y}{1 + \tan y} = \frac{2}{1 + \tan y}$ ;

therefore  $2 \int_0^{\frac{\pi}{4}} \log(1 + \tan y) dy = \frac{\pi}{4} \log 2$ .

See *Cambridge Mathematical Journal*, Vol. III. p. 168.

52. The remainder after  $n + 1$  terms of the expansion of  $\phi(a + h)$  in powers of  $h$ , may be expressed by a definite integral. For let

$$F(z) = \phi(x - z) + z\phi'(x - z) + \frac{z^2}{2} \phi''(x - z) \dots + \frac{z^n}{n} \phi^n(x - z).$$

Differentiate with respect to  $z$ , then

$$F'(z) = -\frac{z^n}{n} \phi^{n+1}(x - z).$$

Integrate both members of this equation between the limits 0 and  $h$ ; thus

$$F(h) - F(0) = -\frac{1}{n} \int_0^h z^n \phi^{n+1}(x - z) dz,$$

that is

$$\begin{aligned} \phi(x - h) + h\phi'(x - h) + \frac{h^2}{2} \phi''(x - h) \dots + \frac{h^n}{n} \phi^n(x - h) - \phi(x) \\ = -\frac{1}{n} \int_0^h z^n \phi^{n+1}(x - z) dz. \end{aligned}$$

Put  $a + h$  for  $x$  and transpose, then

$$\begin{aligned} \phi(a + h) = \phi(a) + h\phi'(a) + \frac{h^2}{2} \phi''(a) \dots + \frac{h^n}{n} \phi^n(a) \\ + \frac{1}{n} \int_0^h z^n \phi^{n+1}(a + h - z) dz. \end{aligned}$$

53. *Bernoulli's Series.* By integration by parts we have

$$\int \phi(x) dx = x \phi(x) - \int x \phi'(x) dx,$$

$$\int x \phi'(x) dx = \frac{x^2}{2} \phi'(x) - \int \frac{x^2}{2} \phi''(x) dx,$$

$$\int x^3 \phi''(x) dx = \frac{x^3}{3} \phi''(x) - \int \frac{x^3}{3} \phi'''(x) dx.$$

.....

$$\text{Thus, } \int \phi(x) dx = x \phi(x) - \frac{x^2}{1.2} \phi'(x) + \frac{x^3}{\underline{3}} \phi''(x) \dots\dots$$

$$+ \frac{(-1)^{n-1} x^n}{\underline{n}} \phi^{(n-1)}(x) + \frac{(-1)^n}{\underline{n}} \int x^n \phi^{(n)}(x) dx.$$

Therefore,

$$\int_0^a \phi(x) dx = a \phi(a) - \frac{a^2}{1.2} \phi'(a) + \frac{a^3}{\underline{3}} \phi''(a) \dots\dots$$

$$+ \frac{(-1)^{n-1} a^n \phi^{(n-1)}(a)}{\underline{n}} + \frac{(-1)^n}{\underline{n}} \int_0^a x^n \phi^{(n)}(x) dx.$$

This series on the right hand is called Bernoulli's series. In some cases this process might be of use in obtaining  $\int_0^a \phi(x) dx$ ; for example, if  $\phi(x)$  be any rational algebraical function of the  $(n-1)^{\text{th}}$  degree,  $\phi^{(n)}x$  is zero; or it might happen that  $\int x^n \phi^{(n)}(x) dx$  could be found more easily than  $\int \phi(x) dx$ . Or again, we may require only an *approximate* value of  $\int_0^a \phi(x) dx$  and the integral  $\int_0^a x^n \phi^{(n)}(x) dx$  might be small enough to be neglected.

54. By adopting different methods of integrating a function, we may apparently sometimes arrive at different results. But we know (*Dif. Calc.* Art. 102) that two functions which have the same differential coefficient can only differ by a

constant, so that any two results which we obtain must either be identical or differ by a constant. Take for example

$$\int (ax + b)(a'x + b') dx;$$

integrate by parts, thus we obtain

$$\frac{(ax + b)^2}{2a} (a'x + b') - \int \frac{a'}{2a} (ax + b)^2 dx,$$

that is, 
$$\frac{(ax + b)^2 (a'x + b')}{2a} - \frac{a' (ax + b)^3}{6a^2}.$$

If we integrate by parts in another way, we can obtain

$$\frac{(a'x + b')^2 (ax + b)}{2a'} - \frac{a (a'x + b')^3}{6a'^2}.$$

Hence

$$\frac{(ax + b)^2 \{3a (a'x + b') - a' (ax + b)\}}{6a^2}$$

and

$$\frac{(a'x + b')^2 \{3a' (ax + b) - a (a'x + b')\}}{6a'^2},$$

can only differ by a constant. Hence multiplying by  $6a^2a'^2$  we have

$$a'^2 (ax + b)^2 \{3a (a'x + b') - a' (ax + b)\} - a^2 (a'x + b')^2 \{3a' (ax + b) - a (a'x + b')\} = C$$

where  $C$  is some constant. This might of course be verified by common reduction. We may easily determine the value of  $C$ ; for since it is independent of  $x$  we may suppose  $ax + b = 0$ , that is,  $x = -\frac{b}{a}$ ; then the left hand member becomes  $(ab' - a'b)^2$ , which is consequently the value of  $C$ .

Similarly from

$$\int (ax + b) dx + \int (a'x + b') dx = \int \{(a + a')x + b + b'\} dx$$



we infer

$$\frac{(ax+b)^2}{2a} + \frac{(a'x+b')^2}{2a'} = \frac{\{(a+a')x+b+b'\}^2}{2(a+a')} + \text{constant.}$$

Multiply by  $2aa'(a+a')$  and then determine the constant by supposing  $x=0$ ; thus we obtain the identity

$$\begin{aligned} a'(a+a')(ax+b)^2 + a(a+a')(a'x+b')^2 \\ = aa'\{(a+a')x+b+b'\}^2 + (ba'-b'a)^2. \end{aligned}$$

55. By  $\int \phi(x) dx$  we indicate the function of which  $\phi(x)$  is the differential coefficient; suppose this to be  $\psi(x)$ . Then we may require the function of which  $\psi(x)$  is the differential coefficient, which we denote by  $\int \psi(x) dx$ , or by  $\iint \phi(x) dx dx$ , and so on. For example, the integral of  $e^{kx}$  is  $\frac{1}{k} e^{kx} + C_1$  where  $C_1$  is a constant; the integral of this is

$$\frac{1}{k^2} e^{kx} + C_1 x + C_2;$$

the integral of this is

$$\frac{1}{k^3} e^{kx} + C_1 \frac{x^2}{2} + C_2 x + C_3,$$

where  $\frac{C_1}{2}$  being still a constant may be denoted for simplicity by  $B$  if we please. Proceeding thus we should find as the result of integrating  $e^{kx}$  successively for  $n$  times

$$\frac{e^{kx}}{k^n} + A_1 x^{n-1} + A_2 x^{n-2} + \dots + A_{n-1} x + A_n$$

where  $A_1, A_2, \dots, A_n$  are constants.

MISCELLANEOUS EXAMPLES.

1.  $\int_0^a \frac{x^{\frac{1}{2}} dx}{\sqrt{a-x}} = \frac{5\pi a^{\frac{3}{2}}}{16}$ . (Assume  $x = a \sin^2 \theta$ ).

2.  $\int_0^{2a} \frac{x dx}{\sqrt{2ax - x^2}} = \pi a$ .

3.  $\int_0^a \frac{(a^2 - e^2 x^2) dx}{\sqrt{(a^2 - x^2)}} = \frac{\pi a^2}{2} \left(1 - \frac{e^2}{2}\right)$ .

4.  $\int_0^{\infty} \frac{dx}{(a^2 + x^2)(b^2 + x^2)} = \frac{\pi}{2ab(a+b)}$ .

5. If  $\phi(x) = \phi(a+x)$ , shew that

$$\int_0^{na} \phi(x) dx = n \int_0^a \phi(x) dx.$$

6. Shew that  $\int_a^b \phi(x) dx = \frac{b-a}{2c} \int_{-c}^c \phi\left(\frac{b+a}{2} + \frac{b-a}{2c} x\right) dx$ .

7. Shew that  $\int_0^{\pi} \frac{x \sin x dx}{1 + \cos^2 x} = \frac{\pi^2}{4}$ . (See Art. 41.)

8. Shew that  $\int_0^{2a} \sqrt{2ax - x^2} \operatorname{vers}^{-1} \frac{x}{a} dx = \frac{\pi^2 a^2}{4}$ .

(Change  $x$  into  $2a - x'$ ; see Art. 41.)

9. Find the limit when  $n$  is infinite of

$$\frac{1}{n} + \frac{1}{\sqrt{(n^2 - 1)}} + \frac{1}{\sqrt{(n^2 - 2^2)}} + \dots + \frac{1}{\sqrt{\{n^2 - (n-1)^2\}}}.$$

Result  $\frac{\pi}{2}$ .

10. Find the limit when  $n$  is infinite of

$$\frac{\left(\frac{1}{n}\right)^p + \left(\frac{2}{n}\right)^p + \left(\frac{3}{n}\right)^p \dots + 1}{\left(\frac{1}{2} + \frac{1}{n}\right)^p + \left(\frac{1}{2} + \frac{2}{n}\right)^p \dots + 1} \quad \text{Result. } \frac{1}{1 - \left(\frac{1}{2}\right)^{p+1}}.$$

11. Find the limit when  $n$  is infinite of  $\left\{\frac{[n]}{n^n}\right\}^{\frac{1}{n}}$ .

*Result.*  $\frac{1}{e}$ . (Take the logarithm of the expression.)

12. Shew that  $\int_0^{\frac{\pi}{2}} \log \tan x \, dx = 0$ .

13. Prove that

$$\int_a^{a+\theta c} \phi(x) \chi(x) \, dx = \phi(a + \theta c) \int_a^{a+\theta c} \chi(x) \, dx,$$

where  $\theta$  is a proper fraction, provided that  $\phi(x)$  and  $\chi(x)$  are finite and continuous from  $x = a$  to  $x = a + c$ , and that  $\chi(x)$  is of invariable sign between these limits.

14. If  $f(x)$  be positive and finite from  $x = a$  to  $x = a + c$ , shew how to find the limit of

$$\left\{f(a) f\left(a + \frac{c}{n}\right) \dots f\left(a + \frac{n-1}{n} c\right)\right\}^{\frac{1}{n}}$$

when  $n$  is infinite; and prove that the limit in question is less than  $\frac{1}{c} \int_a^{a+c} f(x) \, dx$ , assuming that the geometric mean of a finite number of positive quantities which are not all equal is less than the arithmetic.

Hence prove that  $e^{\int_0^1 u \, dx}$  is less than  $\int_0^1 e^u \, dx$ , unless  $u$  be constant from  $x = 0$  to  $x = 1$ .

15. The value of the definite integral  $\int_0^{\frac{\pi}{2}} \log(1 + n \cos^2 \theta) d\theta$  may be found whatever positive value is given to  $n$  from the formula

$$\int_0^{\frac{\pi}{2}} \log(1 + n \cos^2 \theta) d\theta = \frac{\pi}{4} \log \{(1+n)(1+n_1)(1+n_2)\dots\}$$

where  $n, n_1, n_2, \dots$  are quantities connected by the equation

$$n_{r+1} = \frac{n_r^2}{4(n_r + 1)}.$$

(Put  $\theta = \frac{\pi}{2} - \theta'$ ; see Art. 41.)

16. Shew that

$$\int e^{ax} \cos ax \, dx = \frac{e^{ax} \cos(ax - \phi)}{(a^2 + c^2)^{\frac{1}{2}}} + \text{a constant},$$

where  $\tan \phi = \frac{a}{c}$ . Hence shew that if  $e^{ax} \cos ax$  be integrated  $n$  times successively the result is

$$\frac{e^{ax} \cos(ax - n\phi)}{(a^2 + c^2)^{\frac{n}{2}}} + C + C_1 x + C_2 x^2 \dots + C_{n-1} x^{n-1}.$$

## CHAPTER V.

## DOUBLE INTEGRATION.

56. LET  $\phi(x)$  denote any function of  $x$ ; then we have seen that the *integral* of  $\phi(x)$  is a quantity  $u$  such that  $\frac{du}{dx} = \phi(x)$ . The integral may also be regarded as the limit of a certain sum (see Arts. 2—6), and hence is derived the symbol  $\int \phi(x) dx$  by which the integral is denoted. We now proceed to extend these conceptions of an integral to cases where we have more than one independent variable.

57. Suppose we have to find the value of  $u$  which satisfies the equation  $\frac{d^2u}{dy dx} = \phi(x, y)$ , where  $\phi(x, y)$  is a function of the independent variables  $x$  and  $y$ . The equation may be written

$$\frac{d}{dy} \left( \frac{du}{dx} \right) = \phi(x, y),$$

or 
$$\frac{dv}{dy} = \phi(x, y),$$

if  $v = \frac{du}{dx}$ . Thus  $v$  must be a function such that if we differentiate it with respect to  $y$ , considering  $x$  as constant, the result will be  $\phi(x, y)$ . We may therefore put

$$v = \int \phi(x, y) dy,$$

that is, 
$$\frac{du}{dx} = \int \phi(x, y) dy.$$

Hence  $u$  must be such a function that if we differentiate it with respect to  $x$ , considering  $y$  constant, the result will be the function denoted by  $\int \phi(x, y) dy$ . Hence

$$u = \int \left\{ \int \phi(x, y) dy \right\} dx.$$

The method of obtaining  $u$  may be described by saying that we first integrate  $\phi(x, y)$  with respect to  $y$ , and then integrate the result with respect to  $x$ .

The above expression for  $u$  may be more concisely written thus

$$\iint \phi(x, y) dy dx, \quad \text{or} \quad \iint \phi(x, y) dx dy.$$

On this point of notation writers are not quite uniform; we shall in the present work adopt the latter form, that is, of the two symbols  $dx$  and  $dy$  we shall put  $dy$  to the right, when we consider the integration with respect to  $y$  performed before the integration with respect to  $x$ , and *vice versa*.

58. We might find  $u$  by integrating first with respect to  $x$  and then with respect to  $y$ ; this process would be indicated by the equation

$$u = \iint \phi(x, y) dy dx.$$

59. Since we have thus *two methods* of finding  $u$  from the equation  $\frac{d^2 u}{dx dy} = \phi(x, y)$ , it will be desirable to investigate if more than *one result* can be obtained. Suppose then that  $u_1$  and  $u_2$  are two functions either of which when put for  $u$  satisfies the given equation, so that

$$\frac{d^2 u_1}{dx dy} = \phi(x, y) \quad \text{and} \quad \frac{d^2 u_2}{dx dy} = \phi(x, y).$$

We have, by subtraction,

$$\frac{d^2 u_1}{dx dy} - \frac{d^2 u_2}{dx dy} = 0,$$

that is,  $\frac{d}{dx} \left( \frac{dv}{dy} \right) = 0$  where  $v = u_1 - u_2$ .

Now from an equation  $\frac{dw}{dx} = 0$  we infer that  $w$  must be a *constant*, that is, must be a *constant* so far as relates to  $x$ ; in other words,  $w$  cannot be a function of  $x$ , but *may* be a function of any other variable which occurs in the question we are considering.

Thus from the equation  $\frac{d}{dx} \left( \frac{dv}{dy} \right) = 0$  we infer that  $\frac{dv}{dy}$  cannot be a function of  $x$ , but *may* be any arbitrary function of  $y$ . Thus we may put

$$\frac{dv}{dy} = f(y).$$

By integration we deduce

$$v = \int f(y) dy + \text{constant}.$$

Here the constant, as we call it, must not contain  $y$ , but may contain  $x$ ; we may denote it by  $\chi(x)$ . And  $\int f(y) dy$  we will denote by  $\psi(y)$ ; thus finally

$$v = \psi(y) + \chi(x).$$

Therefore two values of  $u$  which satisfy the equation  $\frac{d^2u}{dx dy} = \phi(x, y)$  can only differ by the sum of two arbitrary functions, one of  $x$  only and the other of  $y$  only.

60. We shall now shew the connexion between double integration and summation. Let  $\phi(x, y)$  be a function of  $x$  and  $y$ , which remains finite and continuous so long as  $x$  lies between the fixed values  $a$  and  $b$ , and  $y$  between the fixed values  $\alpha$  and  $\beta$ . Let  $a, x_1, x_2, \dots, x_{n-1}, b$  be a series of quantities in order of magnitude; also let  $\alpha, y_1, y_2, \dots, y_{m-1}, \beta$  be another series of quantities in order of magnitude.

$$\text{Let } x_1 - a = h_1, x_2 - x_1 = h_2, \dots, b - x_{n-1} = h_n;$$

$$\text{also let } y_1 - \alpha = k_1, y_2 - y_1 = k_2, \dots, \beta - y_{m-1} = k_m.$$





Now diminish indefinitely each term of which  $h$  is the type, then  $\Sigma h\rho$  vanishes, and we have finally

$$\int_a^b \psi(x) dx;$$

that is 
$$\int_a^b \left\{ \int_a^\beta \phi(x, y) dy \right\} dx.$$

This is more concisely written

$$\int_a^b \int_a^\beta \phi(x, y) dx dy,$$

$dy$  being placed to the right of  $dx$  because the integration is performed first with respect to  $y$ .

61. We may again remind the student that writers are not all agreed as to the notation for double integrals. Thus we use  $\int_a^b \int_a^\beta \phi(x, y) dx dy$  to imply the following order of operations—integrate  $\phi(x, y)$  with respect to  $y$  between the limits  $\alpha$  and  $\beta$ ; then integrate the result with respect to  $x$  between the limits  $a$  and  $b$ . Some writers would denote the same order of operations by  $\int_a^b \int_a^\beta \phi(x, y) dy dx$ .

62. We might have found the limit of the sum in Art. 60 by first taking all the terms in one vertical column, and then taking all the columns. In this way we should obtain as the sum  $\int_a^\beta \int_a^b \phi(x, y) dy dx$ ; and consequently

$$\int_a^\beta \int_a^b \phi(x, y) dy dx = \int_a^b \int_a^\beta \phi(x, y) dx dy.$$

63. Hitherto we have integrated both with respect to  $x$  and  $y$  between constant limits; the limits however in the first integration may be functions of the other variable. Thus, for example, the symbol  $\int_a^b \int_{\chi(x)}^{\psi(x)} \phi(x, y) dx dy$  will denote the following operations—first integrate with respect to  $y$  so that  $x$

is constant; suppose  $F(x, y)$  to be the integral; then by taking the integral between the assigned limits we have the result

$$F\{x, \psi(x)\} - F\{x, \chi(x)\}.$$

We have finally to obtain the integral indicated by

$$\int_a^b [F\{x, \psi(x)\} - F\{x, \chi(x)\}] dx.$$

The only difference which is required in the summatory process of Art. 60 is, that the quantities  $\alpha, y_1, y_2, \dots, y_{m-1}$  will not have the same meaning in each horizontal line. In the  $(r+1)^{\text{th}}$  line, for example, that is in

$$h_{r+1} \{k_1 \phi(x_r, \alpha) + k_2 \phi(x_r, y_1) + k_3 \phi(x_r, y_2) \dots + k_m \phi(x_r, y_{m-1})\}$$

we must consider  $\alpha$  as standing for  $\chi(x_r)$ , and  $y_1, y_2, \dots$  as a series of quantities, such that  $\chi(x_r), y_1, y_2, \dots, y_{m-1}, \psi(x_r)$ , are in order of magnitude, and that the difference between any consecutive two ultimately vanishes. Hence, proceeding as

before, we get  $\int_{\chi(x_r)}^{\psi(x_r)} \phi(x_r, y) dy$  for the limit of the sum of the terms in the  $(r+1)^{\text{th}}$  line.

64. It is not necessary to suppose the same number of terms in all the horizontal rows; for  $m$  is ultimately made indefinitely great, so that we obtain the same expression for the limit of the  $(r+1)^{\text{th}}$  line whatever may be the number of terms with which we start.

65. When the limits in the first integration are functions of the other variable we cannot perform the integrations in a different order, as in Art. 62, without special investigation to determine what the limits will then be. This question will be considered in a subsequent chapter.

66. From the definition of double integration, it follows that when the limits of both integrations are constant,

$$\iint \phi(x) \psi(y) dx dy = \int \phi(x) dx \times \int \psi(y) dy,$$

supposing that the limits in  $\int \psi(y) dy$  are the same as in the

integration with respect to  $y$  in the left hand member, and the limits in  $\int \phi(x) dx$  the same as in the integration with respect to  $x$  in the left hand member. For the left hand member is the limit of the sum of a series of terms, such as

$$h_{r+1} k_{s+1} \phi(x_r) \psi(y_s),$$

and the right hand member is the limit of the product of

$$h_1 \phi(x_0) + h_2 \phi(x_1) + h_3 \phi(x_2) \dots + h_n \phi(x_{n-1}),$$

and  $k_1 \psi(y_0) + k_2 \psi(y_1) + k_3 \psi(y_2) \dots + k_m \psi(y_{m-1})$ .

67. The reader will now be able to extend the processes given in this chapter to *triple* integrals and to *multiple* integrals generally. The symbol

$$\int_{\xi_0}^{\xi_1} \int_{\eta_0}^{\eta_1} \int_{\zeta_0}^{\zeta_1} \phi(x, y, z) dx dy dz$$


will indicate that the following series of operations must be performed—integrate  $\phi(x, y, z)$  with respect to  $z$  between the limits  $\zeta_0$  and  $\zeta_1$ ; next integrate the result with respect to  $y$  between the limits  $\eta_0$  and  $\eta_1$ ; lastly integrate this result with respect to  $x$  between the limits  $\xi_0$  and  $\xi_1$ . Here  $\zeta_0$  and  $\zeta_1$  may be functions of both  $x$  and  $y$ ; and  $\eta_0$  and  $\eta_1$  may be functions of  $x$ . This triple integral is the limit of a certain series which may be denoted by  $\Sigma \phi(x, y, z) \Delta x \Delta y \Delta z$ .

## CHAPTER VI.

## LENGTHS OF CURVES.

*Plane Curves. Rectangular co-ordinates.*

68. Let  $P$  be any point on the curve  $APQ$ , and let  $x$ ,  $y$  be its co-ordinates; let  $s$  denote the length of the arc  $AP$  measured from a fixed point  $A$  up to  $P$ ;



then, (*Dif. Cal.* Art. 307)

$$\frac{ds}{dx} = \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}}.$$

Hence,

$$s = \int \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}} dx.$$

From the equation to the curve we may express  $\frac{dy}{dx}$  in terms of  $x$ , and thus by integration  $s$  becomes known.

69. The process of finding the length of a curve is called the *rectification of the curve*, because we may suppose the question to be this: Find a *right line* equal in length to any assigned portion of the curve.

In the preceding article we have shewn that the length of an arc of a curve will be known if a certain integral can be obtained. It may happen in many cases that this integral cannot be obtained. Whenever the length of an arc of a curve can be expressed in terms of one or both of the co-ordinates of the variable extremity of the arc, the curve is said to be *rectifiable*.

70. *Application to the Parabola.*

The equation to the parabola is  $y = \sqrt{4ax}$ ; hence

$$\frac{dy}{dx} = \sqrt{\frac{a}{x}}, \quad \frac{ds}{dx} = \sqrt{\left(\frac{x+a}{x}\right)};$$

thus

$$s = \int \sqrt{\left(\frac{x+a}{x}\right)} dx \quad (\text{See Ex. 6, p. 18.})$$

$$= \sqrt{ax + x^2} + a \log \{\sqrt{x} + \sqrt{a+x}\} + C.$$

Here  $C$  denotes some *constant* quantity, that is, some quantity which does not depend upon  $x$ ; its value will depend upon the position of the fixed point from which the arc  $s$  is measured. If we measure from the vertex then  $s$  vanishes with  $x$ ; hence to determine  $C$  we have

$$a \log \sqrt{a} + C = 0;$$

and thus  $s = \sqrt{ax + x^2} + a \log \{\sqrt{x} + \sqrt{a+x}\} - a \log \sqrt{a}$

$$= \sqrt{ax + x^2} + a \log \frac{\sqrt{x} + \sqrt{a+x}}{\sqrt{a}}.$$

If then we require the length of the curve measured from the vertex to the point which has any assigned abscissa, we have only to put that assigned abscissa for  $x$  in the last expression. Thus, for example, for an extremity of the latus rectum  $x = a$ ; hence the length of the arc between the vertex and one extremity of the latus rectum is

$$a\sqrt{2} + a \log (1 + \sqrt{2}).$$

71. In the preceding article we have found the value of the constant  $C$ , but in applying the formula to ascertain the lengths of assigned portions of curves this is not necessary.

For suppose it required to find the length of the arc of a curve measured from the point whose abscissa is  $x_1$  up to the point whose abscissa is  $x_2$ . Let  $\psi(x)$  denote the integral of  $\sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}}$ , and let  $s_1$  and  $s_2$  be the lengths of arcs of the curve measured from any fixed point up to the points whose abscissæ are  $x_1$  and  $x_2$ , respectively, so that  $s_2 - s_1$  is the required length; then

$$s = \int \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}} dx = \psi(x) + C;$$

hence  $s_1 = \psi(x_1) + C; \quad s_2 = \psi(x_2) + C;$

therefore  $s_2 - s_1 = \psi(x_2) - \psi(x_1).$

Hence to find the required length we have to put  $x_1$  and  $x_2$  successively for  $x$  in  $\psi(x)$  and subtract the first result from the second. Thus we need not take any notice of the constant  $C$ ; in fact our result may be written

$$s_2 - s_1 = \int_{x_1}^{x_2} \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}} dx.$$

### 72. Application to the Cycloid.

In the cycloid, if the origin be at the vertex and the axis of  $y$  the tangent at that point, we have (*Dif. Cal. Art. 358*)

$$\frac{ds}{dx} = \sqrt{\left(\frac{2a}{x}\right)},$$

therefore,  $s = \sqrt{8ax} + C.$

The constant will be zero if we measure the arc  $s$  from the vertex.

### 73. Application to the Catenary.

The equation to the catenary is  $y = \frac{c}{2} (e^{\frac{x}{c}} + e^{-\frac{x}{c}})$ ; hence

$$\frac{dy}{dx} = \frac{1}{2} (e^{\frac{x}{c}} - e^{-\frac{x}{c}}), \quad \frac{ds}{dx} = \frac{1}{2} (e^{\frac{x}{c}} + e^{-\frac{x}{c}});$$

thus  $s = \frac{1}{2} \int (e^{\frac{x}{c}} + e^{-\frac{x}{c}}) dx = \frac{c}{2} (e^{\frac{x}{c}} - e^{-\frac{x}{c}}) + C.$

The constant will be zero if we measure the arc  $s$  from the point for which  $x = 0$ .

74. *Application to the Curve given by the equation*

$$x^{\frac{3}{2}} + y^{\frac{3}{2}} = a^{\frac{3}{2}}.$$

Here 
$$\frac{dy}{dx} = -\frac{y^{\frac{1}{2}}}{x^{\frac{1}{2}}}, \quad \frac{ds}{dx} = \left( \frac{x^{\frac{3}{2}} + y^{\frac{3}{2}}}{x^{\frac{3}{2}}} \right)^{\frac{1}{2}} = \frac{a^{\frac{3}{2}}}{x^{\frac{3}{2}}},$$

thus 
$$s = a^{\frac{3}{2}} \int \frac{dx}{x^{\frac{3}{2}}} = \frac{3a^{\frac{3}{2}}x^{\frac{1}{2}}}{2} + C.$$

The constant will be zero if we measure the arc from the point for which  $x = 0$ . The curve is an hypocycloid in which the radius of the revolving circle is one-fourth of the radius of the fixed circle. (See *Dif. Cal. Art. 360*, and put  $b = \frac{a}{4}$ ).

75. Since

$$\begin{aligned} \int \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}} dx &= \int \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}} \frac{dx}{dy} dy \\ &= \int \sqrt{\left\{1 + \left(\frac{dx}{dy}\right)^2\right\}} dy, \end{aligned}$$

we have 
$$s = \int \sqrt{\left\{1 + \left(\frac{dx}{dy}\right)^2\right\}} dy.$$

From the equation to the curve we may express  $\frac{dx}{dy}$  in terms of  $y$ , and thus by integration  $s$  becomes known. In some cases this formula may be more convenient than that in *Art. 68*.

76. *Application to the Logarithmic Curve.*

The equation to this curve is  $y = ba^x$ , or  $y = be^{\frac{x}{c}}$  if we suppose  $a = e^{\frac{1}{c}}$ ; thus  $x = c \log \frac{y}{b}$ ,

therefore  $\frac{dx}{dy} = \frac{c}{y}$ ,  $\frac{ds}{dy} = \frac{\sqrt{(c^2 + y^2)}}{y}$ ,

and  $s = \int \frac{\sqrt{(c^2 + y^2)}}{y} dy = \int \frac{c^2 dy}{y \sqrt{(c^2 + y^2)}} + \int \frac{y dy}{\sqrt{(c^2 + y^2)}}.$

The latter integral is  $\sqrt{(c^2 + y^2)}$ ; the former is

$$c \log \frac{y}{c + \sqrt{(c^2 + y^2)}}, \quad (\text{Art. 14}).$$

Hence  $s = c \log \frac{y}{c + \sqrt{(c^2 + y^2)}} + \sqrt{(c^2 + y^2)} + C.$

77. If  $x$  and  $y$  are each functions of a third variable  $t$ , we have (*Dif. Cal.* Art. 307),

$$\frac{ds}{dt} = \sqrt{\left\{ \left( \frac{dx}{dt} \right)^2 + \left( \frac{dy}{dt} \right)^2 \right\}};$$

thus  $s = \int \sqrt{\left\{ \left( \frac{dx}{dt} \right)^2 + \left( \frac{dy}{dt} \right)^2 \right\}} dt.$

78. The equation to the ellipse is  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ . We may therefore assume  $x = a \sin \phi$ ,  $y = b \cos \phi$ , so that  $\phi$  is the complement of the *excentric angle*, (*Plane Co-ordinate Geometry*, Art. 168). Therefore, by the preceding article,

$$\frac{ds}{d\phi} = \sqrt{(a^2 \cos^2 \phi + b^2 \sin^2 \phi)},$$

and  $s = \int \sqrt{(a^2 \cos^2 \phi + b^2 \sin^2 \phi)} d\phi = a \int \sqrt{(1 - e^2 \sin^2 \phi)} d\phi.$

The exact integral cannot be obtained; we may however expand  $\sqrt{(1 - e^2 \sin^2 \phi)}$  in a series, so that

$$s = a \int \left\{ 1 - \frac{1}{2} e^2 \sin^2 \phi - \frac{1.1}{2.4} e^4 \sin^4 \phi - \frac{1.1.3}{2.4.6} e^6 \sin^6 \phi \dots \right\} d\phi$$

and each term can be integrated separately. To obtain the length of the elliptic quadrant we must integrate between the limits 0 and  $\frac{\pi}{2}$ .



*Plane Curves. Polar Co-ordinates.*

79. Let  $r$ ,  $\theta$  be the polar co-ordinates of any point of a curve, and  $s$  the length of the arc measured from any fixed point up to this point; then (*Dif. Cal.* Art. 311)

$$\frac{ds}{d\theta} = \sqrt{\left\{r^2 + \left(\frac{dr}{d\theta}\right)^2\right\}};$$

hence 
$$s = \int \sqrt{\left\{r^2 + \left(\frac{dr}{d\theta}\right)^2\right\}} d\theta.$$

80. *Application to the Spiral of Archimedes.*

In this curve  $r = a\theta$ , thus  $\frac{dr}{d\theta} = a$ ;

hence 
$$s = \int \sqrt{(r^2 + a^2)} d\theta = a \int \sqrt{(1 + \theta^2)} d\theta$$

$$= \frac{a\theta}{2} \sqrt{(1 + \theta^2)} + \frac{a}{2} \log \{\theta + \sqrt{(1 + \theta^2)}\} + C.$$

The constant will be zero if we measure the arc  $s$  from the pole, that is from the point where  $\theta = 0$ .

81. *Application to the Cardioid.*

The equation to this curve is  $r = a(1 + \cos \theta)$ ; thus

$$s = \int \sqrt{\{a^2(1 + \cos \theta)^2 + a^2 \sin^2 \theta\}} d\theta = a \int \sqrt{(2 + 2 \cos \theta)} d\theta$$

$$= 2a \int \cos \frac{\theta}{2} d\theta = 4a \sin \frac{\theta}{2} + C.$$

The constant will be zero if we measure the arc  $s$  from the point for which  $\theta = 0$ , that is, from the point where the curve crosses the initial line.

The length of that part of the curve which is comprised between the initial line and a line through the pole at right angles to the initial line is  $4a \sin \frac{\pi}{4}$ . The length of half the perimeter of the curve is  $4a \sin \frac{\pi}{2}$ , that is,  $4a$ .

82. Suppose we require the length of the complete perimeter of the cardioid; we might at first suppose that it would be equal to  $2a \int_0^{2\pi} \cos \frac{\theta}{2} d\theta$ ; but this would give zero as the result, which is obviously inadmissible. The reason of this may be easily seen; we have in fact shewn that

$$\frac{ds}{d\theta} = a \sqrt{2 + 2 \cos \theta},$$

and this ought not to be put equal to  $2a \cos \frac{\theta}{2}$  but to  $\pm 2a \cos \frac{\theta}{2}$ ,

and the proper sign should be determined in any application of the formula. Now by  $s$  we understand a positive quantity, and we may measure  $s$  so that it increases with  $\theta$ , and thus

$\frac{ds}{d\theta}$  is positive. Hence when  $\cos \frac{\theta}{2}$  is positive, we take the

upper sign and put  $\frac{ds}{d\theta} = 2a \cos \frac{\theta}{2}$ ; when  $\cos \frac{\theta}{2}$  is negative, we

take the lower sign and put  $\frac{ds}{d\theta} = -2a \cos \frac{\theta}{2}$ . Hence the

length of the complete perimeter is not  $2a \int_0^{2\pi} \cos \frac{\theta}{2} d\theta$ , but

$2a \int_0^{\pi} \cos \frac{\theta}{2} d\theta - 2a \int_{\pi}^{2\pi} \cos \frac{\theta}{2} d\theta$ , that is,  $8a$ . This result might

have been anticipated, for it will be obvious from the symmetry of the figure that the length of the complete perimeter is double the length of the part which is situated on one side of the initial line, and this was shewn to be  $4a$  in the preceding article.

83. It may sometimes be more convenient to find the length of a curve from the formula

$$s = \int \sqrt{\left\{ r^2 \left( \frac{d\theta}{dr} \right)^2 + 1 \right\}} dr,$$

which follows immediately from that in Art. 79.

84. *Application to the Logarithmic Spiral.*

The equation to this curve is  $r = ba^\theta$ , or  $r = be^{\frac{\theta}{a}}$  if we suppose  $a = e^{\frac{1}{a}}$ ; thus  $\theta = c \log \frac{r}{b}$ ; therefore  $\frac{d\theta}{dr} = \frac{c}{r}$  and

$$s = \int \sqrt{(1 + c^2)} dr = \sqrt{(1 + c^2)} r + C.$$

Thus the length of the portion of the curve which has  $r_1$  and  $r_2$  for the radii vectores of its extreme points is

$$\int_{r_1}^{r_2} \sqrt{(1 + c^2)} dr, \text{ that is, } \sqrt{(1 + c^2)} (r_2 - r_1).$$

The angle between the radius vector and the corresponding tangent at any point of this curve is constant, (*Dif. Cal. Art. 354*); and if that angle be denoted by  $\alpha$  we have  $c = \tan \alpha$ ; thus  $\sqrt{(1 + c^2)} = \sec \alpha$ ,  $\frac{ds}{dr} = \sec \alpha$ , and  $s = r \sec \alpha + C$ . Hence  $(r_2 - r_1) \sec \alpha$  is the length of the portion mentioned above.

*Formulae involving the radius vector and perpendicular.*

85. Let  $\phi$  be the angle between the radius vector  $r$  of any point of a curve and the tangent at that point; then  $\cos \phi = \frac{dr}{ds}$ , (*Dif. Cal. Art. 310*). Let  $p$  be the perpendicular from the pole on the same tangent; then

$$\sin \phi = \frac{p}{r}, \text{ therefore } \cos \phi = \frac{\sqrt{(r^2 - p^2)}}{r};$$

thus 
$$\frac{dr}{ds} = \frac{\sqrt{(r^2 - p^2)}}{r};$$

therefore 
$$\frac{ds}{dr} = \frac{r}{\sqrt{(r^2 - p^2)}},$$

and 
$$s = \int \frac{r dr}{\sqrt{(r^2 - p^2)}}.$$

86. *Application to the Epicycloid.*

With the notation and figure in *Dif. Cal.* Art. 360, it may be shewn that the equation to the tangent to the epicycloid at  $P$  is

$$y' - y = - \frac{\cos \theta - \cos \frac{a+b}{b} \theta}{\sin \theta - \sin \frac{a+b}{b} \theta} (x' - x),$$

where  $x$  and  $y$  are the co-ordinates of  $P$ , and  $x'$  and  $y'$  the variable co-ordinates. Hence it will be found that the perpendicular  $p$  from the origin on the tangent at  $P$  is given by

$$p = (a + 2b) \sin \frac{a\theta}{2b};$$

also 
$$r^2 = a^2 + 4b(a+b) \sin^2 \frac{a\theta}{2b};$$

thus 
$$p^2 = \frac{c^2(r^2 - a^2)}{c^2 - a^2} \text{ where } c = a + 2b.$$

Hence, by Art. 85,

$$s = \frac{\sqrt{(c^2 - a^2)}}{a} \int \frac{r dr}{\sqrt{(c^2 - r^2)}} = - \frac{\sqrt{(c^2 - a^2)}}{a} \sqrt{(c^2 - r^2)} + C.$$

At a cusp  $r = a$ , and at a vertex  $r = c$ ; thus the length of the portion of the curve between a cusp and the adjacent vertex is

$$\frac{\sqrt{(c^2 - a^2)}}{a} \int_a^c \frac{r dr}{\sqrt{(c^2 - r^2)}}, \text{ that is } \frac{c^2 - a^2}{a}, \text{ that is } \frac{4b(a+b)}{a}.$$

Hence the length of the portion between two consecutive cusps is  $\frac{8b(a+b)}{a}$ .

87. A remark may be made here similar to that in Art. 82. If we apply the formula

$$s = - \frac{\sqrt{(c^2 - a^2)}}{a} \sqrt{(c^2 - r^2)} + C$$

to find the length between two consecutive cusps, we arrive at the result zero, since  $r = a$  at both limits. The reason is that we have used the formula

$$\frac{ds}{dr} = \frac{\sqrt{(c^2 - a^2)}}{a} \frac{r}{\sqrt{(c^2 - r^2)}}$$

while the true formula is

$$\frac{ds}{dr} = \pm \frac{\sqrt{(c^2 - a^2)}}{a} \frac{r}{\sqrt{(c^2 - r^2)}}.$$

Since  $s$  may be taken to increase continually, it follows that  $\frac{ds}{dr}$  is positive when  $r$  is increasing, and negative when  $r$  is diminishing. Now in passing along the curve from a cusp to the adjacent vertex  $r$  increases, thus  $\frac{ds}{dr}$  is positive, and we should take the *upper* sign in the formula for  $\frac{ds}{dr}$ ; then in passing from the vertex to the next cusp  $r$  diminishes, thus  $\frac{ds}{dr}$  is negative and the *lower* sign must be taken. Hence the length from one cusp to the next cusp is

$$\begin{aligned} &= \frac{\sqrt{(c^2 - a^2)}}{a} \int_a^c \frac{r dr}{\sqrt{(c^2 - r^2)}} - \frac{\sqrt{(c^2 - a^2)}}{a} \int_c^a \frac{r dr}{\sqrt{(c^2 - r^2)}} \\ &= \frac{2\sqrt{(c^2 - a^2)}}{a} \int_a^c \frac{r dr}{\sqrt{(c^2 - r^2)}} = \frac{8b(a + b)}{a}. \end{aligned}$$

88. From what is stated in the preceding article, it appears that if the arc  $s$  begin at a vertex the proper formula is

$$\frac{ds}{dr} = -\frac{\sqrt{(c^2 - a^2)}}{a} \frac{r}{\sqrt{(c^2 - r^2)}},$$

therefore  $s = -\frac{\sqrt{(c^2 - a^2)}}{a} \int \frac{r dr}{\sqrt{(c^2 - r^2)}} = \frac{\sqrt{(c^2 - a^2)}}{a} \sqrt{(c^2 - r^2)}$ .

No constant is required since we begin to measure at the point for which  $r = c$ ; the formula holds for values of  $s$  less than  $\frac{4b(a + b)}{a}$ .

It may be observed that thus

$$s = \frac{c^2 - a^2}{a^2} \sqrt{(r^2 - p^2)}.$$

89. Similarly for the hypocycloid we may shew that

$$p^2 = \frac{c^2(a^2 - r^2)}{a^2 - c^2} \text{ where } c = a - 2b.$$

Suppose  $c^2$  less than  $a^2$ ; then we may shew that

$$\frac{ds}{dr} = \pm \frac{\sqrt{(a^2 - c^2)}}{a} \frac{r}{\sqrt{(r^2 - c^2)}},$$

and thus  $s$  may be found. The length of the curve between two adjacent cusps is  $\frac{8b(a-b)}{a}$ .

Next suppose  $c^2$  greater than  $a^2$ ; then we should write the value of  $\frac{ds}{dr}$  thus

$$\frac{ds}{dr} = \pm \frac{\sqrt{(c^2 - a^2)}}{a} \frac{r}{\sqrt{(c^2 - r^2)}};$$

in this case  $b$  is greater than  $a$ , and we shall find the length of the curve between two adjacent cusps to be  $\frac{8b(b-a)}{a}$ .

When  $a = 2b$  we have  $c = 0$  and  $p = 0$ ; in this case the hypocycloid becomes a straight line coinciding with a diameter of the fixed circle.

If  $a = b$  we have  $c^2 = a^2$ ; in this case the denominator in the value of  $p^2$  vanishes; it will be found that the hypocycloid is then reduced to a point, and  $r = a$ .


It may be shewn as in Art. 88, that if  $s$  be measured from a vertex to a point not beyond the adjacent cusp, we have

$$s = \pm \frac{c^2 - a^2}{a^2} \sqrt{(r^2 - p^2)},$$

the upper or lower sign being taken according as  $c$  is greater or less than  $a$ .

*Formulae involving the Perpendicular and its Inclination.*

90. Another method of expressing the length of a curve is worthy of notice.



Let  $P$  be a point in a curve;  $x, y$  its co-ordinates. Let  $s$  be the length of the arc measured from a fixed point  $A$  up to  $P$ . Draw  $OY$  a perpendicular from the origin  $O$  on the tangent at  $P$ ; suppose  $OY = p$ ,  $PY = u$ ,  $YOx = \theta$ ; then

$$p = x \cos \theta + y \sin \theta,$$

$$u = x \sin \theta - y \cos \theta,$$

$$\frac{dy}{dx} = -\cot \theta, \quad \frac{ds}{dx} = -\operatorname{cosec} \theta;$$

therefore

$$\frac{dp}{d\theta} = -x \sin \theta + y \cos \theta + \cos \theta \frac{dx}{d\theta} + \sin \theta \frac{dy}{d\theta} = -u,$$

$$\frac{d^2p}{d\theta^2} = -\frac{du}{d\theta} = -x \cos \theta - y \sin \theta - \sin \theta \frac{dx}{d\theta} + \cos \theta \frac{dy}{d\theta}$$

$$= -p - \operatorname{cosec} \theta \frac{dx}{d\theta} = -p + \frac{ds}{d\theta};$$

therefore, by integration,

$$\frac{dp}{d\theta} = -\int p d\theta + s,$$

therefore 
$$s = \frac{dp}{d\theta} + \int p d\theta;$$

this may also be written

$$s + u = \int p d\theta.$$

Suppose  $s_1$  and  $u_1$  the values of  $s$  and  $u$  when  $\theta$  has the value  $\theta_1$ , and  $s_2$  and  $u_2$  their values when  $\theta$  has the value  $\theta_2$ , then

$$s_2 - s_1 + u_2 - u_1 = \int_{\theta_1}^{\theta_2} p d\theta.$$

We have measured  $u$  in the direction of revolution from  $P$  and have taken it as positive in this case; when  $u$  is negative it will indicate that  $Y$  is on the other side of  $P$ .

91. The preceding article may be used for different purposes, among which two may be noticed.

(1) To determine the length of any portion of a curve when the equation to the curve is given; for from that equation together with  $\frac{dy}{dx} = -\cot \theta$  we can find  $x$  and  $y$  in terms of  $\theta$ , and therefore  $p$  which is equal to  $x \cos \theta + y \sin \theta$ ; then  $s$  may be found from the equation

$$s = \frac{dp}{d\theta} + \int p d\theta.$$

(2) To find a curve such that by means of its arc a proposed integral may be represented; for if the proposed integral be  $\int p d\theta$  where  $p$  is a function of  $\theta$ , the required curve is found by eliminating  $\theta$  between the equations


$$x = p \cos \theta - \frac{dp}{d\theta} \sin \theta, \quad y = p \sin \theta + \frac{dp}{d\theta} \cos \theta,$$

and then the integral may be represented by  $s - \frac{dp}{d\theta}$ .

Arts. 90 and 91 have been derived from Hymers's *Integral Calculus*, Art. 136.



92. *Application to the Ellipse.*



Let  $APB$  be a quadrant of an ellipse,  $CY$  the perpendicular on the tangent at  $P$ ; let  $ACY = \theta$ . Then (*Plane Coordinate Geometry*, Art. 196),  $CY = a\sqrt{1 - e^2 \sin^2 \theta}$ ;

therefore  $AP + PY = a \int \sqrt{1 - e^2 \sin^2 \theta} d\theta$ ,

the constant to be added to the integral is supposed to be so taken that the integral may vanish with  $\theta$ . If  $R$  be a point such that its excentric angle is  $\frac{\pi}{2} - \theta$ , we have by Art. 78,

$$BR = a \int \sqrt{1 - e^2 \sin^2 \theta} d\theta ;$$

thus  $AP + PY = BR \dots \dots \dots (1)$ .

And  $PY = -\frac{dp}{d\theta} = \frac{ae^2 \sin \theta \cos \theta}{\sqrt{1 - e^2 \sin^2 \theta}}$ .

Let  $x$  be the abscissa of  $P$ ; then by Art. 90,

$$x = p \cos \theta - \frac{dp}{d\theta} \sin \theta$$

$$= a\sqrt{1 - e^2 \sin^2 \theta} \cos \theta + \frac{ae^2 \sin^2 \theta \cos \theta}{\sqrt{1 - e^2 \sin^2 \theta}} = \frac{a \cos \theta}{\sqrt{1 - e^2 \sin^2 \theta}}.$$

Thus  $PY = e^2 x \sin \theta$ ; and if  $x'$  be the abscissa of  $R$  we have  $x' = a \cos (\frac{\pi}{2} - \theta)$  so that  $PY = \frac{e^2 x x'}{a}$ . Thus (1) may be written

$$BR - AP = \frac{e^2}{a} x x' \dots \dots \dots (2);$$

this result is called Fagnani's Theorem.

From the ascertained values of  $x$  and  $x'$  we have

$$x^2 = \frac{a^2 - a^2 \sin^2 \theta}{1 - e^2 \sin^2 \theta} = \frac{a^2 - x'^2}{1 - \frac{e^2 x'^2}{a^2}};$$


therefore  $e^2 x^2 x'^2 - a^2 (x^2 + x'^2) + a^4 = 0$ .

Thus the equation which connects  $x$  and  $x'$  involves these quantities *symmetrically*; hence from (2) we can infer that

$$BP - AR = \frac{e^2}{a} xx'.$$

This is also obvious from the figure.

93. *Application to the Hyperbola.*



Let  $C$  be the centre and  $A$  the vertex of an hyperbola,  $CY$  the perpendicular on the tangent at  $P$ . Let  $\angle ACY = \theta$  and  $CY = p$ ; then it may be proved that

$$PY - AP = a \int \sqrt{(1 - e^2 \sin^2 \theta)} d\theta;$$

this may be proved in the same manner as the corresponding result of the preceding article; we may either make the requisite changes of sign in the formulæ of Art. 90, which are produced by difference of figure; or may begin from the beginning again in the manner of that article. The constant to be added to the integral is supposed to be so taken that the integral may vanish with  $\theta$ .

Suppose  $\alpha$  the greatest value which  $\theta$  can have, then (*Plane Co-ordinate Geometry*, Art. 257)  $\cot \alpha = \sqrt{e^2 - 1}$ . When  $P$  moves off to an infinite distance  $PY - AP$  becomes the difference between the length of the asymptote from  $C$  and the infinite hyperbolic arc from  $A$ . Thus this difference is

$$\alpha \int_0^{\alpha} \sqrt{1 - e^2 \sin^2 \theta} d\theta.$$

*Inverse questions on the lengths of Curves.*

94. In the preceding articles we have shewn how the length of an arc of a known curve is to be found in terms of the abscissa of its variable extremity; we will now briefly notice the inverse problem—to find a curve such that the arc shall be a given function of the abscissa of its variable extremity.

Suppose  $\phi(x)$  the given function; then  $s = \phi(x)$ ;

therefore 
$$\phi'(x) = \frac{ds}{dx} = \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}};$$

thus 
$$\frac{dy}{dx} = [(\phi'(x))^2 - 1]^{\frac{1}{2}},$$

and 
$$y = \int [(\phi'(x))^2 - 1]^{\frac{1}{2}} dx.$$

95. As an example of the preceding method, suppose  $\phi(x) = \sqrt{4cx}$ ; thus  $\phi'(x) = \sqrt{\frac{c}{x}}$ ; therefore

$$\begin{aligned} y &= \int \left[ \frac{c}{x} - 1 \right]^{\frac{1}{2}} dx = \int \frac{(c-x) dx}{\sqrt{(cx-x^2)}} \\ &= \int \frac{\left(\frac{c}{2} - x\right) dx}{\sqrt{(cx-x^2)}} + \frac{c}{2} \int \frac{dx}{\sqrt{(cx-x^2)}} \\ &= \sqrt{(cx-x^2)} + \frac{c}{2} \text{vers}^{-1} \frac{2x}{c} + C. \end{aligned}$$

We may write  $y'$  for  $y - C$  and thus we find that the curve is a cycloid. (*Dif. Cal.* Art. 358.)

96. For another example suppose  $\phi(x) = a \log x$ ; thus  $\phi'(x) = \frac{a}{x}$ .

Here 
$$y = \int \sqrt{\left(\frac{a^2}{x^2} - 1\right)} dx = \int \frac{(a^2 - x^2) dx}{x \sqrt{(a^2 - x^2)}}$$

$$= \int \frac{a^2 dx}{x \sqrt{(a^2 - x^2)}} - \int \frac{x dx}{\sqrt{(a^2 - x^2)}}$$

$$= a \log \frac{x}{a + \sqrt{(a^2 - x^2)}} + \sqrt{(a^2 - x^2)} + C.$$

*Involutes and Evolutes.*

97. We may express the length of an arc of a curve without integration when we know the equation to the involute of the curve. Suppose  $s'$  to represent the length of an arc of a curve,  $\rho$  the radius of curvature at that point of the involute which corresponds to the variable extremity of  $s'$ , then (*Dif. Cal. Art. 331*)  $s' \pm \rho = l$ , where  $l$  is a constant. If the equation to the involute is known,  $\rho$  can be found in terms of the co-ordinates of the point in the involute; then these co-ordinates can be expressed in terms of the co-ordinates of the corresponding point of the evolute, and thus  $s'$  is known. By this method we have to perform the processes of differentiation and algebraical reduction instead of integration.

98. *Application to the Evolute of the Parabola.*

Take for the involute the parabola which has for its equation  $y^2 = 4ax$ ; let  $x', y'$  be the co-ordinates of the point of the evolute which corresponds to the point  $(x, y)$  on the parabola. Then by the ordinary methods (*Dif. Cal. Art. 330*) we have

$$x' = 2a + 3x, \quad y' = -\frac{y^3}{4a^2},$$

and

$$\rho = 2a \left(\frac{a+x}{a}\right)^{\frac{3}{2}}.$$

Thus we shall obtain for the equation to the evolute

$$27ay'^2 = 4(x' - 2a)^3;$$

and

$$\rho = 2a \left(\frac{x' + a}{3a}\right)^{\frac{3}{2}};$$

therefore  $s' \pm 2a \left( \frac{x' + a}{3a} \right)^{\frac{2}{3}} = l.$

Suppose we measure  $s'$  from the point for which  $x' = 2a$ , that is from the point which corresponds to the vertex of the parabola; then we see that  $s'$  increases with  $x'$ , so that we must take the lower sign in the last equation; also by supposing  $x' = 2a$  and  $s' = 0$  we find  $l = -2a$ ; thus

$$s' = 2a \left( \frac{x' + a}{3a} \right)^{\frac{2}{3}} - 2a.$$

This value of  $s'$  may also be obtained by the application of the ordinary method of integration.

99. When the length of the arc of a curve is known in terms of the co-ordinates of its variable extremity, the equation to the involute can be found by ordinary processes of elimination.

For we have (*Dif. Cal.* Art. 331),

$$\frac{\frac{dx'}{dx}}{x' - x} = \frac{1}{\rho} \frac{ds'}{dx}$$

where the accented letters refer to a point in a curve, and the unaccented letters to the corresponding point in the involute. Thus

$$x = x' - \rho \frac{dx'}{ds'} \dots\dots\dots(1).$$

Similarly  $y = y' - \rho \frac{dy'}{ds'} \dots\dots\dots(2).$

If then  $s'$  is known in terms of  $x'$ , or of  $y'$ , or of both, by means of this relation and the known equation to the curve we may find  $\frac{dx'}{ds'}$  and  $\frac{dy'}{ds'}$ ; and  $\rho$  is known from the equation  $s' \pm \rho = l.$  It only remains then to eliminate  $x'$  and  $y'$  from (1) and (2) and the known equation to the curve; we obtain thus an equation between  $x$  and  $y$ , which is the required equation to the involute.



100. *Application to the Catenary.*

The equation to the Catenary is

$$y' = \frac{c}{2} \left( e^{\frac{x'}{c}} + e^{-\frac{x'}{c}} \right),$$

and

$$s' = \frac{c}{2} \left( e^{\frac{x'}{c}} - e^{-\frac{x'}{c}} \right),$$

supposing  $s'$  measured from the point for which  $x' = 0$  and  $y' = c$ ; we shall now find the equation to that involute to the catenary which begins at the point of the curve just specified.

We have then

$$\frac{dy'}{dx'} = \frac{s'}{c}, \quad \frac{ds'}{dx'} = \frac{y'}{c};$$

thus

$$\frac{dy'}{ds'} = \frac{s'}{y'}, \quad \frac{dx'}{ds'} = \frac{c}{y'};$$

and  $\rho = s'$ , no constant being required, because by supposition  $\rho$  vanishes with  $s'$ .

Hence equations (1) and (2) of the preceding article become

$$x = x' - \frac{s'c}{y'};$$

$$y = y' - \frac{s'^2}{y'} = \frac{y'^2 - s'^2}{y'} = \frac{c^2}{y'}.$$

And

$$s' = \sqrt{y'^2 - c^2} = \sqrt{\left(\frac{c^2}{y'} - c^2\right)} = \frac{c}{y'} \sqrt{c^2 - y'^2};$$

therefore

$$\frac{s'}{y'} = \frac{\sqrt{c^2 - y'^2}}{c};$$

thus  $x = x' - \sqrt{c^2 - y'^2}$ ; therefore  $x' = \sqrt{c^2 - y'^2} + x$ .

We have then to substitute these values of  $x'$  and  $y'$  in the equation to the catenary, and thus obtain the required relation between  $x$  and  $y$ . The substitution may be conveniently performed thus

$$y' = \frac{c}{2} \left( e^{\frac{x}{c}} + e^{-\frac{x}{c}} \right);$$

therefore  $\sqrt{(y'^2 - c^2)} = \frac{c}{2} \left( e^{\frac{x}{c}} - e^{-\frac{x}{c}} \right);$

therefore  $y' + \sqrt{(y'^2 - c^2)} = ce^{\frac{x}{c}},$

therefore  $x' = c \log \frac{y' + \sqrt{(y'^2 - c^2)}}{c}.$

Thus finally,  $x + \sqrt{(c^2 - y^2)} = c \log \frac{c + \sqrt{(c^2 - y^2)}}{y}.$

This curve is called the *tractory*; on account of the radical, there are two values of  $x$  for every value of  $y$  less than  $c$ , these two values being numerically equal, but of opposite signs. There is a cusp at the point for which  $x = 0$  and  $y = c$ ; and the axis of  $x$  is an asymptote.

101. The polar formulæ may also be used in like manner to determine the involute when the length of an arc of the evolute can be expressed in terms of the polar co-ordinates of its variable extremity. We have (*Dif. Cal.* Art. 332),

$$r'^2 = \rho^2 + r^2 - 2\rho p \dots \dots \dots (1),$$

$$p'^2 = r^2 - p^2 \dots \dots \dots (2).$$

Here, as before, the accented letters belong to the known curve, that is, to the evolute, and the unaccented letters to the required involute; thus since the evolute is known, there is a known relation between  $p'$  and  $r'$ . And  $s' \pm p = l$ , so that if  $s'$  can be expressed in terms of  $p'$  and  $r'$  we may eliminate  $p'$  and  $r'$  by means of (1), (2), and the known equation to the evolute. Thus we obtain an equation connecting  $p$  and  $r$ , which serves to determine the involute.

### 102. *Application to the Equiangular Spiral.*

In this curve  $p' = r' \sin \alpha$ , where  $\alpha$  is the constant angle of the spiral. If we suppose the involute to begin from the

pole of the spiral, and  $s'$  to be measured from that point, we have  $\rho = s' = r' \sec \alpha$ , (Art. 84). Thus (1) of the preceding article becomes

$$\begin{aligned} r'^2 &= r'^2 \sec^2 \alpha + r'^2 - 2r'p \sec \alpha \\ &= r'^2 \sec^2 \alpha + r'^2 \sin^2 \alpha + p^2 - 2r'p \sec \alpha, \text{ by (2).} \end{aligned}$$

From this quadratic for  $p$  we obtain

$$p - r' \sec \alpha = \pm r' \cos \alpha.$$

If we take the upper sign we find  $p = \frac{r'(1 + \cos^2 \alpha)}{\cos \alpha}$ , and then from (2) we find  $r'^2 = \frac{1 + 3 \cos^2 \alpha}{\cos^2 \alpha} r'^2$ . But this solution must be rejected, because from it we should find  $\rho$  or  $r' \frac{dr}{dp} = \frac{1 + 3 \cos^2 \alpha}{\cos \alpha (1 + \cos^2 \alpha)} r'$ , which is inconsistent with the equation  $\rho = r' \sec \alpha$ .

If we take the lower sign we find  $p = \frac{r' \sin^2 \alpha}{\cos \alpha}$ , and then from (2) we find  $r'^2 = \frac{r'^2 \sin^2 \alpha}{\cos^2 \alpha}$ ; thus  $p = r' \sin \alpha$ . Hence the involute is an equiangular spiral with the same constant angle as the evolute has.

### *Intrinsic Equation to a Curve.*

103. Let  $s$  denote the length of an arc of a curve measured from some fixed point,  $\phi$  the inclination of the tangent at the variable extremity to the tangent at some fixed point of the curve; then the equation which determines the relation between  $s$  and  $\phi$  is called the *intrinsic equation* to the curve. In some investigations, especially those relating to involutes and evolutes, this method of determining a curve is simpler than the ordinary method of referring the curve to rectangular axes which are *extrinsic* lines.



104. We will first shew how the *intrinsic* equation may be obtained from the ordinary equation.

Suppose  $y = f(x)$  the equation to a curve, the origin being a point on the curve, and the axis of  $y$  a tangent at that point; from the given equation we have

$$\frac{dy}{dx} = f'(x) = \frac{1}{\tan \phi} \text{ by hypothesis;}$$

thus  $x$  is known in terms of  $\tan \phi$ , say  $x = F(\tan \phi)$ ; then

$$\frac{dx}{d\phi} = F'(\tan \phi) \sec^2 \phi;$$

also 
$$\frac{ds}{dx} = \operatorname{cosec} \phi;$$

therefore 
$$\frac{ds}{d\phi} = F'(\tan \phi) \sec^2 \phi \operatorname{cosec} \phi;$$

from this equation  $s$  may be found in terms of  $\phi$  by integration. A similar result will be obtained if at the origin the axis of  $x$  be the axis which we suppose to coincide with a tangent.

### 105. *Application to the Cycloid.*

By *Dif. Cal.* Art. 358, we have

$$\frac{dy}{dx} = \sqrt{\left(\frac{2a-x}{x}\right)} = \frac{1}{\tan \phi};$$

therefore 
$$\frac{2a}{x} = \frac{1}{\sin^2 \phi}, \quad x = 2a \sin^2 \phi,$$

$$\frac{dx}{d\phi} = 4a \sin \phi \cos \phi,$$

$$\frac{ds}{d\phi} = \operatorname{cosec} \phi \frac{dx}{d\phi} = 4a \cos \phi;$$

therefore 
$$s = 4a \sin \phi + C.$$

The constant will be zero if we suppose  $s$  measured from the fixed point where the first tangent is drawn, that is, from the vertex of the curve.

106. Having given the intrinsic equation to deduce the ordinary equation.

We have 
$$\frac{dx}{ds} = \sin \phi ;$$

therefore 
$$x = \int ds \sin \phi .$$

Similarly 
$$y = \int ds \cos \phi .$$

Now  $s$  is by supposition known in terms of  $\phi$ ; thus by integration we may find  $x$  and  $y$  in terms of  $\phi$ , and then by eliminating  $\phi$  we obtain the ordinary equation to the curve in terms of  $x$  and  $y$ .

107. *Application to the Cycloid.*

Here  $s = 4a \sin \phi ;$

thus 
$$x = \int ds \sin \phi = 4a \int \sin \phi \cos \phi d\phi = C - a \cos 2\phi ,$$

$$y = \int ds \cos \phi = 4a \int \cos^2 \phi d\phi = C' + 2a\phi + a \sin 2\phi .$$

Hence by eliminating  $\phi$  we can obtain the ordinary equation; if the origin of the rectangular axes is the vertex of the curve, we shall have  $C = a$  and  $C' = 0$ .

108. We shall now give some miscellaneous examples of intrinsic equations.

The intrinsic equation to the circle is obviously  $s = a\phi$ .

109. The equation to the catenary is

$$y + c = \frac{c}{2} (e^{\frac{x}{c}} + e^{-\frac{x}{c}}),$$

the origin being on the curve. Hence

$$\frac{dy}{dx} = \frac{1}{2} (e^{\frac{x}{c}} - e^{-\frac{x}{c}}), \quad s = \frac{c}{2} (e^{\frac{x}{c}} - e^{-\frac{x}{c}});$$

thus if  $\phi$  be the angle which the tangent at any point make with the tangent at the origin

$$s = c \tan \phi.$$

110. We have seen in Art. 86, that for the epicycloid

$$\frac{dy}{dx} = \frac{\cos \theta - \cos \frac{a+b}{b} \theta}{\sin \frac{a+b}{b} \theta - \sin \theta} = \tan \phi \text{ suppose,}$$

thus 
$$\phi = \frac{a+2b}{2b} \theta.$$

Again, from the same article,

$$\begin{aligned} s &= -\frac{\sqrt{(c^2 - a^2)}}{a} \sqrt{(c^2 - r^2)} + C \\ &= -\frac{4b(a+b)}{a} \cos \frac{a\theta}{2b} + C \\ &= \frac{4b(a+b)}{a} \left(1 - \cos \frac{a\theta}{2b}\right), \end{aligned}$$

if we suppose  $s$  measured from the point for which  $\theta = 0$ .

Thus 
$$s = \frac{4b(a+b)}{a} \left(1 - \cos \frac{a\phi}{a+2b}\right).$$

We may simplify this result by putting

$$\phi = \frac{\pi(a+2b)}{2a} + \phi', \quad \text{and } s = \frac{4b(a+b)}{a} + s';$$

this amounts to measuring the arc from a vertex instead of from a cusp. Thus

$$s' = \frac{4b(a+b)}{a} \sin \frac{a\phi'}{a+2b}$$

where the accent may now be dropped.

111. Similarly the intrinsic equation to the hypocycloid may be written

$$s = \frac{4b(a-b)}{a} \sin \frac{a\phi}{a-2b}.$$

112. It appears from the last two articles that  $s = l \sin n\phi$  represents an epicycloid or hypocycloid, according as  $n$  is less or greater than unity. For example, if

$$s = l \sin \frac{\phi}{2}, \quad s = l \sin \frac{\phi}{3}, \quad s = l \sin \frac{\phi}{4}, \quad s = l \sin \frac{\phi}{5}, \dots$$

we have epicycloids in which  $\frac{b}{a} = \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$

If  $s = l \sin 2\phi, s = l \sin 3\phi, s = l \sin 4\phi, s = l \sin 5\phi, \dots$

we have hypocycloids in which  $\frac{b}{a} = \frac{1}{4}, \frac{1}{3}, \frac{3}{8}, \frac{2}{5}, \dots$

113. If  $\rho$  be the radius of curvature of the curve at the point determined by  $s$  and  $\phi$ , we have (*Dif. Cal. Art. 324*),

$$\rho = \frac{ds}{d\phi}.$$

In the logarithmic spiral we know that  $\rho$  varies as  $s$  if the arc be measured from the pole; thus

$$\rho = ks = \frac{ds}{d\phi};$$

therefore


$$s = ae^{k\phi}$$

where  $a$  is a constant. If we put  $s = s' + a$  we have

$$s' = a(e^{k\phi} - 1),$$

and now  $s'$  is measured from the point for which  $\phi = 0$ .

114. If the intrinsic equation to a curve be known, that to the evolute can be found.



Let  $AP$  be a curve,  $BQ$  the evolute; let  $s$  be the length of an arc of  $AP$  measured from some fixed point up to  $P$ ;  $s'$  the length of an arc of  $BQ$  measured from some fixed point up to  $Q$ . It is evident that  $\phi$  is the same both for  $s$  and  $s'$ , if in  $BQ$  we measure  $\phi$  from  $BA$ , which is perpendicular to the line from which  $\phi$  is measured in  $AP$ .

$$\text{In the left-hand figure } s' = \rho - C = \frac{ds}{d\phi} - C.$$

$$\text{In the right-hand figure } s' = C - \rho = C - \frac{ds}{d\phi}.$$

Thus if  $s$  be known in terms of  $\phi$ , we can find  $s'$  in terms of  $\phi$ . The constant  $C$  is equal to the value of  $\rho$  at the point corresponding to that for which  $s' = 0$ .

115. For example, in the cycloid  $s = 4a \sin \phi$ ; thus

$$s' = C - 4a \cos \phi.$$

Put  $\phi = \psi + \frac{\pi}{2}$  and  $s' = \sigma + C$ ; thus

$$\sigma = 4a \sin \psi.$$

This shews that the evolute is an equal cycloid.

116. Similarly if the intrinsic equation to a curve be known, that to the involute may be found. For by Art. 114

$$\frac{ds}{d\phi} = C \pm s';$$

therefore

$$s = \int (C \pm s') d\phi.$$

Thus if  $s'$  be known in terms of  $\phi$ , we can find  $s$  in terms of  $\phi$ .

117. For example, in the circle  $s' = a\phi$ . Thus

$$s = \int (C \pm a\phi) d\phi = C\phi \pm \frac{a\phi^2}{2} + C'.$$

If we suppose  $s$  to begin where  $\phi = 0$  we have  $C' = 0$ , and further, if  $s$  begins where the involute meets the circle  $C = 0$ ; thus  $s = \frac{a\phi^2}{2}$ . (See *Dif. Cal.* Art. 333.)

118. It is obvious that by the methods of Arts. 114 and 116 we may find the evolute of the evolute of a curve, or the involute of the involute of a curve, and so on.

119. The student may exercise himself in tracing curves from their intrinsic equations; he will find it useful to take such a curve as the cycloid, the form of which is well known and ascertain that the intrinsic equation does lead to that form; he may then take some of the epicycloids or hypocycloids given in Art. 112. For further information on this subject, and for illustrative figures, the student is referred to two memoirs by Dr Whewell, published in the *Cambridge Philosophical Transactions*, Vol. VIII. page 659, and Vol. IX. page 150.

### *Curves of double Curvature.*

120. Let  $x, y, z$  be the co-ordinates of a point on a curve in space;  $x + \Delta x, y + \Delta y, z + \Delta z$  the co-ordinates of an adjacent point on the curve. Then it is known by the principles of solid geometry, that the length of the chord joining

these two points is  $\sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2}$ . Let  $s$  be the length of the arc of the curve measured from some fixed point up to  $(x, y, z)$ ; and let  $s + \Delta s$  be the length of the arc measured from the same fixed point up to  $(x + \Delta x, y + \Delta y, z + \Delta z)$ . We shall assume that  $\Delta s$  bears to the chord joining the adjacent points a ratio which is ultimately equal to unity when the second point moves along the curve up to the first point. Thus the limit of

$$\frac{\Delta s}{\sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2}}, \text{ that is of } \frac{\frac{\Delta s}{\Delta x}}{\sqrt{\left\{1 + \left(\frac{\Delta y}{\Delta x}\right)^2 + \left(\frac{\Delta z}{\Delta x}\right)^2\right\}}}$$

is unity. Hence

$$\frac{ds}{dx} = \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2 + \left(\frac{dz}{dx}\right)^2\right\}},$$

therefore 
$$s = \int \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2 + \left(\frac{dz}{dx}\right)^2\right\}} dx.$$

From the equations to the curve  $\frac{dy}{dx}$  and  $\frac{dz}{dx}$  may be expressed in terms of  $x$ , and then by integration  $s$  is known in terms of  $x$ .

121. With respect to the assumption in the preceding article, the student may refer to *Dif. Cal. Arts.* 307, 308; he may also hereafter consult De Morgan's *Differential and Integral Calculus*, page 444, and Homersham Cox's *Integral Calculus*, page 95.

122. Suppose, for example, that the curve is determined by the equations

$$y^2 = 4ax \dots \dots \dots (1),$$

$$z = \sqrt{(2cx - x^2)} + c \operatorname{vers}^{-1} \frac{x}{c} \dots \dots \dots (2),$$

so that the curve is formed by the intersection of two cylinders, namely a cylinder which has its generating lines parallel to the axis of  $z$ , and which stands upon the parabola in the plane of  $(x, y)$  given by (1), and a cylinder which has its

generating lines parallel to the axis of  $y$ , and which stands on the cycloid in the plane of  $(x, z)$  given by (2). Then

$$\frac{dy}{dx} = \sqrt{\left(\frac{a}{x}\right)}, \quad \frac{dz}{dx} = \sqrt{\left(\frac{2c-x}{x}\right)};$$

hence 
$$\frac{ds}{dx} = \sqrt{\left(1 + \frac{a}{x} + \frac{2c}{x} - 1\right)} = \sqrt{\left(\frac{2c+a}{x}\right)};$$

therefore 
$$s = \sqrt{(2c+a)} \int \frac{dx}{\sqrt{x}} = 2\sqrt{(2c+a)} \sqrt{x}.$$

No constant is required if we measure the arc from the origin of co-ordinates.

123. The formula given in Art. 120 may be changed into

$$s = \int \sqrt{\left\{1 + \left(\frac{dx}{dy}\right)^2 + \left(\frac{dz}{dy}\right)^2\right\}} dy,$$

and 
$$s = \int \sqrt{\left\{1 + \left(\frac{dx}{dz}\right)^2 + \left(\frac{dy}{dz}\right)^2\right\}} dz,$$

and in some cases these forms may be more convenient than that in Art. 120.

124. Sometimes a curve in space is determined by three equations, which express  $x, y, z$  respectively in terms of an auxiliary variable; then by eliminating this variable we may, if necessary, obtain two equations connecting  $x, y$ , and  $z$ , and thus determine the curve in the ordinary way. Suppose then  $x, y, z$  each a known function of  $t$ ; then

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}, \quad \text{and} \quad \frac{dz}{dx} = \frac{\frac{dz}{dt}}{\frac{dx}{dt}};$$

and 
$$\begin{aligned} s &= \int \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2 + \left(\frac{dz}{dx}\right)^2\right\}} dx \\ &= \int \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2 + \left(\frac{dz}{dx}\right)^2\right\}} \frac{dx}{dt} dt \\ &= \int \sqrt{\left\{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2\right\}} dt. \end{aligned}$$



125. *Application to the Helix.*

This curve may be determined by the equations

$$x = a \cos t, \quad y = a \sin t, \quad z = ct;$$

thus 
$$s = \sqrt{(a^2 + c^2)} \int dt = t \sqrt{(a^2 + c^2)} + C.$$

126. When polar co-ordinates are used to determine position of a point in space, we have the following equations connecting the rectangular and polar co-ordinates of point,

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad z = r \cos \theta.$$

And as a curve in space is determined by two equations between  $x$ ,  $y$ , and  $z$ , it may also be determined by two equations between  $r$ ,  $\theta$ , and  $\phi$ . Thus we may conceive  $r$ ,  $\phi$  to be known functions of  $\theta$ , and therefore  $x$ ,  $y$ , and  $z$  become known functions of  $\theta$ .

Hence

$$\frac{dx}{d\theta} = \sin \theta \cos \phi \frac{dr}{d\theta} - r \sin \theta \sin \phi \frac{d\phi}{d\theta} + r \cos \theta \cos \phi,$$

$$\frac{dy}{d\theta} = \sin \theta \sin \phi \frac{dr}{d\theta} + r \sin \theta \cos \phi \frac{d\phi}{d\theta} + r \cos \theta \sin \phi,$$

$$\frac{dz}{d\theta} = \cos \theta \frac{dr}{d\theta} - r \sin \theta.$$

Therefore 
$$\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 + \left(\frac{dz}{d\theta}\right)^2 = \left(\frac{dr}{d\theta}\right)^2 + r^2 \sin^2 \theta \left(\frac{d\phi}{d\theta}\right)^2 + r^2 \cos^2 \theta$$

and 
$$s = \int \sqrt{\left\{r^2 + \left(\frac{dr}{d\theta}\right)^2 + r^2 \sin^2 \theta \left(\frac{d\phi}{d\theta}\right)^2\right\}} d\theta.$$

This may be transformed into

$$s = \int \sqrt{\left\{r^2 \left(\frac{d\theta}{dr}\right)^2 + 1 + r^2 \sin^2 \theta \left(\frac{d\phi}{dr}\right)^2\right\}} dr$$

or into 
$$s = \int \sqrt{\left\{ r^2 \left( \frac{d\theta}{d\phi} \right)^2 + \left( \frac{dr}{d\phi} \right)^2 + r^2 \sin^2 \theta \right\}} d\phi.$$

127. If  $p$  be the perpendicular from the origin on the tangent to a curve in space, then the equation

$$\frac{ds}{dr} = \frac{r}{\sqrt{(r^2 - p^2)}},$$

which was proved for a *plane* curve in Art. 85 will still hold. For each member of the equation expresses the secant of the angle which the tangent makes with the radius vector at the point of contact.

EXAMPLES.

1. For what values of  $m$  and  $n$  are the curves  $a^m y^n = x^{m+n}$  *rectifiable*? (See Art. 14.)

*Result.* If  $\frac{n}{2m}$  or  $\frac{n}{2m} + \frac{1}{2}$  is an integer,

2. Shew that the length of the arc of a Tractory measured from the cusp is determined by  $s = c \log \frac{c}{y}$ .
3. Shew that the Cissoïd is rectifiable.
4. Shew that the whole length of the curve whose equation is  $4(x^2 + y^2) - a^2 = 3a^{\frac{1}{2}}y^{\frac{3}{2}}$  is equal to  $6a$ .

[It may be proved that  $\left( \frac{ds}{dy} \right)^2 = \frac{a^{\frac{1}{2}}}{4y^{\frac{3}{2}}(a^{\frac{1}{2}} - y^{\frac{3}{2}})}$ ].

5. The length of the arc of the curve

$$(x + y)^{\frac{3}{2}} - (x - y)^{\frac{3}{2}} = a^{\frac{3}{2}}$$

between the limits  $(x_1, y_1)$  and  $(x, y)$  is

$$\frac{1}{2\sqrt{2}} \{ (x + y)^{\frac{3}{2}} + (x - y)^{\frac{3}{2}} \}^{\frac{1}{2}} - \frac{1}{2\sqrt{2}} \{ (x_1 + y_1)^{\frac{3}{2}} + (x_1 - y_1)^{\frac{3}{2}} \}^{\frac{1}{2}}.$$

6. If  $s = ae^{\frac{x}{a}}$ , find the relation between  $x$  and  $y$ .

7. Shew that the intrinsic equation to the parabola is

$$\frac{ds}{d\phi} = \frac{2a}{\cos^3 \phi} \text{ or } s = \frac{a}{2} \log \frac{1 + \sin \phi}{1 - \sin \phi} + \frac{a \sin \phi}{1 - \sin^2 \phi}.$$

8. The intrinsic equation to the curve  $y^3 = ax^3$  is

$$s = \frac{8a}{27} (\sec^3 \phi - 1).$$

9. Draw the curve determined by the intrinsic equation

$$\phi = n \sin s.$$

10. The evolute of an epicycloid is an epicycloid, the radius of the fixed circle being  $\frac{a^2}{a+2b}$  and the radius of


generating circle  $\frac{ab}{a+2b}$ . (Arts. 110 and 114.)

11. Determine the length of a spiral drawn on the surface of a cone, such that any generating line of the cone cuts it at points equidistant from one another.

## CHAPTER VII.

## AREAS OF PLANE CURVES AND OF SURFACES.

*Plane Areas. Rectangular Formulæ. Single Integration.*



128. Let  $DPE$  be a curve, of which the equation is  $y = \phi(x)$ , and suppose  $x, y$  to be the co-ordinates of a point  $P$ . Let  $A$  denote the area included between the curve, the axis of  $x$ , the ordinate  $PM$ , and some fixed ordinate  $AD$ , then (*Dif. Cal.* Art. 43)

$$\frac{dA}{dx} = \phi(x);$$

hence

$$A = \int \phi(x) dx.$$

Let  $\psi(x) + C$  be the integral of  $\phi(x)$ ; thus

$$A = \psi(x) + C.$$

Let  $A_1$  denote the area when the variable ordinate is at a distance  $x_1$  from the axis of  $y$ , and let  $A_2$  denote the area when

the variable ordinate is at a distance  $x_2$  from the axis of  $y$ , then

$$A_1 = \psi(x_1) + C, \quad A_2 = \psi(x_2) + C;$$

therefore  $A_2 - A_1 = \psi(x_2) - \psi(x_1) = \int_{x_1}^{x_2} \phi(x) dx$ .

### 129. Application to the Circle.

The equation to the circle referred to its centre as origin is  $y^2 = a^2 - x^2$ ; here  $\phi(x) = \sqrt{a^2 - x^2}$ ; thus

$$A = \int \phi(x) dx = \int \sqrt{a^2 - x^2} dx = \frac{x \sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + C$$

The constant  $C$  vanishes if we suppose the *fixed ordinate* to coincide with the axis of  $y$ . It will be seen by drawing a figure, that the area comprised between the axis of  $x$ , the axis of  $y$ , the circle, and the ordinate at the distance  $x$  from the axis of  $y$ , may be divided into a triangle and a sector, the values of which are given by the first and second terms in the above expression for  $A$ . This remark may serve to assist the student in remembering the important integral

$$\int \sqrt{a^2 - x^2} dx = \frac{x \sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}.$$

### 130. Application to the Ellipse.

Suppose it required to find the whole area of the ellipse. The equation to the ellipse may be written  $y^2 = \frac{b^2}{a^2}(a^2 - x^2)$ . Hence the area of one quadrant of the ellipse

$$= \int_0^a \frac{b}{a} \sqrt{a^2 - x^2} dx = \frac{b}{a} \int_0^a \sqrt{a^2 - x^2} dx = \frac{b}{a} \frac{\pi a^2}{4} = \frac{\pi ab}{4};$$

hence the area of the ellipse is  $\pi ab$ .

### 131. Application to the Parabola.

The equation to the parabola is  $y^2 = 4ax$ ; here then

$$\phi(x) = \sqrt{4ax},$$

and 
$$\int \sqrt{4ax} \, dx = \frac{4\sqrt{a}}{3} x^{\frac{3}{2}} + C;$$

thus with the notation of Art. 128

$$A_2 - A_1 = \int_{x_1}^{x_2} \sqrt{4ax} \, dx = \frac{4\sqrt{a}}{3} (x_2^{\frac{3}{2}} - x_1^{\frac{3}{2}}).$$

If  $x_1 = 0$  we have for the area  $\frac{4\sqrt{a}}{3} x_2^{\frac{3}{2}}$ , that is, two thirds of the product of the abscissa  $x_2$  and the ordinate  $\sqrt{4ax_2}$ .

### 132. Application to the Cycloid.

The integration required by the formula  $\int y \, dx$  becomes sometimes more easy if we express  $x$  and  $y$  in terms of a new variable. Thus, for example, in the cycloid we can put, (*Dif. Cal.* Art. 358)

$$x = a(1 - \cos \theta), \quad y = a(\theta + \sin \theta);$$

therefore 
$$\int y \, dx = a^2 \int (\theta + \sin \theta) \sin \theta \, d\theta$$

$$= a^2 \int \theta \sin \theta \, d\theta + \frac{a^2}{2} \int (1 - \cos 2\theta) \, d\theta;$$

this gives 
$$a^2 \left( -\theta \cos \theta + \sin \theta \right) + \frac{a^2}{2} \left( \theta - \frac{\sin 2\theta}{2} \right).$$

If we take this between the limits 0 and  $\pi$  for  $\theta$ , we obtain the area of half a cycloid; the result is  $\frac{3a^2\pi}{2}$ . Hence the area of the whole cycloid is equal to three times that of the generating circle.

### 133. The equations to the companion to the cycloid are

$$x = a(1 - \cos \theta), \quad y = a\theta;$$

hence it may be shewn that the area of the whole curve is twice that of the generating circle.

134. If a curve be determined by the equation  $x = \phi(y)$ , then the area contained between the curve, the axis of  $y$ , and

lines drawn parallel to the axis of  $x$  at distances respectively equal to  $y_1$  and  $y_2$  is  $\int_{y_1}^{y_2} \phi(y) dy$ . This is obvious after the proof of the similar proposition in Art. 128.

135. The formulæ in Arts. 128 and 134 furnish one of the most simple and important examples of the application of the Integral Calculus. As we have already remarked, the problem of determining the areas of curves was one of those which gave rise to the Integral Calculus, and the symbols used are very expressive of the process necessary for solving the problem. In the figure to Art. 128, the student will see that the rectangle  $PpMN$  may be appropriately denoted by  $y\Delta x$ , and the process of finding the area of  $ADEB$  amounts to this: we first effect the addition denoted by  $\Sigma y\Delta x$ , and then diminish  $\Delta x$  indefinitely.

136. Suppose we require the area contained between the curve  $y = c \sin \frac{x}{a}$ , the axis of  $x$ , and ordinates at the distances  $x_1$  and  $x_2$  respectively from the axis of  $y$ . We have

$$c \int_{x_1}^{x_2} \sin \frac{x}{a} dx = ca \left( \cos \frac{x_1}{a} - \cos \frac{x_2}{a} \right).$$

Suppose then  $x_1 = 0$  and  $x_2 = a\pi$ ; the area is  $2ca$ . Next suppose  $x_1 = 0$  and  $x_2 = 2a\pi$ ; the result

$$ca \left( \cos \frac{x_1}{a} - \cos \frac{x_2}{a} \right)$$

becomes zero in this case, which is obviously inadmissible, since the area must be some positive quantity. In fact  $\sin \frac{x}{a}$  is *negative* from  $x = a\pi$  to  $x = 2a\pi$ , but in the proof that the area is equal to  $\int y dx$ , it is supposed that  $y$  is *positive*. If  $y$  be really negative the area will be  $\int (-y) dx$ .

Thus in the present example the area will not be

$$\int_0^{2a\pi} \sin \frac{x}{a} dx \quad \text{but} \quad c \int_0^{a\pi} \sin \frac{x}{a} dx + c \int_{a\pi}^{2a\pi} \left( -\sin \frac{x}{a} \right) dx,$$




that is, 
$$c \int_0^{a\pi} \sin \frac{x}{a} dx - c \int_{a\pi}^{2a\pi} \sin \frac{x}{a} dx ;$$

this will give  $2ca + 2ca$ , that is  $4ca$ .

*Plane Areas. Rectangular Formulæ. Double Integration.*

137. In Art. 128 we have obtained a formula for finding the area of a curve; that formula supposes the area to be the limit of a number of elemental areas, each element being a quantity of which  $y\Delta x$  is the type. We shall now proceed to explain another mode of decomposing the required area into elemental areas.



Suppose we require the area included between the curves  $BPQE$  and  $bpqe$ , and the straight lines  $Bb$  and  $Ee$ . Let a series of lines be drawn parallel to the axis of  $y$ , and another series parallel to the axis of  $x$ . Let  $st$  represent one of the rectangles thus formed, and suppose  $x$  and  $y$  to be the co-ordinates of  $s$ , and  $x + \Delta x$  and  $y + \Delta y$  the co-ordinates of  $t$ ; then the area of the rectangle  $st$  is  $\Delta x \Delta y$ . Hence the required area may be found by summing up all the values of  $\Delta x \Delta y$ , and then proceeding to the limit obtained by supposing  $\Delta x$  and  $\Delta y$  to diminish indefinitely.

We effect the required summation of such terms as  $\Delta x \Delta y$  in the following way: we first collect all the rectangles



similar to  $st$  which are contained in the strip  $PQqp$ , and we thus obtain the area of this strip; then we sum up all the strips similar to this strip which lie between  $Bb$  and  $Ee$ . The error we may make by neglecting the element of area which lies at the top and bottom of each strip, and which is not a complete rectangle, will disappear in the limit when  $\Delta x$  and  $\Delta y$  are indefinitely diminished.

Let  $y = \phi(x)$  be the equation to the upper curve, and  $y = \psi(x)$  the equation to the lower curve; let  $OC = c$  and  $OH = h$ , then if  $A$  denote the required area, we have

$$A = \int_c^h \int_{\psi(x)}^{\phi(x)} dx dy;$$

for the symbolical expression here given denotes the process which we have just stated in words.

Now  $\int dy = y$ , therefore  $\int_{\psi(x)}^{\phi(x)} dy = \phi(x) - \psi(x)$ ; thus we have

$$A = \int_c^h \{ \phi(x) - \psi(x) \} dx.$$

In this form we can at once see the truth of the expression, for  $\phi(x) - \psi(x) = PL - pL = Pp$ ; thus  $\{ \phi(x) - \psi(x) \} \Delta x$  may be taken for the area of the strip  $PQqp$ , and the formula asserts that  $A$  is equal to the limit of the sum of such strips.

The lines in the figure are not necessarily equidistant: that is the elements of which  $\Delta x \Delta y$  is the type are not necessarily all of the same area.

138. The result of the preceding article is, that the area  $A$  is found from the equation

$$A = \int_c^h \{ \phi(x) - \psi(x) \} dx.$$

This result may be obtained in a very simple manner as shewn in the latter part of the preceding article, so that it was not absolutely *necessary* to introduce the formula of double integration. We have however drawn attention to the formula

$$A = \int_c^h \int_{\psi(x)}^{\phi(x)} dx dy$$



because of the illustration which is here given of the process of double integration; the student may thus find it easier to apply the processes of double integration to those cases where it is absolutely necessary, of which examples will occur hereafter.

139. If the area which is to be evaluated is bounded by the curves  $x = \psi(y)$ , and  $x = \phi(y)$ , and straight lines parallel to the axis of  $x$  at distances respectively equal to  $c$  and  $h$ , we have in a similar manner

$$A = \int_c^h \int_{\psi(y)}^{\phi(y)} dy dx = \int_c^h \{\phi(y) - \psi(y)\} dy.$$

Some examples of the formulæ of Arts. 137 and 139 will now be considered; we shall see that either of these formulæ may be used in an example, though generally one will be more simple than the other.

140. Required the area included between the parabola  $y^2 = ax$  and the circle  $y^2 = 2ax - x^2$ .

The curves pass through the origin and meet at the point for which  $x = a$ ; thus if we take only that area which lies on the positive side of the axis of  $x$ , we have

$$A = \int_0^a \{\sqrt{(2ax - x^2)} - \sqrt{(ax)}\} dx = \frac{\pi a^2}{4} - \frac{2a^2}{3}.$$

The whole area will therefore be  $2 \left( \frac{\pi a^2}{4} - \frac{2a^2}{3} \right)$ .


Suppose that we wish in this example to integrate with respect to  $x$  first. From the equation  $y^2 = 2ax - x^2$  we deduce  $x = a \pm \sqrt{(a^2 - y^2)}$ , and it will appear at once from a figure that we must take the lower sign in the present question.

Thus let  $x_1$  stand for  $a - \sqrt{(a^2 - y^2)}$ , and  $x_2$  for  $\frac{y^2}{a}$ , then

$$\begin{aligned} A &= \int_0^a \int_{x_1}^{x_2} dy dx = \int_0^a \left\{ \frac{y^2}{a} - a + \sqrt{(a^2 - y^2)} \right\} dy \\ &= \frac{a^2}{3} - a^2 + \frac{\pi a^2}{4} = \frac{\pi a^2}{4} - \frac{2a^2}{3}, \text{ as before.} \end{aligned}$$

The reader should draw the figure and pay close attention to the *limits* of the integrations.

141. In the accompanying figure  $S$  is the centre of a circle  $BLD$ ,  $S$  is also the focus of a parabola  $ALC$ ; we shall



indicate the integrations that should be performed in order to obtain the areas  $ALB$  and  $LDC$ . This example is introduced for the purpose of illustrating the processes of double integration, and not for any interest in the results: the areas can be easily ascertained by means of formulæ already given; thus  $ALB$  is the difference of the parabolic area  $ALS$  and the quadrant  $SLB$ ; and similarly  $LDC$  is known.

In finding the area  $ALB$  it will be convenient to suppose the positive direction of the axis of  $x$  to be that towards the left hand; thus if  $4a$  be the latus rectum of the parabola, and therefore  $2a$  the radius of the circle, the equation to the parabola is  $y^2 = 4a(a - x)$ , and that to the circle  $y^2 = 4a^2 - x^2$ .

Suppose we integrate with respect to  $x$  first, then

$$\text{area } ALB = \int_0^{2a} \int_{x_1}^{x_2} dy dx,$$

where  $x_1 = a - \frac{y^2}{4a}, \quad x_2 = \sqrt{4a^2 - y^2}.$

For here  $(x_2 - x_1) \Delta y$  represents a strip included between the two curves and two lines parallel to the axis of  $x$ ; and

strips are situated at distances from the axis of  $x$  ranging between 0 and  $2a$ , so that the integration with respect to  $y$  is taken between the limits 0 and  $2a$ .

Suppose we integrate with respect to  $y$  first; we shall then have to divide the area into two parts by the line  $AF$ . Let

$$y_1 = \sqrt{(4a^2 - 4ax)}, \quad y_2 = \sqrt{(4a^2 - x^2)};$$

$$\text{then area } ALF = \int_0^a \int_{y_1}^{y_2} dx dy = \int_0^a (y_2 - y_1) dx;$$

$$\text{area } AFB = \int_a^{2a} \int_0^{y_2} dx dy = \int_a^{2a} y_2 dx;$$

the sum of these two parts expresses the area  $ALB$ .

Next take the area  $LDC$ ; suppose now the positive direction of the axis of  $x$  to be that towards the right hand, then the equation to the parabola is  $y^2 = 4a(a+x)$ , and that to the circle  $y^2 = 4a^2 - x^2$ .

Suppose we integrate with respect to  $y$  first; let

$$y_1 = \sqrt{(4a^2 - x^2)} \quad \text{and} \quad y_2 = \sqrt{(4a^2 + 4ax)};$$

$$\text{then area } DLC = \int_0^{2a} \int_{y_1}^{y_2} dx dy.$$

Suppose we integrate with respect to  $x$  first; we shall then have to divide the area into two parts by the line  $LK$ . Let

$$x_1 = \sqrt{(4a^2 - y^2)}, \quad x_2 = \frac{y^2}{4a} - a;$$

then we shall find that  $DC = 2a\sqrt{3} = b$  suppose; thus

$$\text{area } DLK = \int_0^{2a} \int_{x_1}^{2a} dy dx,$$

$$\text{area } CLK = \int_{2a}^b \int_{x_2}^{2a} dy dx;$$

the sum of these two parts expresses the area  $LDC$ .

142. One case in which the formulæ of Art. 137 are useful is that in which the bounding curves are different branches of the same curve. Suppose the equation to a curve to be  $(y - mx - c)^2 = a^2 - x^2$ ; thus

$$y = mx + c \pm \sqrt{a^2 - x^2}.$$

Here we may put

$$\psi(x) = mx + c - \sqrt{a^2 - x^2},$$

$$\phi(x) = mx + c + \sqrt{a^2 - x^2};$$

thus  $\phi(x) - \psi(x) = 2\sqrt{a^2 - x^2}$ , and the complete area of the curve is

$$\int_{-a}^a 2\sqrt{a^2 - x^2} dx, \text{ that is } \pi a^2.$$

143. We have hitherto supposed the axes rectangular, but if they are oblique and inclined at an angle  $\omega$ , the formula in Art. 128 becomes

$$A = \sin \omega \int \phi(x) dx,$$


and a similar change is made in all the other formulæ. It is obvious that such elements of area as are denoted by  $y\Delta x$  and  $\Delta y\Delta x$  when the axes are rectangular will be denoted by  $\sin \omega y\Delta x$  and  $\sin \omega \Delta y\Delta x$  when the axes are inclined at an angle  $\omega$ .

For example, the equation to the parabola is  $y^2 = 4a'x$  when the axes are the oblique system formed by a diameter and the tangent at its extremity; hence the area included between the curve, the axis of  $x$ , and an ordinate at the point for which  $x = c$ , is

$$\sin \omega \int_0^c \sqrt{4a'x} dx = \frac{4 \sin \omega \sqrt{a'c^{\frac{3}}{3}}}{3},$$

that is two-thirds of the parallelogram which has the abscissa  $c$  and the ordinate at its extremity for adjacent sides.

*Plane Areas. Polar Formulae. Single Integration.*



144. Let  $APQ$  be a curve, of which the polar equation is  $r = \phi(\theta)$ , and suppose  $r, \theta$  to be the co-ordinates of a point  $P$ . Let  $A$  denote the area included between the curve, the radius vector  $SA$  drawn to a fixed point  $A$ , and the radius vector  $SP$ , then (*Dif. Cal. Art. 313*)

$$\frac{dA}{d\theta} = \frac{\{\phi(\theta)\}^2}{2}.$$

Hence 
$$A = \frac{1}{2} \int \{\phi(\theta)\}^2 d\theta.$$

Let  $\psi(\theta)$  be the integral of  $\frac{\{\phi(\theta)\}^2}{2}$ , then

$$A = \psi(\theta) + C.$$

Let  $A_1$  denote the area when the variable radius vector is at an angular distance  $\theta_1$  from the initial line, and let  $A_2$  denote the area when the variable radius vector is at an angular distance  $\theta_2$  from the initial line; then

$$A_1 = \psi(\theta_1) + C, \quad A_2 = \psi(\theta_2) + C,$$

therefore 
$$A_2 - A_1 = \psi(\theta_2) - \psi(\theta_1) = \frac{1}{2} \int_{\theta_1}^{\theta_2} \{\phi(\theta)\}^2 d\theta.$$

145. *Application to the Equiangular Spiral.*

In this curve  $r = be^{\frac{\theta}{c}}$ ; thus

$$A = \frac{1}{2} \int b^2 e^{\frac{2\theta}{c}} d\theta = \frac{b^2 c}{4} e^{\frac{2\theta}{c}} + C,$$

$$\text{and } A_2 - A_1 = \frac{1}{2} \int_{\theta_1}^{\theta_2} b^2 e^{2\theta} d\theta = \frac{b^2 c}{4} (e^{\frac{2\theta_2}{c}} - e^{\frac{2\theta_1}{c}}) = \frac{c}{4} (r_2^2 - r_1^2),$$

where  $r_1$  and  $r_2$  are the extreme radii vectores of the area considered.

#### 146. *Application to the Parabola.*

Let the focus be the pole, then

$$r = \frac{a}{\cos^2 \frac{\theta}{2}}; \text{ thus } A = \frac{a^2}{2} \int \frac{d\theta}{\cos^4 \frac{\theta}{2}}$$

$$= \frac{a^2}{2} \int \left(1 + \tan^2 \frac{\theta}{2}\right) \sec^2 \frac{\theta}{2} d\theta = a^2 \tan \frac{\theta}{2} + \frac{a^2}{3} \tan^3 \frac{\theta}{2} + C.$$

$$\text{Hence } A_2 - A_1 = a^2 \left( \tan \frac{\theta_2}{2} - \tan \frac{\theta_1}{2} \right) + \frac{a^2}{3} \left( \tan^3 \frac{\theta_2}{2} - \tan^3 \frac{\theta_1}{2} \right).$$

Suppose that  $\theta_1 = 0$  and  $\theta_2 = \frac{\pi}{2}$ , then we obtain for the area  $a^2 + \frac{a^2}{3}$ , that is  $\frac{4a^2}{3}$ ; this agrees with Art. 131.

For another example we will suppose the Parabola referred to the intersection of the directrix and the axis as pole, the axis being the initial line. Here

$$r = 2a \frac{\cos \theta - \sqrt{(\cos 2\theta)}}{\sin^2 \theta},$$

$$\begin{aligned} \text{thus } A &= 2a^2 \int \frac{\cos^2 \theta + \cos 2\theta - 2 \cos \theta \sqrt{(\cos 2\theta)}}{\sin^4 \theta} d\theta \\ &= 2a^2 \int \frac{2 \cos^2 \theta - \sin^2 \theta}{\sin^4 \theta} d\theta - 4a^2 \int \frac{\cos \theta \sqrt{(\cos 2\theta)}}{\sin^4 \theta} d\theta. \end{aligned}$$

Now 
$$\int \frac{2 \cos^2 \theta - \sin^2 \theta}{\sin^4 \theta} d\theta = \int (2 \cot^2 \theta - 1) \operatorname{cosec}^2 \theta d\theta$$

$$= -\frac{2}{3} \cot^3 \theta + \cot \theta.$$

And 
$$\int \frac{\cos \theta \sqrt{(\cos 2\theta)} d\theta}{\sin^4 \theta} = \int \frac{\sqrt{(1 - 2 \sin^2 \theta)} d \sin \theta}{\sin^4 \theta};$$

assume  $\sin \theta = \frac{1}{t}$ , then the integral becomes

$$-\int \sqrt{(t^2 - 2)} t dt, \text{ that is, } -\frac{1}{8} (t^2 - 2)^{\frac{3}{2}}.$$

Hence, adding the constant, we have

$$A = \frac{4a^2}{3} (\operatorname{cosec}^2 \theta - 2)^{\frac{3}{2}} - \frac{4a^2}{3} \cot^3 \theta + 2a^2 \cot \theta + C$$

$$= 2a^2 \cot \theta + \frac{4a^2}{3} \frac{(\cos 2\theta)^{\frac{3}{2}} - \cos^3 \theta}{\sin^3 \theta} + C.$$

The constant will be zero if  $A$  commences from the initial line.

147. *Application to the curve*  $r = a(\theta + \sin \theta)$ . Here

$$A = \frac{a^2}{2} \int (\theta + \sin \theta)^2 d\theta = \frac{a^2}{2} \int (\theta^2 + 2\theta \sin \theta + \sin^2 \theta) d\theta;$$

and 
$$\int \theta \sin \theta d\theta = -\theta \cos \theta + \sin \theta$$

$$\int \sin^2 \theta d\theta = \frac{1}{2} \int (1 - \cos 2\theta) d\theta = \frac{\theta}{2} - \frac{\sin 2\theta}{4},$$

thus 
$$A = \frac{a^2}{2} \left\{ \frac{\theta^3}{3} - 2\theta \cos \theta + 2 \sin \theta + \frac{\theta}{2} - \frac{\sin 2\theta}{4} \right\} + C.$$

Suppose we require the area of the smallest portion which is bounded by the curve and by a radius vector which is




inclined to the initial line at a right angle; then we have 0 and  $\frac{1}{2}\pi$  as the limits of the integration. Thus the required area is

$$\frac{a^2}{2} \left\{ \frac{\pi^3}{24} + \frac{\pi}{4} + 2 \right\}.$$

*Plane Curves. Polar Formulæ. Double Integration.*

148. In Art. 144 we have obtained a formula for finding the area of a curve; that formula supposes the area to be the limit of a number of elemental areas, each element being a quantity of which  $\frac{1}{2}r^2 \Delta\theta$  is the type. We shall now proceed to explain another mode of decomposing the required area into elemental areas.



Suppose we require the area included between the curves  $BPQE$  and  $bpqe$ , and the straight lines  $Bb$  and  $Ee$ . Let a series of radii vectores be drawn from  $O$ , and a series of circles with  $O$  as centre; thus the plane area is divided into a series of curvilinear quadrilaterals. Let  $st$  represent one of these elements, and suppose  $r$  and  $\theta$  to be the polar co-ordinates of  $s$ , and  $r + \Delta r$  and  $\theta + \Delta\theta$  the polar co-ordinates of  $t$ ; then the area of the element  $st$  will be ultimately  $r\Delta\theta \Delta r$ . Hence the required area is to be found by summing up all the values of  $r\Delta\theta \Delta r$ , and then proceeding to the limit obtained by supposing  $\Delta\theta$  and  $\Delta r$  to diminish indefinitely.

We effect the required summation of such terms as  $r\Delta\theta\Delta r$  in the following way: we first collect all the elements similar to  $st$  which are contained in the strip  $PQqp$ , and thus obtain the area of the strip; then we sum up all the strips similar to this strip which lie between  $Bb$  and  $Ee$ .

Let  $r = \phi(\theta)$  be the equation to the curve  $BPQE$  and  $r = \psi(\theta)$  the equation to the curve  $bpqe$ , let  $\alpha$  and  $\beta$  be the angles which  $OB$  and  $OE$  make respectively with  $Ox$ ; and let  $A$  denote the required area, then

$$A = \int_{\alpha}^{\beta} \int_{\psi(\theta)}^{\phi(\theta)} r d\theta dr;$$

for the symbolical expression here given denotes the process which we have just stated in words.

Now  $\int r dr = \frac{r^2}{2}$ , therefore

$$\int_{\psi(\theta)}^{\phi(\theta)} r dr = \frac{1}{2} [\{\phi(\theta)\}^2 - \{\psi(\theta)\}^2],$$

thus we have

$$A = \frac{1}{2} \int_{\alpha}^{\beta} [\{\phi(\theta)\}^2 - \{\psi(\theta)\}^2] d\theta.$$

In this form we can see at once the truth of the expression, for  $OP = \phi(\theta)$  and  $Oq = \psi(\theta)$ , and thus

$$\frac{1}{2} \{\phi(\theta)\}^2 \Delta\theta - \frac{1}{2} \{\psi(\theta)\}^2 \Delta\theta$$

may be taken for the area of the strip  $PQqp$ , and the formula asserts that the area  $A$  is equal to the limit of the sum of such strips.

149. The remark made in Art. 138 may be repeated here; we have introduced the process in the former part of the preceding article, not because double integration is absolutely necessary for finding the area of a curve, but because the process of finding the area of a curve illustrates double integration.


150. If the area which is to be evaluated is bounded by the curves whose equations are  $\theta = \phi(r)$ ,  $\theta = \psi(r)$  respectively,

and by the circles whose equations are  $r = a$  and  $r = b$  respectively; it will be convenient to integrate with respect to  $\theta$  first. In this case, instead of first summing up all the elements like  $st$ , which form the strip  $PQqp$ , we first sum up all the elements similar to  $st$  which are included between the two circles which bound  $st$  and the curves determined by  $\theta = \phi(r)$  and  $\theta = \psi(r)$ . Thus we have

$$A = \int_a^b \int_{\psi(r)}^{\phi(r)} r dr d\theta.$$

Some examples of the formulæ in Arts. 148 and 150 will now be considered; we shall see that either of these formulæ may be used in an example, although one may be more convenient than the other.

151. We will apply the formulæ to find the area between the two semicircles  $OPB$  and  $Opb$  and the straight line  $bB$ .



Let  $Ob = c$ ,  $OB = h$ , then the equation to  $OPB$  is  $r = h \cos \theta$ , and the equation to  $Opb$  is  $r = c \cos \theta$ . Thus the area

$$= \int_0^{\frac{\pi}{2}} \int_{c \cos \theta}^{h \cos \theta} r dr d\theta.$$

Now 
$$\int_{c \cos \theta}^{h \cos \theta} r dr = \frac{1}{2} (h^2 - c^2) \cos^2 \theta;$$

thus the area 
$$= \frac{1}{2} (h^2 - c^2) \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta = \frac{\pi}{8} (h^2 - c^2).$$

Suppose we wish to integrate with respect to  $\theta$  first; we shall then have to divide the area into two parts by describing an arc of a circle from  $O$  as centre, with radius  $Ob$ . Then

the area bounded by this arc, the straight line  $Bb$ , and the larger semicircle is

$$\int_c^h \int_0^{\cos^{-1} \frac{r}{h}} r dr d\theta.$$

The area bounded by the aforesaid arc, the semicircle  $Opb$ , and the larger semicircle is

$$\int_0^e \int_{\cos^{-1} \frac{r}{c}}^{\cos^{-1} \frac{r}{h}} r dr d\theta.$$

The sum of these two parts expresses the required area.

152. Let us apply polar formulæ to the example in Art. 141. With  $S$  as pole, the polar equation to the parabola is  $r(1 + \cos \theta) = 2a$  or  $r \cos^2 \frac{\theta}{2} = a$  where  $\theta$  is measured from  $SB$ ; and the polar equation to the circle is  $r = 2a$ . Hence, if we integrate with respect to  $r$  first,

$$\text{area } ALB = \int_0^{\frac{\pi}{2}} \int_{a \sec^2 \frac{\theta}{2}}^{2a} r d\theta dr.$$

If we integrate with respect to  $\theta$  first, we shall have if  $\theta_1 = \cos^{-1} \frac{2a-r}{r}$

$$\text{area } ALB = \int_a^{2a} \int_0^{\theta_1} r dr d\theta.$$

Next consider the area  $DLC$ . The equation to  $DC$  is  $r \cos \theta = -2a$ ; the length of  $SC$  is  $4a$ , and the angle  $BSC$  is  $\frac{2\pi}{3}$ . Let  $\theta_1 = \cos^{-1} \frac{2a-r}{r}$ ,  $\theta_2 = \cos^{-1} \left( \frac{-2a}{r} \right)$ . Then if we integrate with respect to  $\theta$  first,

$$\text{area } DLC = \int_{2a}^{4a} \int_{\theta_1}^{\theta_2} r dr d\theta.$$

If we integrate with respect to  $r$  first, we shall have to divide the area into two parts, by the line joining  $S$  with  $C$ .

The area of the portion which has  $LC$  for one of its boundaries is


$$\int_{\frac{\pi}{2}}^{\frac{2\pi}{3}} \int_{2a}^{a \sec^2 \frac{\theta}{2}} r d\theta dr.$$

The area of the remaining portion is

$$\int_{\frac{2\pi}{3}}^{\pi} \int_{2a}^{2a \sec \theta} r d\theta dr.$$

The sum of these two parts expresses the required area.

153. A good example is supplied by the problem of finding the area included between two radii vectores and two different branches of the same polar curve.



Suppose  $BPpb$ ,  $CQqc$  to be two different arcs of a spiral, and that the area is to be evaluated which is bounded by these arcs and the straight lines  $BC$  and  $bc$ ; then the area is

$$\frac{1}{2} \int (r_2^2 - r_1^2) d\theta,$$

where  $r$ , denotes any radius vector of the exterior arc, as  $SQ$ , and  $r$ , the corresponding radius vector  $SP$  of the interior arc. The limits of  $\theta$  will be given by the angles which  $SB$  and  $Sb$  respectively make with the initial line.

Take for example the spiral of Archimedes; let  $\theta$  be the whole angle which the radius vector has revolved through from the initial line until it takes the position  $SP$ ; so that  $\theta$  may be an angle of any magnitude. From the nature of the curve we have  $SP$  or  $r = a\theta$ , where  $a$  is some constant. If then  $CQ$  is the next branch to  $BP$  we shall have

$$SQ = a(\theta + 2\pi).$$

Suppose  $\theta_1$  and  $\theta_2$  the values of  $\theta$  for  $SB$  and  $Sb$  respectively; thus the area  $BbcC$ .

$$\begin{aligned} &= \frac{a^2}{2} \int_{\theta_1}^{\theta_2} \{(\theta + 2\pi)^2 - \theta^2\} d\theta \\ &= \frac{a^2}{2} \{2\pi(\theta_2^2 - \theta_1^2) + 4\pi^2(\theta_2 - \theta_1)\}. \end{aligned}$$

154. The student will remark a certain difference between the formulæ  $\iint dx dy$  and  $\iint r d\theta dr$ , which express the area of a plane figure. The former supposes the area decomposed into a number of rectangles and  $\Delta x \Delta y$  represents the true area of one rectangle. Hence in taking the aggregate of these rectangles to represent the required area the only error that can arise is owing to the neglect of the irregular elements which occur at the top and bottom of each strip; as we have already remarked in Art. 137. But in the second case  $r \Delta\theta \Delta r$  is not the *accurate value* of the area of one of the elements, so that an error is made in the case of every element. It is therefore important to shew formally that the error disappears in the limit, which may be done as follows. The element  $st$  in the figure of Art. 148 is the difference of two circular sectors, and its exact area is

$$\frac{1}{2}(r + \Delta r)^2 \Delta\theta - \frac{1}{2}r^2 \Delta\theta,$$

that is,

$$r \Delta r \Delta\theta + \frac{1}{2}(\Delta r)^2 \Delta\theta.$$

In taking the former term to represent the area we neglect  $\frac{1}{2}(\Delta r)^2 \Delta \theta$ . Hence the ratio of the term neglected to the term retained

$$= \frac{\frac{1}{2}(\Delta r)^2 \Delta \theta}{r \Delta r \Delta \theta} = \frac{\Delta r}{2r}.$$

By taking  $\Delta r$  small enough this ratio may be made as small as we please. Hence we may infer that the sum of the neglected terms will ultimately vanish in comparison with the sum of the terms retained, that is all error disappears in the limit.

*Other Polar Formulæ.*

155. Let  $s$  be the length of the arc of a curve measured from some fixed point up to the point whose co-ordinates are  $r$  and  $\theta$ ; let  $p$  be the perpendicular from the origin on the tangent at the latter point; then the sine of the angle between this tangent and the corresponding radius vector is  $r \frac{d\theta}{ds}$ , (*Dif.*

*Cal.* Art. 310); also  $\frac{p}{r}$  is another expression for this sine;

hence,  $r \frac{d\theta}{ds} = \frac{p}{r}$ . Let  $A$  denote the area between the curve and certain limiting radii vectors; then

$$A = \frac{1}{2} \int r^2 d\theta = \frac{1}{2} \int r^2 \frac{d\theta}{ds} ds = \frac{1}{2} \int r \frac{p}{r} ds = \frac{1}{2} \int p ds;$$

the limits of  $s$  in the latter integral must be such as correspond to the limiting radii vectores of the area considered.

The result can be illustrated geometrically; suppose  $P, Q$  adjacent points on a curve,  $S$  the pole,  $p'$  the perpendicular from  $S$  on the chord  $PQ$ ; then, the area of the triangle  $PQS$

$$= \frac{1}{2} p' \times \text{chord } PQ.$$

Now suppose  $Q$  to approach indefinitely near to  $P$ , then  $p' = p$ , and the limit of the ratio of the chord  $PQ$  to the arc  $PQ$  is unity.

156. Since  $\int p ds = \int p \frac{ds}{dr} dr = \int \frac{pr dr}{\sqrt{(r^2 - p^2)}} \text{ (Art. 85),}$

we have  $A = \frac{1}{2} \int \frac{pr dr}{\sqrt{(r^2 - p^2)}} .$

157. *Application to the Epicycloid.*

Here  $p^2 = \frac{c^2 (r^2 - a^2)}{c^2 - a^2}$ ; thus

$$\begin{aligned} A &= \frac{1}{2} \int \frac{c \sqrt{(r^2 - a^2)} r dr}{a \sqrt{(c^2 - r^2)}} = \frac{c}{2a} \int \frac{\sqrt{(r^2 - a^2)} r dr}{\sqrt{\{c^2 - a^2 - (r^2 - a^2)\}}} \\ &= \frac{c}{2a} \int \frac{z^2 dz}{\sqrt{(c^2 - a^2 - z^2)}} \text{ where } z^2 = r^2 - a^2. \end{aligned}$$

Now

$$\begin{aligned} \int \frac{z^2 dz}{\sqrt{(c^2 - a^2 - z^2)}} &= \int \frac{z^2 - (c^2 - a^2)}{\sqrt{(c^2 - a^2 - z^2)}} dz + (c^2 - a^2) \int \frac{dz}{\sqrt{(c^2 - a^2 - z^2)}} \\ &= (c^2 - a^2) \int \frac{dz}{\sqrt{(c^2 - a^2 - z^2)}} - \int \sqrt{(c^2 - a^2 - z^2)} dz \\ &= \frac{c^2 - a^2}{2} \sin^{-1} \frac{z}{\sqrt{(c^2 - a^2)}} - \frac{z \sqrt{(c^2 - a^2 - z^2)}}{2} \\ &= \frac{c^2 - a^2}{2} \sin^{-1} \frac{\sqrt{(r^2 - a^2)}}{\sqrt{(c^2 - a^2)}} - \frac{\sqrt{(r^2 - a^2)} \sqrt{(c^2 - r^2)}}{2} . \end{aligned}$$

Taking this between the limits  $r = a$  and  $r = c$ , we get  $\frac{c^2 - a^2}{2} \frac{\pi}{2}$ , that is,  $b(a + b)\pi$ . Hence the area is  $\frac{c}{2a} b(a + b)\pi$ ,

that is,  $\frac{(a + 2b)b(a + b)\pi}{2a}$ . By doubling this result we obtain the area between the curve and the radii vectors drawn to two consecutive cusps, which is therefore  $\frac{(a + 2b)b(a + b)\pi}{a}$ .

The area of the circular sector which forms part of this area is  $\pi ab$ ; subtract the latter and we obtain the area between an arc of the epicycloid extending from one cusp to the next cusp



and the fixed circle on which the generating circle rolls; the result is

$$\frac{\pi b^3}{a} (3a + 2b).$$

158. Similarly in the hypocycloid the area between the fixed circle and the part of the curve which extends between two consecutive cusps may be found. If  $a$  is greater than  $b$  the result is

$$\frac{\pi b^3}{a} (3a - 2b).$$

*Area between a Curve and its Evolute.*

159. In the figures to Art. 114, if we suppose the string or line  $PQ$  to move through a small angle  $\Delta\phi$ , the figure between the two positions of the line and the curve  $AP$  may be considered ultimately as a sector of a circle; its area will therefore be  $\frac{1}{2}\rho^2\Delta\phi$ , where  $\rho = PQ$ . Thus if  $A$  denote the whole area bounded by the curve, its evolute, and two radii of curvature corresponding to the values  $\phi_1$  and  $\phi_2$  of  $\phi$ , we have

$$A = \frac{1}{2} \int_{\phi_1}^{\phi_2} \rho^2 d\phi.$$

Since  $\frac{d\phi}{ds} = \frac{1}{\rho}$ , we may also write this

$$A = \frac{1}{2} \int \rho ds,$$

the limits of  $s$  being properly taken so as to correspond with the known limits of  $\phi$ .

160. *Application to the Catenary.*


Here  $s = c \tan \phi$ , Art. 109,

therefore  $\rho = c \sec^2 \phi$ ,  $A = \frac{1}{2} \int_{\phi_1}^{\phi_2} c^2 \sec^4 \phi d\phi$ ;

and 
$$\int \sec^4 \phi \, d\phi = \tan \phi + \frac{1}{3} \tan^3 \phi + C;$$

thus  $A$  is known.

*Area of Surfaces of Revolution.—Rectangular Formulæ.*



161. Let  $A$  be a fixed point in the curve  $APQ$ ; let  $x, y$  be the co-ordinates of any point  $P$ , and  $s$  the length of the arc  $AP$ . Suppose the curve to revolve round the axis of  $x$ , and let  $S$  denote the area of the surface formed by the revolution of  $AP$ ; then (*Dif. Cal.* Art. 315)

$$\frac{dS}{ds} = 2\pi y;$$

therefore 
$$S = \int 2\pi y \, ds \dots\dots\dots(1),$$

thus 
$$S = \int 2\pi y \frac{ds}{dx} \, dx \dots\dots\dots(2),$$

and 
$$S = \int 2\pi y \frac{ds}{dy} \, dy \dots\dots\dots(3).$$

Of these three forms we can choose in any particular example that which is most convenient. If  $y$  can be easily expressed in terms of  $s$  we may use (1); if  $\frac{ds}{dx}$  can be easily expressed in terms of  $y$  we may use (3); in some cases it may be more convenient to express  $y$  and  $\frac{ds}{dx}$  in terms of  $x$  and use (2).

In each case the area of the surface generated by the arc of the curve which lies between assigned points will be found by integrating between appropriate limits.

162. *Application to the Cylinder.*

Suppose a straight line parallel to the axis of  $x$  to revolve round the axis of  $x$ , thus generating a right circular cylinder: let  $a$  be the distance of the revolving line from the axis of  $x$ ;

then 
$$y = a, \text{ and } \frac{ds}{dx} = 1;$$

thus by equation (2) of Art. 161,

$$S = 2\pi \int a \, dx = 2\pi ax + C.$$

Suppose the abscissæ of the extreme points of the portion of the line which revolves to be  $x_1$  and  $x_2$ ; then the surface generated

$$= 2\pi a \int_{x_1}^{x_2} dx = 2\pi a (x_2 - x_1).$$

163. *Application to the Cone.*

Let a straight line which passes through the origin and is inclined to the axis of  $x$  at an angle  $\alpha$  revolve round the axis of  $x$ , and thus generate a conical surface. Then

$$y = x \tan \alpha, \text{ and } \frac{ds}{dx} = \sec \alpha;$$

thus by equation (2) of Art. 161,

$$S = 2\pi \int \tan \alpha \sec \alpha x \, dx = \pi \tan \alpha \sec \alpha x^2 + C.$$

Hence the surface of the frustum of a cone cut off by planes perpendicular to its axis at distances  $x_1$ ,  $x_2$  respectively from the vertex is

$$\pi \tan \alpha \sec \alpha (x_2^2 - x_1^2).$$

Suppose  $x_1 = 0$ , and let  $r$  be the radius of the section made

by the plane at the distance  $x_2$ , then  $r = x_2 \tan \alpha$ , and the area is

$$\pi \operatorname{cosec} \alpha r^2.$$

164. *Application to the Sphere.*

Let the circle given by the equation  $y^2 = a^2 - x^2$  revolve round the axis of  $x$ ; here

$$\frac{dy}{dx} = -\frac{x}{y},$$

and 
$$\frac{ds}{dx} = \sqrt{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}} = \sqrt{\left(1 + \frac{x^2}{y^2}\right)} = \frac{a}{y}.$$

Hence by equation (2) of Art. 161,

$$S = 2\pi \int y \frac{a}{y} dx = 2\pi a \int dx = 2\pi ax + C;$$

Thus the surface included between the planes determined by

$$x = x_1 \text{ and } x = x_2 \text{ is } 2\pi a (x_2 - x_1).$$

Hence the area of a zone of a sphere depends only on the height of the zone and the radius of the sphere, and is equal to the area which the planes that bound it would cut off from a cylinder having its axis perpendicular to the planes and circumscribing the sphere.

165. *Application to the Prolate Spheroid.*

Let the ellipse given by  $a^2y^2 + b^2x^2 = a^2b^2$  revolve round the axis of  $x$  which is supposed to coincide with the major axis of the ellipse; here

$$\frac{dy}{dx} = -\frac{b^2x}{a^2y},$$

and 
$$\frac{ds}{dx} = \sqrt{\left(1 + \frac{b^4x^2}{a^4y^2}\right)} = \frac{b\sqrt{(a^2 - e^2x^2)}}{ay}.$$

Hence by equation (2) of Art. 161,

$$\begin{aligned} S &= \frac{2\pi b}{a} \int \sqrt{(a^2 - e^2x^2)} dx = \frac{2\pi b e}{a} \int \sqrt{\left(\frac{a^2}{e^2} - x^2\right)} dx \\ &= \frac{\pi b e}{a} \left\{ x \sqrt{\left(\frac{a^2}{e^2} - x^2\right)} + \frac{a^2}{e^2} \sin^{-1} \frac{ex}{a} \right\}. \end{aligned}$$

The surface generated by the revolution of a quadrant of the ellipse will be obtained by taking 0 and  $a$  as the limits of  $x$  in the integration. This gives

$$\pi ab \left\{ \sqrt{1 - e^2} + \frac{\sin^{-1} e}{e} \right\}.$$

166. Suppose one curve to have for its equation  $y = \phi(x)$ , and another curve to have for its equation  $y = \psi(x)$ , and let both curves revolve round the axis of  $x$ . Let  $s_1$  and  $s_2$  denote the lengths of arcs measured from fixed points in the two curves up to the point whose abscissa is  $x$ . Let  $S$  denote the sum of the areas of both surfaces intercepted between two planes perpendicular to the axis of  $x$  at the distances  $x_1$  and  $x_2$  respectively from the origin. Then, by Art. 161,

$$S = 2\pi \int_{x_1}^{x_2} \left\{ \phi(x) \frac{ds_1}{dx} + \psi(x) \frac{ds_2}{dx} \right\} dx.$$

Suppose, for example, that there is a curve which is bisected by the line  $y = a$ , so that we may put  $y = a + \chi(x)$  for the upper branch and  $y = a - \chi(x)$  for the lower branch. Hence

$$\frac{ds_1}{dx} = \frac{ds_2}{dx},$$

and 
$$S = 4\pi a \int_{x_1}^{x_2} \frac{ds_1}{dx} dx = 4\pi a \int ds_1,$$

the limits for  $s_1$  being taken so as to correspond with the assigned limits of  $x$ .

Hence, if there be any complete curve which is bisected by a straight line and made to revolve round an axis which is parallel to this line at a distance  $a$  from it and which does not cut the curve, the area of the whole surface generated is equal to the length of the curve multiplied by  $2\pi a$ .

167. For example, take the circle given by the equation

$$(x - h)^2 + (y - k)^2 - c^2 = 0.$$

Here the area of the whole surface generated by the revolution of the circle round the axis of  $x$  will be  $2\pi k \times 2\pi c$ .

There is no difficulty in this example in obtaining separately the two portions of the surface. For the part above the line  $y = k$ , we have  $2\pi \int y ds$ , that is,

$$2\pi \int [k + \sqrt{c^2 - (x - h)^2}] ds,$$

that is,  $2\pi \int k ds + 2\pi \int \sqrt{c^2 - (x - h)^2} ds$ .

The former of these integrals is  $2\pi ks$ ; the latter is equal to

$$2\pi \int \sqrt{c^2 - (x - h)^2} \frac{ds}{dx} dx,$$

which will reduce to  $2\pi \int c dx$ , that is,  $2\pi cx$ . Hence the surface required is found by taking the expression  $2\pi ks + 2\pi cx$  between proper limits.

#### *Area of Surfaces of Revolution. Polar Formulae.*

168. It may be sometimes convenient to use polar coordinates; thus from Art. 161 we deduce

$$S = \int 2\pi y ds = \int 2\pi y \frac{ds}{d\theta} d\theta = \int 2\pi r \sin \theta \frac{ds}{d\theta} d\theta$$

where  $\frac{ds}{d\theta} = \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2}$ .

#### 169. *Application to the Cardioid.*

Here  $r = a(1 + \cos \theta)$ , thus

$$\frac{ds}{d\theta} = a \sqrt{(1 + \cos \theta)^2 + \sin^2 \theta} = a \sqrt{2 + 2 \cos \theta} = 2a \cos \frac{\theta}{2};$$


therefore

$$\begin{aligned} S &= 4\pi a^2 \int (1 + \cos \theta) \cos \frac{\theta}{2} \sin \theta d\theta = 16\pi a^2 \int \cos^4 \frac{\theta}{2} \sin \frac{\theta}{2} d\theta \\ &= -\frac{32\pi a^2}{5} \cos^5 \frac{\theta}{2} + C. \end{aligned}$$

The surface formed by the revolution of the complete curve about the initial line will be obtained by taking 0 and  $\pi$  as the limits of  $\theta$  in the integral. This gives  $\frac{32\pi a^2}{5}$ .

*Any Surface. Double Integration.*

170. Let  $x, y, z$  be the co-ordinates of any point  $p$  of a surface;  $x + \Delta x, y + \Delta y, z + \Delta z$  the co-ordinates of an ad-



jacent point  $q$ . Through  $p$  draw a plane parallel to that of  $(x, z)$ , and a plane parallel to that of  $(y, z)$ ; also through  $q$  draw a plane parallel to that of  $(x, z)$  and a plane parallel to that of  $(y, z)$ . These planes will intercept an element  $pq$  of the curved surface, and the projection of this element on the plane of  $(x, y)$  will be the rectangle  $PQ$ . Suppose the tangent plane to the surface at  $p$  to be inclined to the plane of  $(x, y)$  at an angle  $\gamma$ , then it is known from solid geometry that

$$\sec \gamma = \sqrt{\left\{ 1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2 \right\}}$$

where  $\frac{dz}{dx}$  and  $\frac{dz}{dy}$  must be found from the known equation to

the surface. Now the area of  $PQ$  is  $\Delta x \Delta y$ , hence by solid geometry the area of the element of the tangent plane at  $p$  of which  $PQ$  is the projection is  $\Delta x \Delta y \sec \gamma$ . We shall assume that the limit of the sum of such terms as  $\Delta x \Delta y \sec \gamma$  for all values of  $x$  and  $y$  comprised between assigned limits is the area of the surface corresponding to those limits. Let then  $S$  denote this surface, thus

$$S = \iint \sqrt{\left\{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2\right\}} dx dy$$

the limits of the integrations being dependent upon the portion of the surface considered.

171. With respect to the point assumed in the preceding article, the reader is referred to the remarks on a similar point in *Dif. Cal.* Art. 308. He may also hereafter consult De Morgan's *Differential and Integral Calculus*, page 444, and Homersham Cox's *Integral Calculus*, page 96.

172. *Application to the Sphere.*

Let it be required to find the area of the eighth part of the surface of the sphere given by the equation

$$x^2 + y^2 + z^2 = a^2.$$

Here 
$$\frac{dz}{dx} = -\frac{x}{z}, \quad \frac{dz}{dy} = -\frac{y}{z},$$

thus 
$$S = \iint \sqrt{\left(1 + \frac{x^2}{z^2} + \frac{y^2}{z^2}\right)} dx dy = \iint \frac{a dx dy}{\sqrt{a^2 - x^2 - y^2}}.$$

Now in the figure we suppose  $OL = x$ ; put  $y_1$  for  $Ll$ , then  $y_1 = \sqrt{a^2 - x^2}$ , for the value of  $y_1$  is obtained from the equation to the surface by supposing  $z = 0$ . If we integrate with respect to  $y$  between the limits 0 and  $y_1$ , we sum up all the elements comprised in a strip of which  $LMml$  is the projection on the plane of  $(x, y)$ . Now

$$\int_0^{y_1} \frac{dy}{\sqrt{a^2 - x^2 - y^2}} = \int_0^{y_1} \frac{dy}{\sqrt{y_1^2 - y^2}} = \frac{\pi}{2};$$



thus 
$$S = \frac{\pi a}{2} \int dx.$$

If we integrate with respect to  $x$  from 0 to  $a$ , we sum up all the strips comprised in the surface of which  $OAB$  is the projection. Thus  $\frac{\pi a^2}{2}$  is the required result; and therefore the whole surface of the sphere is  $4\pi a^2$ .

If we integrate with respect to  $x$  first, we shall have

$$S = \int_0^a \int_0^{x_1} \frac{a \, dy \, dx}{\sqrt{(a^2 - x^2 - y^2)}},$$

where  $x_1 = \sqrt{(a^2 - y^2)}$ .

173. As another example let it be required to find the area of that part of the surface given by the equation

$$z^2 + (x \cos \alpha + y \sin \alpha)^2 - a^2 = 0,$$

which is situated in the positive compartment of co-ordinates. This surface is a right circular cylinder, having for its axis the line determined by  $z = 0$ ,  $x \cos \alpha + y \sin \alpha = 0$ , and  $a$  is the radius of a circular section of it. Here

$$\frac{dz}{dx} = -\frac{\cos \alpha (x \cos \alpha + y \sin \alpha)}{z},$$

$$\frac{dz}{dy} = -\frac{\sin \alpha (x \cos \alpha + y \sin \alpha)}{z},$$

thus 
$$S = \iint \frac{a \, dx \, dy}{z} = \iint \frac{a \, dx \, dy}{\sqrt{a^2 - (x \cos \alpha + y \sin \alpha)^2}}.$$

The co-ordinate plane of  $(x, y)$  cuts the surface in the straight lines  $a = \pm (x \cos \alpha + y \sin \alpha)$ , and if the upper sign be taken, we have a line lying in the positive quadrant of the plane of  $(x, y)$ .

To obtain the value of  $S$  we integrate first with respect to  $y$  between the limits  $y = 0$  and  $y = (a - x \cos \alpha) \operatorname{cosec} \alpha$ ; now

$$\int \frac{dy}{\sqrt{a^2 - (x \cos \alpha + y \sin \alpha)^2}} = \frac{1}{\sin \alpha} \sin^{-1} \frac{x \cos \alpha + y \sin \alpha}{a}$$

take this between the assigned limits, and we obtain

$$\frac{1}{\sin \alpha} \left( \frac{\pi}{2} - \sin^{-1} \frac{x \cos \alpha}{a} \right);$$

therefore 
$$S = \frac{a}{\sin \alpha} \int \left\{ \frac{\pi}{2} - \sin^{-1} \frac{x \cos \alpha}{a} \right\} dx,$$

and the limits of the integration are 0 and  $\frac{a}{\cos \alpha}$ . Hence we shall find

$$S = \frac{a^2}{\sin \alpha \cos \alpha}.$$

174. Instead of taking the element of the tangent plane at any point of a surface, so that its projection shall be the rectangle  $\Delta x \Delta y$ , it may be in some cases more convenient to take it so that its projection shall be the polar element  $r \Delta \theta \Delta r$ . Thus we shall have

$$S = \iint \sec \gamma \, r d\theta \, dr.$$

For example, suppose we require the area of the surface  $xy = az$ , which is cut off by the surface  $x^2 + y^2 = c^2$ ; here

$$\sec \gamma = \sqrt{1 + \frac{x^2}{a^2} + \frac{y^2}{a^2}} = \frac{\sqrt{a^2 + r^2}}{a} \text{ since } x^2 + y^2 = r^2.$$

Thus 
$$S = \int_0^{2\pi} \int_0^c \frac{\sqrt{a^2 + r^2}}{a} r d\theta \, dr = \frac{2\pi}{3a} \{(c^2 + a^2)^{\frac{3}{2}} - a^3\}.$$

175. Suppose  $x = r \sin \theta \cos \phi$ ,  $y = r \sin \theta \sin \phi$ ,  $z = r \cos \theta$ , so that  $r$ ,  $\theta$ ,  $\phi$  are the usual polar co-ordinates of a point in space; then we shall shew hereafter that the equation

$$S = \iint \sqrt{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2} \, dx \, dy$$

may be transformed into

$$S = \iint \sqrt{r^2 \sin^2 \theta + \left(\frac{dr}{d\theta}\right)^2 \sin^2 \theta + \left(\frac{dr}{d\phi}\right)^2} \, r d\theta \, d\phi.$$

An independent geometrical proof will be found in the *Cambridge and Dublin Mathematical Journal*, Vol. IX., and also in Carmichael's *Treatise on the Calculus of Operations*. It will be remembered that in this formula  $r = \sqrt{(x^2 + y^2 + z^2)}$ , while in Art. 174 we denote  $\sqrt{(x^2 + y^2)}$  by  $r$ .

*Approximate Values of Integrals.*

176. Suppose  $y$  a function of  $x$ , and that we require  $\int_a^c y dx$ . If the *indefinite* integral  $\int y dx$  is known we can at once ascertain the required definite integral. If the *indefinite* integral is unknown, we may still determine approximately the value of the definite integral. This process of approximation is best illustrated by supposing  $y$  to be an ordinate of a curve so that  $\int_a^c y dx$  represents a certain area. Divide  $c - a$  into  $n$  parts each equal to  $h$  and draw  $n - 1$  ordinates at equal distances between the initial and final ordinates; then the ordinates may be denoted by  $y_1, y_2, \dots, y_n, y_{n+1}$ . Hence we may take

$$h (y_1 + y_2 + \dots + y_n)$$

as an approximate value of the required area. Or we may take

$$h (y_2 + y_3, \dots + y_{n+1})$$

as an approximate value.

We may obtain another approximation thus; suppose the extremities of the  $r^{\text{th}}$  and  $r + 1^{\text{th}}$  ordinates joined; thus we have a trapezium, the area of which is  $(y_r + y_{r+1}) \frac{h}{2}$ . The sum of all such trapeziums gives as an approximate value of the area

$$h \left\{ \frac{y_1}{2} + y_2 + y_3, \dots + y_n + \frac{y_{n+1}}{2} \right\}.$$

This result is in fact half the sum of the two former results. It is obvious we may make the approximation as close as we please by sufficiently increasing  $n$ .

177. The following is another method of approximation. Let a parabola be drawn having its axis parallel to that of  $y$ ; let  $y_1, y_2, y_3$  represent three equidistant ordinates,  $h$  the distance between  $y_1$  and  $y_2$ , and therefore also between  $y_2$  and  $y_3$ . Then it may be proved that the area contained between the parabola, the axis of  $x$ , and the two extreme ordinates is

$$\frac{h}{3} (y_1 + 4y_2 + y_3).$$

This will be easily shewn by a figure, as the area consists of a trapezium and a parabolic segment, and the area of the latter is known by Art. 143.

Let us now suppose that  $n$  is even, so that the area we have to estimate is divided into an even number of pieces. Then assume that the area of the first two pieces is

$$\frac{h}{3} (y_1 + 4y_2 + y_3),$$

that the area of the third and fourth pieces is

$$\frac{h}{3} (y_3 + 4y_4 + y_5),$$

and so on. Thus we shall have finally as an approximate result

$$\frac{h}{3} \{y_1 + 2(y_3 + y_5 + \dots + y_{n-1}) + y_{n+1} + 4(y_2 + y_4 + \dots + y_n)\}.$$

Hence we have the following rule: add together the first ordinate, the last ordinate, twice the sum of all the other odd ordinates, and four times the sum of all the even ordinates; then multiply the result by one-third the common distance of the ordinates.

### EXAMPLES.

1. If  $A$  denote the area contained between the catenary, the axis of  $x$ , the axis of  $y$ , and an ordinate at the extremity of the arc  $s$ , shew that  $A = cs$ . The arc  $s$  begins at the lowest point of the curve.

2. The whole area of the curve

$$\left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} = 1,$$

is  $\frac{3}{8}\pi ab$ . (The integration may be effected by assuming  $x = a \cos^3 \phi$ .)

3. The area of the curve  $y(x^2 + a^2) = c^2(a - x)$  from  $x = 0$  to  $x = a$  is  $c^2\left(\frac{\pi}{4} - \frac{1}{2}\log 2\right)$ .
4. The area of the curve  $y^2x = 4a^2(2a - x)$  from  $x = 0$  to  $x = 2a$  is  $4\pi a^2$ .
5. Shew that the whole area of the curve  $y^2 = \frac{x^2(a+x)}{a-x}$ , supposing it bounded on one side by the asymptote, is  $4a^2$ . (Estimate the area of the loop and the other portion separately.)
6. Find the whole area between the curve  $y^2(x^2 + a^2) = a^2x^2$  and its asymptotes. *Result.*  $4a^2$ .
7. Find the whole area between the curve  $xy^2 = 4a^2(2a - x)$  and its asymptote. *Result.*  $4\pi a^2$ .
8. Find the whole area between the curve  $y^2(2a - x) = x^3$  and its asymptote. *Result.*  $3\pi a^2$ .
9. Find the whole area of the curve  $y = x \pm \sqrt{a^2 - x^2}$ . *Result.*  $\pi a^2$ .
10. Find the area included between the curves
- $$y^2 - 4ax = 0, \quad x^2 - 4ay = 0. \quad \text{Result. } \frac{16a^2}{3}.$$
11. Find the whole area of the curve  $a^4y^2 + b^2x^4 = a^2b^2x^2$ . *Result.*  $\frac{4}{3}ab$ .

12. Find the area of the loop of the curve  $a^3y^4 = x^4(a^3 - x^3)$ .

$$\text{Result. } \frac{4a^3}{5}.$$

13. The area between the tractory, the axis of  $y$ , and the asymptote is  $\frac{\pi c^2}{4}$ . (See Art. 100).

14. Find the area of the loop of the curve

$$y^2(a^2 + x^2) = x^2(a^2 - x^2). \quad \text{Result. } \frac{a^2}{2}(\pi - 2).$$

15. Find the area of the loop of the curve

$$16a^4y^2 = b^2x^2(a^2 - 2ax). \quad \text{Result. } \frac{ab}{30}.$$

16. Find the area of the loop of the curve

$$2y^2(a^2 + x^2) = (a^2 - x^2)^2.$$

$$\text{Result. } a^2\{3\sqrt{2} \log(1 + \sqrt{2}) - 2\}.$$

17. Find the whole area of the curve

$$2y^2(a^2 + x^2) - 4ay(a^2 - x^2) + (a^2 - x^2)^2 = 0.$$

$$\text{Result. } a^2\pi \left\{4 - \frac{5\sqrt{2}}{2}\right\}.$$

18. Find the area of the curve

$$y = c \sin \frac{x}{a} \cdot \log \sin \frac{x}{a}$$

from  $x=0$  to  $x=a\pi$ .  $\text{Result. } 2ac(1 - \log 2).$

19. Find the area of the curve  $\frac{y}{c} = \left(\frac{x}{a}\right)^n$  between  $x=\alpha$  and  $x=\beta$ , and from the result deduce the area of the hyperbola  $xy = a^2$  between the same limits.

20. Find the area of the ellipse whose equation is

$$ax^2 + 2bxy + cy^2 = 1. \quad \text{Result. } \frac{\pi}{\sqrt{(ac - b^2)}}.$$

21. Find the area of a loop of the curve
- $r^2 = a^2 \cos 2\theta$
- .

$$\text{Result. } \frac{a^2}{2}.$$

22. Find the area contained by all the loops of the curve

$$r = a \sin n\theta.$$

$$\text{Result. } \frac{\pi a^2}{4} \text{ or } \frac{\pi a^2}{2} \text{ according as } n \text{ is odd or even.}$$

23. Find the area between the curves
- $r = a \cos n\theta$
- and
- $r = a$
- .

24. Find the area of a loop of the curve
- $r^2 \cos \theta = a^2 \sin 3\theta$
- .

$$\text{Result. } \frac{3a^2}{4} - \frac{a^2}{2} \log 2.$$

25. Find the whole area of the curve
- $r = a (\cos 2\theta + \sin 2\theta)$
- .

$$\text{Result. } \pi a^2.$$

26. Find the area of a loop of the curve
- $(x^2 + y^2)^3 = 4a^2 x^2 y^2$
- .

$$\text{Result. } \frac{\pi a^2}{8}.$$

27. Find the whole area of the curve

$$(x^2 + y^2)^2 = 4a^2 x^2 + 4b^2 y^2. \quad \text{Result. } 2\pi (a^2 + b^2).$$

28. Find the whole area of the curve

$$\frac{x^2}{a^4} + \frac{y^2}{b^4} = \frac{1}{c^2} \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} \right)^2. \quad \text{Result. } \frac{\pi c^2}{2ab} (a^2 + b^2).$$

29. Find the area of the loop of the curve

$$y^3 - 3axy + x^3 = 0. \quad \text{Result. } \frac{3a^3}{2}.$$

30. Find the area of the loop of the curve

$$r \cos \theta = a \cos 2\theta. \quad \text{Result. } \left( 2 - \frac{\pi}{2} \right) a^2.$$

31. Find the area of the curve

$$r = \frac{a^2}{\sqrt{(a^2 - b^2 \cos^2 \theta)}} + b \cos \theta,$$

$a$  being  $> b$ .

*Result.*  $\frac{\pi a^2}{\sqrt{(a^2 - b^2)}} + \frac{\pi b^2}{2}.$

32. In a logarithmic spiral find the area between the curve and two radii vectores drawn from the pole.
33. Find the area between the conchoid  $r = a + b \operatorname{cosec} \theta$  and two radii vectores drawn from the pole.
34. In an ellipse find the area between the curve and two radii vectores drawn from the centre.
35. In a parabola find the area between the curve and two radii vectores drawn from the vertex.
36. Find the area included between the curve

$$r = a (\sec \theta + \tan \theta)$$

and its asymptote  $r \cos \theta = 2a$ . *Result.*  $\left(\frac{\pi}{2} + 2\right) a^2.$

37. The whole area of the curve  $r = a(2 \cos \theta + 1)$  is  $a^2 \left(2\pi + \frac{3\sqrt{3}}{2}\right)$ , and the area of the inner loop is  $a^2 \left(\pi - \frac{3\sqrt{3}}{2}\right)$ .

38. Find the whole area of the curve  $r = a \cos \theta + b$  where  $a$  is greater than  $b$ . Also find the area of the inner loop.

39. If  $x$  and  $y$  be the co-ordinates of an equilateral hyperbola  $x^2 - y^2 = a^2$ , shew that

$$x = \frac{a}{2} \left( e^{\frac{2u}{a^2}} + e^{-\frac{2u}{a^2}} \right), \quad y = \frac{a}{2} \left( e^{\frac{2u}{a^2}} - e^{-\frac{2u}{a^2}} \right),$$

where  $u$  is the area intercepted between the curve, the central radius vector, and the axis.



40. Find the whole area of the curve which is the locus of the intersection of two normals to an ellipse at right angles.

*Result.*  $\pi (a-b)^2$ .

It may be shewn that the equation to the curve is

$$r^2 (a^2 + b^2) (a^2 \sin^2 \theta + b^2 \cos^2 \theta)^2 \\ = (a^2 - b^2)^2 (a^2 \sin^2 \theta - b^2 \cos^2 \theta)^2.$$

41. Find the area included within any arc traced by the extremity of the radius vector of a spiral in a complete revolution, and the straight line joining the extremities of the arc. If, for example, the equation to the spiral be  $r = a \left(\frac{\theta}{2\pi}\right)^n$ , prove that the area corresponding to any value of  $\theta$  greater than  $2\pi$  is

$$\frac{\pi a^2}{2n+1} \left\{ \left(\frac{\theta}{2\pi}\right)^{2n+1} - \left(\frac{\theta}{2\pi} - 1\right)^{2n+1} \right\}.$$

42. Find the area contained between a parabola, its evolute, and two radii of curvature of the parabola. (Art. 159.)
43. Find the area contained between a cycloid, its evolute, and two radii of curvature of the cycloid.
44. Find the area of the surface generated by the revolution round the axis of  $x$  of the curve  $xy = k^2$ .

45. Also of the curve  $y = ae^{\frac{x}{c}}$ .

46. Also of the catenary  $y = \frac{c}{2} (e^{\frac{x}{c}} + e^{-\frac{x}{c}})$ .

47. Shew that the whole surface of an oblate spheroid is

$$2\pi a^2 \left\{ 1 + \frac{1-e^2}{2e} \log \frac{1+e}{1-e} \right\}.$$

48. A cycloid revolves round the tangent at the vertex; shew that the whole surface generated is  $\frac{32}{3} \pi a^2$ .

- I A cycloid revolves round its base; shew that the whole surface generated is  $\frac{64}{3}\pi a^2$ .
50. A cycloid revolves round its axis; shew that the whole surface generated is  $8\pi a^2(\pi - \frac{2}{3})$ .
51. The whole surface generated by the revolution of the tractory round the axis of  $x$  is  $4\pi c^2$ .
52. A sphere is pierced perpendicularly to the plane of one of its great circles by two right cylinders, of which the diameters are equal to the radius of the sphere and the axes pass through the middle points of two radii that compose a diameter of this great circle. Find the surface of that portion of the sphere not included within the cylinders.

*Result.* Twice the square of the diameter of the sphere.

53. Find the surface generated by the portion of the curve  $y = a \pm a \log \frac{x}{a}$  between the limits  $x = a$  and  $x = ae$ .

*Result.*  $4\pi a^2 \left\{ 1 + \sqrt{1 + e^2} - \sqrt{2} + \log \frac{1 + \sqrt{2}}{1 + \sqrt{1 + e^2}} \right\}$ .


54. Find  $\int \frac{dS}{p}$ , where  $dS$  represents an element of surface, and  $p$  the perpendicular from the origin upon the tangent plane of the element, the integral being extended over the whole of the ellipsoid  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ .

*Result.*  $\frac{4\pi}{3abc} (a^2b^2 + b^2c^2 + c^2a^2)$ .

## CHAPTER VIII.

## VOLUMES OF SOLIDS.

*Formulae involving Single Integration. Solid of Revolution.*



178. LET  $A$  be a fixed point in a curve  $APQ$ , and  $P$  any other point on the curve whose co-ordinates are  $x$  and  $y$ . Let the curve revolve round the axis of  $x$ , and let  $V$  denote the volume of the solid bounded by the surface generated by the curve and by two planes perpendicular to the axis of  $x$ , one through  $A$  and the other through  $P$ ; then (*Dif. Cal. Art. 314*),

$$\frac{dV}{dx} = \pi y^2,$$

therefore

$$V = \int \pi y^2 dx.$$

From the equation to the curve  $y$  is a known function of  $x$ ; suppose  $\psi(x)$  to be the integral of  $\pi y^2$ ; then

$$V = \psi(x) + C.$$

Let  $V_1$  denote the volume when the point  $P$  has  $x_1$  for its abscissa, and  $V_2$  the volume when the point  $P$  has  $x_2$  for its abscissa; thus

$$V_1 = \psi(x_1) + C,$$

$$V_2 = \psi(x_2) + C,$$

therefore 
$$V_2 - V_1 = \psi(x_2) - \psi(x_1) = \pi \int_{x_1}^{x_2} y^2 dx.$$

179. *Application to the Right Circular Cone.*

Let a straight line pass through the origin and make an angle  $\alpha$  with the axis of  $x$ ; then this straight line will generate a right circular cone by revolving round the axis of  $x$ . Here  $y = x \tan \alpha$ ; thus

$$V = \int \pi \tan^2 \alpha x^2 dx = \frac{\pi \tan^2 \alpha}{3} x^3 + C,$$

$$V_2 - V_1 = \frac{\pi \tan^2 \alpha}{3} (x_2^3 - x_1^3).$$

Suppose  $x_1 = 0$ , and let  $r = x_2 \tan \alpha$ ; thus the volume becomes  $\frac{\pi \tan^2 \alpha x_2^3}{3}$ , that is,  $\frac{\pi r^2 x_2}{3}$ . Hence the volume of a right circular cone is one-third the product of the area of the base into the altitude.

180. *Application to the Sphere.*

Here taking the origin at the centre of the sphere we have  $y^2 = a^2 - x^2$ ; thus

$$\int \pi y^2 dx = \pi \left( a^2 x - \frac{x^3}{3} \right) + C.$$

The volume of a hemisphere  $= \int_0^a \pi y^2 dx = \frac{2\pi a^3}{3}.$

181. *Application to the Paraboloid.*

Here the generating curve is the parabola, so that

$$y^2 = 4ax.$$

Thus 
$$V_2 - V_1 = \pi \int_{x_1}^{x_2} 4ax \, dx = 2a\pi (x_2^2 - x_1^2).$$

Suppose  $x_1 = 0$ , then the volume becomes  $2a\pi x_2^2$ , that is  $\frac{1}{2} \pi y_2^2 x_2$ , where  $y_2^2 = 4ax_2$ ; thus the volume is half that of a cylinder which has the same height, namely  $x_2$ , and the same base, namely a circle of which  $y_2$  is the radius.

182. *Application to the Solid formed by a Cycloid.*

Let a cycloid revolve round its axis; here (*Dif. Cal. Art. 358*),

$$y = \sqrt{(2ax - x^2)} + a \operatorname{vers}^{-1} \frac{x}{a}.$$

The integration is best effected by putting for  $x$  and  $y$  their values in terms of  $\theta$ , (*Dif. Cal. Art. 358*.) Thus

$$\pi \int y^2 dx = \pi a^3 \int (\theta + \sin \theta)^2 \sin \theta \, d\theta.$$

To obtain the volume generated by a semi-cycloid the limits for  $x$  would be 0 and  $2a$ ; thus the corresponding limits for  $\theta$  are 0 and  $\pi$ .

Now 
$$\int \theta^2 \sin \theta \, d\theta = -\theta^2 \cos \theta + 2 \int \theta \cos \theta \, d\theta$$

$$= -\theta^2 \cos \theta + 2\theta \sin \theta + 2 \cos \theta,$$

therefore 
$$\int_0^\pi \theta^2 \sin \theta \, d\theta = \pi^2 - 4$$

$$2 \int \theta \sin^2 \theta \, d\theta = \int \theta (1 - \cos 2\theta) \, d\theta = \frac{\theta^2}{2} - \frac{\theta \sin 2\theta}{2} - \frac{\cos 2\theta}{4},$$

therefore 
$$2 \int_0^\pi \theta \sin^2 \theta \, d\theta = \frac{\pi^2}{2}.$$

And 
$$\int_0^\pi \sin^3 \theta \, d\theta = 2 \int_0^{\frac{\pi}{2}} \sin^3 \theta \, d\theta = 2 \cdot \frac{2}{3}. \quad (\text{Art. 35.})$$

Thus the required volume

$$= \pi a^3 \left\{ \pi^2 - 4 + \frac{\pi^2}{2} + \frac{4}{3} \right\} = \pi a^3 \left( \frac{3\pi^2}{2} - \frac{8}{3} \right).$$

183. This formula for the volume of a solid of revolution,  $V = \int \pi y^2 dx$ , like others which we have noticed, is one, the truth of which is obvious, as soon as the notation of the Integral Calculus is understood. In the figure to Art. 128, if  $PM$  be  $y$  and  $MN$  be denoted by  $\Delta x$ , then  $\pi y^2 \Delta x$  is the volume of the solid generated by the revolution of  $MN_pP$  about the axis of  $x$ . Thus  $\Sigma \pi y^2 \Delta x$  will differ from the volume generated by the revolution of  $ADEB$  by the sum of such volumes as are generated by  $PpQ$ , and the latter sum will vanish in the limit. Thus the volume generated by the revolution of  $ADEB$  is equal to the limit of  $\Sigma \pi y^2 \Delta x$ , that is, to  $\int \pi y^2 dx$ .

184. Similarly, if  $V$  denote the volume bounded by the surface formed by a curve which revolves round the axis of  $y$ , and by planes perpendicular to the axis of  $y$ , we shall have

$$V = \int \pi x^2 dy.$$

And, as in Art. 178, we shall have

$$V_2 - V_1 = \int_{y_1}^{y_2} \pi x^2 dy.$$

185. Suppose two curves to revolve round the axis of  $x$ , and thus to generate two surfaces, and that we require the *difference* of two volumes, one bounded by the first surface and by planes perpendicular to the axis of  $x$ , and the other bounded by the second surface and by the planes already assigned. Let  $y = \phi(x)$  be the equation to the first curve, and  $y = \psi(x)$  that to the second. Then if  $V$  denote the required difference, we have

$$\begin{aligned} V &= \int \pi \{ \phi(x) \}^2 dx - \int \pi \{ \psi(x) \}^2 dx \\ &= \pi \int [ \{ \phi(x) \}^2 - \{ \psi(x) \}^2 ] dx. \end{aligned}$$

If the planes which bound the required volume are determined by  $x = x_1$  and  $x = x_2$ , we must integrate between the limits  $x_1$  and  $x_2$  for  $x$ .

186. Suppose, for example, that a closed curve is such that the line  $y = a$  bisects every ordinate parallel to the axis of  $y$ ; then we have  $\phi(x) = a + \chi(x)$  and  $\psi(x) = a - \chi(x)$  where  $\chi(x)$  denotes some function of  $x$ . Thus

$$\{\phi(x)\}^2 - \{\psi(x)\}^2 = 4a\chi(x),$$

and

$$V = \pi \int_{x_1}^{x_2} 4a\chi(x) dx.$$

Suppose the abscissæ of the extreme points of the curve are  $x_2$  and  $x_1$ , then the volume generated by the revolution of the closed curve round the axis of  $x$  is  $4a\pi \int_{x_1}^{x_2} \chi(x) dx$ .

And  $2 \int_{x_1}^{x_2} \chi(x) dx$  is the area of the closed curve, so that the volume is equal to the product of  $2a\pi$  into the area. This demonstration supposes that the generating curve lies entirely on one side of the axis of  $x$ .

If the generating curve be the circle given by

$$(x - h)^2 + (y - k)^2 = c^2,$$

we have  $\pi c^2$  for its area, and therefore  $2kc^2\pi^2$  for the volume generated by the revolution of it round the axis of  $x$ .

187. In a similar way if the curves  $x = \phi(y)$ ,  $x = \psi(y)$ , revolve round the axis of  $y$  we obtain for the volume bounded by these surfaces and by planes perpendicular to the axis of  $y$

$$V = \pi \int [\{\phi(y)\}^2 - \{\psi(y)\}^2] dy.$$

188. The method given in Art. 178 for finding the volume of a *solid of revolution* may be adapted to *any* solid. The method may be described thus: conceive the solid cut up into thin slices by a series of parallel planes, estimate approximately the volume of each slice and add these volumes; the limit of this sum when each slice becomes indefinitely thin is

the volume of the solid required. Suppose that a solid is cut up into slices by planes perpendicular to the axis of  $x$ ; let  $\phi(x)$  be the area of a section of the solid made by a plane which is at a distance  $x$  from the origin, and let  $x + \Delta x$  be the distance of the next plane from the origin; thus these two planes intercept a slice of which the thickness is  $\Delta x$ , and of which the volume may be represented by  $\phi(x) \Delta x$ . The volume of the solid will therefore be the limit of  $\Sigma \phi(x) \Delta x$ , that is, it will be  $\int \phi(x) dx$ ; the limits of the integration will depend upon the particular solid or portion of a solid under consideration.

189. *Application to an Ellipsoid.*

The equation to the ellipsoid is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1;$$

if a section be made by a plane perpendicular to the axis of  $x$  at a distance  $x$  from the origin, the boundary of the section is an ellipse, of which the semi-axes are  $b \sqrt{1 - \frac{x^2}{a^2}}$  and  $c \sqrt{1 - \frac{x^2}{a^2}}$ ; hence the area of this ellipse is  $\pi bc \left(1 - \frac{x^2}{a^2}\right)$ ; this is therefore the value of  $\phi(x)$ . Hence the volume of the ellipsoid

$$= \int_{-a}^a \pi bc \left(1 - \frac{x^2}{a^2}\right) dx = \frac{4\pi abc}{3}.$$

190. *Application to a Pyramid.*

Let there be a pyramid, the base of which is any rectilinear figure; let  $A$  be the area of the base and  $h$  the height. Take the origin of co-ordinates at the vertex of the pyramid, and the axis of  $x$  perpendicular to the base of the pyramid, then the volume of the pyramid

$$= \int_0^h \phi(x) dx.$$



Now the section of the pyramid made by any plane parallel to the base is a rectilinear figure similar to the base, and the areas of similar figures are as the squares of their homologous sides; hence we infer that

$$\phi(x) = \frac{x^2}{h^2} A.$$

Thus the volume of the pyramid

$$= \frac{A}{h^2} \int_0^h x^2 dx = \frac{Ah}{3}.$$

This investigation also holds for a cone, the base of which is any closed curve.

191. As an example we will find the volume lying between an hyperboloid of one sheet, its asymptotic cone and two planes perpendicular to their common axis.

Let the equation to the hyperboloid be

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} - \frac{z^2}{c^2} + 1 = 0,$$

and that to the cone


$$\frac{x^2}{a^2} - \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0.$$

If a section of the former surface be made by a plane perpendicular to the axis of  $x$  and at a distance  $x$  from the origin, the boundary is an ellipse of which the area is  $\pi bc \left( \frac{x^2}{a^2} + 1 \right)$ ; the section of the second surface made by the same plane also has an ellipse for its boundary, and its area is  $\frac{\pi bc x^2}{a^2}$ . Therefore the difference of the areas is  $\pi bc$ . Hence the required volume, supposing it bounded by the planes  $x = x_1$  and  $x = x_2$  is

$$\int_{x_1}^{x_2} \pi bc dx, \quad \text{that is, } \pi bc (x_2 - x_1).$$

192. Sometimes it may be convenient to make sections by parallel planes not perpendicular to the axis of  $x$ . If  $\alpha$  be the inclination of the axis of  $x$  to the parallel planes, then  $\phi(x) \sin \alpha \Delta x$  may be taken as the volume of a slice and the integration performed as before.

*Formula involving Double Integration.*



193. We will first give a formula for the volume of a solid of revolution. In the figure, let  $x, y$  be the co-ordinates of  $s$ , and  $x + \Delta x, y + \Delta y$  those of  $t$ . Suppose the whole figure to revolve round the axis of  $x$ , then the element  $st$  will generate a ring, the volume of which will be ultimately  $2\pi y \Delta x \Delta y$ : this follows from the consideration that  $\Delta x \Delta y$  is the area of  $st$  and  $2\pi y$  the perimeter of the circle described by  $s$ . Hence the volume generated by the figure  $BEeb$ , or by any portion of it, will be the limit of the sum of such terms as  $2\pi y \Delta x \Delta y$ . Let  $V$  denote the required volume, then

$$V = 2\pi \iint y \, dx \, dy;$$

the limits of the integration being so taken as to include all the elements of the required volume.

194. Suppose the volume required that which is obtained by the revolution of all the figure  $BEeb$ ; let  $y = \phi(x)$  be the

equation to the upper curve,  $y = \psi(x)$  that to the lower curve, and let  $OC = x_1$ ,  $OH = x_2$ . We should then integrate first with respect to  $y$  between the limits  $y = \psi(x)$  and  $y = \phi(x)$ ; we thus sum up all the elements like  $2\pi y \Delta x \Delta y$  which are contained in the solid formed by the revolution of the strip  $PQqp$ ; then we integrate with respect to  $x$  between the limits  $x_1$  and  $x_2$ . Thus to express the operation symbolically

$$\begin{aligned} V &= 2\pi \int_{x_1}^{x_2} \int_{\psi(x)}^{\phi(x)} y \, dx \, dy \\ &= \pi \int_{x_1}^{x_2} [\{\phi(x)\}^2 - \{\psi(x)\}^2] \, dx. \end{aligned}$$

The second expression is obtained by effecting the integration with respect to  $y$  between the assigned limits, and it coincides with that already obtained in Art. 185.

195. Thus in the preceding article we divide the solid into elementary rings, of which  $2\pi y \Delta x \Delta y$  is the type; in the first integration we collect a number of these rings, so as to form a figure, which is the difference of two concentric circular slices; in the second integration we collect all these figures and thus obtain the volume of the required solid.

196. Suppose the figure which revolves round the axis of  $x$  to be bounded by the curves  $x = \phi(y)$  and  $x = \psi(y)$ , and by the straight lines  $y = y_1$  and  $y = y_2$ ; then in applying the formula for  $V$  it will be convenient to integrate first with respect to  $x$ ; thus

$$V = 2\pi \int_{y_1}^{y_2} \int_{\psi(y)}^{\phi(y)} y \, dy \, dx.$$

In this case in the integration with respect to  $x$  we collect all the elements like  $2\pi y \Delta y \Delta x$  which have the same radius  $y$ , so that the sum of the elements is a thin cylindrical shell, of which  $\Delta y$  is the thickness,  $y$  is the radius, and  $\phi(y) - \psi(y)$  the height. Thus

$$V = 2\pi \int_{y_1}^{y_2} \{\phi(y) - \psi(y)\} y \, dy.$$

197. As an example of the preceding formulæ, let it be required to find the volume of the solid generated by the revolution of the area  $ALB$  round the axis of  $x$  in the figure already given in Art. 141. This volume is the excess of the hemisphere generated by the revolution of  $SLB$  over the paraboloid generated by the revolution of  $ASL$ ; the result is therefore known, and we propose the example, not for the sake of the result, but for illustration of the formulæ of double integration.

Suppose the positive direction of the axis of  $x$  to the left, then the equation to  $AL$  is  $y^2 = 4a(a - x)$  and that to  $BL$  is  $y^2 = 4a^2 - x^2$ . Let  $V$  be the required volume, then

$$V = \int_0^{2a} \int_{\frac{4a^2 - y^2}{4a}}^{\sqrt{4a^2 - y^2}} 2\pi y \, dy \, dx.$$

If we wish to integrate with respect to  $y$  first, we must, as in Art. 141, suppose the figure  $ALB$  divided into two parts; thus

$$V = \int_0^a \int_{\sqrt{4a^2 - x^2}}^{\sqrt{4a^2 - x^2}} 2\pi y \, dx \, dy + \int_a^{2a} \int_0^{\sqrt{4a^2 - x^2}} 2\pi y \, dx \, dy.$$

Again, let it be required to find the volume generated by the revolution of  $LDC$  about the axis of  $x$ . Let the positive direction of the axis of  $x$  be now to the right, then the equation to  $LC$  is  $y^2 = 4a(a + x)$  and that to  $LD$  is  $y^2 = 4a^2 - x^2$ . Let  $V$  be the required volume, then

$$V = \int_0^{2a} \int_{\sqrt{4a^2 - x^2}}^{\sqrt{4a^2 + 4ax}} 2\pi y \, dx \, dy.$$


If we wish to integrate with respect to  $x$  first, we must, as in Art. 141, suppose the figure  $LDC$  divided into two parts; thus

$$V = \int_0^{2a} \int_0^{2a} 2\pi y \, dy \, dx + \int_{2a}^{2a\sqrt{3}} \int_{\frac{y^2 - 4a^2}{4a}}^{2a} 2\pi y \, dy \, dx.$$

198. Similarly, if a solid is formed by the revolution of a curve round the axis of  $y$ , we have

$$V = \iint 2\pi x \, dy \, dx.$$

199. We now proceed to consider any solid.



Let  $x, y, z$  be the co-ordinates of any point  $p$  of a surface,  $x + \Delta x, y + \Delta y, z + \Delta z$  the co-ordinates of an adjacent point  $q$ . Through  $p$  draw planes parallel to the co-ordinate planes of  $(x, z)$  and  $(y, z)$ ; through  $q$  also draw planes parallel to the same co-ordinate planes. These four planes will include between them a column, of which  $PQ$  is the base and  $Pp$  the height. The volume of this column will be ultimately  $z \Delta x \Delta y$ , and the volume between an assigned portion of the given surface and the plane of  $(x, y)$  will be found by taking the limit of the sum of a series of terms like  $z \Delta x \Delta y$ . Let  $V$  denote this volume, then

$$V = \iint z \, dx \, dy.$$

The equation to the surface gives  $z$  as a function of  $x$  and  $y$ ; the limits of the integration must be taken so as to include all the elements of the proposed solid.

If we integrate first with respect to  $y$ , we sum up the columns which form a slice comprised between two planes perpendicular to the axis of  $x$ ; thus the limits of integration with respect to  $y$  may be functions of  $x$ , and we shall obtain

$$\int z \, dy = f(x),$$

where  $f(x)$  is in fact the area of the section of the solid considered made by a plane perpendicular to the axis of  $x$  at a distance  $x$  from the origin. Then finally

$$V = \int f(x) \, dx;$$

this coincides with the formula already given in Art. 188.

200. *Application to the Ellipsoid.*

Let it be required to find the volume of the eighth part of the ellipsoid determined by the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$

Here we have to find

$$\iint c \sqrt{\left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}\right)} \, dx \, dy.$$

First integrate with respect to  $y$ , then the limits of  $y$  are 0 and  $Ll$ , that is 0 and  $b \sqrt{\left(1 - \frac{x^2}{a^2}\right)}$ ; we thus obtain the sum of all the columns which form the slice between the planes  $Lpl$  and  $Mqm$ . Now between the assigned limits

$$\int \sqrt{\left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}\right)} \, dy = \frac{\pi b}{4} \left(1 - \frac{x^2}{a^2}\right);$$

thus 
$$V = \int \frac{\pi}{4} bc \left(1 - \frac{x^2}{a^2}\right) \, dx.$$

The limits of  $x$  are 0 and  $a$ ; we thus obtain the sum of

all the slices which are comprised in the solid  $OABC$ . Hence

$$V = \frac{\pi abc}{6}.$$

201. Suppose the given surface to be determined by  $xy = az$ , and we require the volume bounded by the plane of  $(x, y)$ , by the given surface, and by the four planes  $x = x_1$ ,  $x = x_2$ ,  $y = y_1$ ,  $y = y_2$ . Here the volume is given by

$$\begin{aligned} V &= \int_{x_1}^{x_2} \int_{y_1}^{y_2} \frac{xy}{a} dx dy \\ &= \frac{1}{4a} (y_2^2 - y_1^2) (x_2^2 - x_1^2) \\ &= \frac{1}{4a} (x_2 - x_1) (y_2 - y_1) \{x_1 y_1 + x_2 y_2 + x_1 y_2 + x_2 y_1\} \\ &= \frac{1}{4} (x_2 - x_1) (y_2 - y_1) (z_1 + z_2 + z_3 + z_4), \end{aligned}$$

where  $z_1, z_2, z_3, z_4$  are the ordinates of the four corner points of the selected portion.

202. Find the volume comprised between the plane  $z = 0$  and the surfaces  $xy = az$  and  $(x - h)^2 + (y - k)^2 = c^2$ .

Here we have to integrate  $\iint \frac{xy}{a} dx dy$  between limits determined by  $(x - h)^2 + (y - k)^2 = c^2$ .

Now  $\int y dy = \frac{y^2}{2}$ , and the limits of  $y$  are

$$k - \sqrt{c^2 - (x - h)^2} \text{ and } k + \sqrt{c^2 - (x - h)^2}.$$

Thus we obtain

$$2k \sqrt{c^2 - (x - h)^2}.$$

Thus finally the required volume

$$= \frac{2k}{a} \int x \sqrt{c^2 - (x - h)^2} dx,$$

where the limits of  $x$  are  $h - c$  and  $h + c$ .

And

$$\int x \sqrt{c^2 - (x-h)^2} dx = \int (x-h) \sqrt{c^2 - (x-h)^2} dx + h \int \sqrt{c^2 - (x-h)^2} dx.$$

Put  $x-h=t$ ; thus we obtain

$$\int t \sqrt{c^2 - t^2} dt + h \int \sqrt{c^2 - t^2} dt.$$

The limits of  $t$  are  $-c$  and  $+c$ ; therefore the result is  $\frac{hc^2\pi}{2}$ ; and the required volume is  $\frac{hkc^2\pi}{a}$ .

203. Instead of dividing a solid into columns standing on rectangular bases, so that  $z \Delta x \Delta y$  is the volume of the column, we may divide it into columns standing upon the polar element of area; hence  $z r \Delta \theta \Delta r$  is the volume of the column. Therefore for the volume  $V$  of a solid we have the formula

$$V = \iint z r d\theta dr.$$

From the equation to the surface  $z$  must be expressed as a function of  $r$  and  $\theta$ .

204. Required the volume of the solid comprised between the plane of  $(x, y)$  and the surface whose equation is

$$z = ae^{-\frac{x^2+y^2}{c^2}}.$$

Here, since  $x^2 + y^2 = r^2$

$$V = a \iint e^{-\frac{r^2}{c^2}} r d\theta dr.$$

The surface extends to an infinite distance from the origin in every direction; thus the limits of  $\theta$  are 0 and  $2\pi$ , and those of  $r$  are 0 and  $\infty$ .

Now 
$$\int e^{-\frac{r^2}{c^2}} r dr = -\frac{e^{-\frac{r^2}{c^2}}}{2} c^2;$$



thus 
$$\int_0^{\infty} e^{-\frac{r^2}{c^2}} r dr = \frac{c^2}{2}.$$

And 
$$\int_0^{2\pi} d\theta = 2\pi.$$

Hence the required volume is  $\pi ac^3$ .

*Formulae involving Triple Integration.*

205. In the figure to Art. 199, suppose we draw a series of planes perpendicular to the axis of  $z$ ; let  $z$  be the distance of one plane from the origin and  $z + \Delta z$  the distance of the next. These planes intercept from the column  $pqPQ$  an elementary rectangular parallelepiped, the volume of which is  $\Delta x \Delta y \Delta z$ . The whole solid may be considered as the limit of the sum of such elements. Hence if  $V$  denote its volume

$$V = \iiint dx dy dz.$$

206. Required the volume of a portion of the cylinder determined by the equation

$$x^2 + y^2 - 2ax = 0,$$

which is intercepted between the planes

$$z = x \tan \alpha \text{ and } z = x \tan \beta.$$

Here if  $y_1$  stand for  $\sqrt{(2ax - x^2)}$ , we have

$$\begin{aligned} V &= \int_0^{2a} \int_{-y_1}^{y_1} \int_{x \tan \alpha}^{x \tan \beta} dx dy dz. \\ &= \int_0^{2a} \int_{-y_1}^{y_1} (\tan \beta - \tan \alpha) x dx dy \\ &= 2 (\tan \beta - \tan \alpha) \int_0^{2a} x \sqrt{(2ax - x^2)} dx \\ &= 2 (\tan \beta - \tan \alpha) \frac{\pi a^3}{2}. \end{aligned}$$

207. The polar element of plane area is, as we have seen in previous articles,  $r\Delta\theta\Delta r$ . Suppose this to revolve round the initial line through an angle  $2\pi$ , then a solid ring is generated, of which the volume is  $2\pi r \sin\theta r\Delta\theta\Delta r$ , since  $2\pi r \sin\theta$  is the circumference of the circle described by the point whose polar co-ordinates are  $r$  and  $\theta$ . Let  $\phi$  denote the angle which the plane of the element in any position makes with the initial position of the plane,  $\phi + \Delta\phi$  the angle which the plane in a consecutive position makes with the initial plane; then the part of the solid ring which is intercepted between the revolving plane in these two positions is to the whole ring in the same proportion as  $\Delta\phi$  is to  $2\pi$ . Hence the volume of this intercepted part is

$$r^3 \sin\theta \Delta\phi \Delta\theta \Delta r.$$

This is therefore an expression in polar co-ordinates for an element of any solid. Hence the volume of the whole solid may be found by taking the limit of the sum of such elements; that is, if  $V$  denote the required volume,

$$V = \iiint r^3 \sin\theta \, d\phi \, d\theta \, dr.$$

The limits of the integration must be so taken as to include in the integration all the elements of the proposed solid. The student will remember that  $r$  denotes the distance of any point from the origin,  $\theta$  the angle which this distance makes with some fixed line through the origin, and  $\phi$  the angle which the plane passing through this distance and the fixed line makes with some fixed plane passing through the fixed line.

208. Suppose, for example, that we apply the formula to find the volume of the eighth part of a sphere. Integrate with respect to  $r$  first; we have

$$\int r^2 \, dr = \frac{r^3}{3}.$$

Suppose  $a$  the radius of the sphere, then the limits of  $r$  are 0 and  $a$ ; thus

$$V = \iint \frac{a^3}{3} \sin\theta \, d\phi \, d\theta.$$

In thus integrating with respect to  $r$ , we collect all the elements like  $r^2 \sin \theta \Delta \phi \Delta \theta \Delta r$  which compose a pyramidal solid, having its vertex at the center of the sphere, and for its base the curvilinear element of spherical surface, which is denoted by  $a^2 \sin \theta \Delta \phi \Delta \theta$ .

Integrate next with respect to  $\theta$ ; we have

$$\int \sin \theta d\theta = -\cos \theta;$$

the limits of  $\theta$  are 0 and  $\frac{\pi}{2}$ ; thus

$$V = \int \frac{a^3}{3} d\phi.$$

In thus integrating with respect to  $\theta$ , we collect all the pyramids similar to  $\frac{a^3}{3} \sin \theta \Delta \phi \Delta \theta$  which form a wedge-shaped slice of the solid contained between the two planes through the fixed line corresponding to  $\phi$  and  $\phi + \Delta \phi$ .

Lastly, integrate with respect to  $\phi$  from 0 to  $\frac{\pi}{2}$ ; thus

$$V = \frac{\pi a^3}{6}.$$

In this example the integrations may be performed in any order, and the student should examine and illustrate them.

209. A right cone has its vertex on the surface of a sphere, and its axis coincident with the diameter of the sphere passing through that point; find the volume common to the cone and the sphere.

Let  $a$  be the radius of the sphere;  $\alpha$  the semi-vertical angle of the cone,  $V$  the required volume, then the polar equation to the sphere with the vertex of the cone as origin is  $r = 2a \cos \theta$ . Therefore

$$V = \int_0^{2\pi} \int_0^\alpha \int_0^{2a \cos \theta} r^2 \sin \theta d\phi d\theta dr.$$

210. The curve  $r = a(1 + \cos \theta)$  revolves round the initial line, find the volume of the solid generated.

Here the required volume

$$\begin{aligned} &= \int_0^\pi \int_0^{2\pi} \int_0^a (1 + \cos \theta) r^2 \sin \theta \, d\theta \, d\phi \, dr \\ &= \frac{2\pi a^3}{3} \int_0^\pi (1 + \cos \theta)^3 \sin \theta \, d\theta. \end{aligned}$$

It will be found that this  $= \frac{8\pi a^3}{3}$ .

EXAMPLES.

1. If the curve  $y^2(x - 4a) = ax(x - 3a)$  revolve round the axis of  $x$  the volume generated from  $x = 0$  to  $x = 3a$  is  $\frac{\pi a^3}{2} (15 - 16 \log 2)$ .
2. A cycloid revolves round the tangent at the vertex, shew that the volume generated is  $\pi^2 a^3$ .
3. A cycloid revolves round its base; shew that the volume generated is  $5\pi^2 a^3$ .
4. The curve  $y^2(2a - x) = x^3$  revolves round its asymptote; shew that the volume generated is  $2\pi^2 a^3$ .
5. The curve  $xy^2 = 4a^2(2a - x)$  revolves round its asymptote; shew that the volume generated is  $4\pi^2 a^3$ .
6. Find the volume of the closed portion of the solid generated by the revolution of the curve  $(y^2 - b^2)^2 = a^2 x$  round the axis of  $y$ .

*Result.*  $\frac{256}{315} \frac{\pi b^9}{a^5}$ .

7. Express the volume of a frustum of a sphere in terms of its height and the radii of its ends.

8. If the curve  $y^2 = 2mx + nx^2$  revolve about the axis of  $x$  find the volume of any frustum; and shew that it may be expressed either by

$$\frac{\pi h}{2} (b^2 + c^2 - \frac{1}{3} nh^2) \quad \text{or by } \pi h \left( x^2 + \frac{nh^2}{12} \right),$$

where  $h$  is the altitude of the frustum and  $b, c, r$  are the radii of its two ends and middle section. Deduce expressions for the volume of a cone and spheroid.

9. Find by integration the volume included between a right cone whose vertical angle is  $60^\circ$ , and a sphere of given radius touching it along a circle.

$$\text{Result. } \frac{\pi r^3}{6}.$$

10. If a paraboloid have its vertex in the base, and axis in the surface of a cylinder, the cylinder will be divided into parts which are as 3 : 5 by the surface of the paraboloid; the altitude and diameter of the base of the cylinder and the latus rectum of the paraboloid being all equal.

11. Determine the volume of the solid generated by the revolution of the curve  $(x^2 + y^2)^2 = a^2x^2 + b^2y^2$  about the axis of  $y$ , supposing  $a$  greater than  $b$ . Shew what the result becomes when  $a = b$ .

12. Find the volume of the solid formed by the revolution of the curve  $(y^2 + x^2)^2 = a^2(x^2 - y^2)$  round the axis of  $x$ .

$$\text{Result. } \frac{\pi a^3}{2} \left\{ \frac{1}{\sqrt{2}} \log(1 + \sqrt{2}) - \frac{1}{3} \right\}.$$

13. A paraboloid of revolution has its axis coincident with a diameter of a sphere, and its vertex outside the sphere; find the volume of the portion of the sphere outside the paraboloid.

*Result.*  $\frac{\pi h^3}{6}$  where  $h$  is the distance of the two planes in which the curves of intersection of the surfaces are situated.

14. Find the volume cut off from the surface

$$\frac{z^2}{c} + \frac{y^2}{b} = 2x$$

by a plane parallel to that of  $(y, z)$  at a distance  $a$  from it.

*Result.*  $\pi a^2 \sqrt{bc}$ .

15. A quadrant of an ellipse revolves round a tangent at the end of the minor axis of the ellipse; shew that the volume included by the surface formed by the curve is

$$\frac{\pi ab^3}{6} (10 - 3\pi).$$

16. Find the volume of the solid generated by the revolution of the closed part of the curve

$$x^3 - 3axy + y^3 = 0$$

round the line  $x + y = 0$ .

*Result.*  $\frac{8\pi^2 a^3}{3\sqrt{6}}$ .

17. If the axes of two equal circular cylinders of radius  $a$  intersect at an angle  $\beta$ , the volume common to both is  $\frac{16}{3} \frac{a^3}{\sin \beta}$ ; and the surface of each intercepted by the other is  $\frac{8a^2}{\sin \beta}$ .

18. Find the volume enclosed by the surfaces defined by the equations

$$x^2 + y^2 = cz, \quad x^2 + y^2 = ax, \quad z = 0,$$

illustrating by figures the progress of the summation.

*Result.*  $\frac{3\pi a^4}{32c}$ .

19. If  $S$  be a closed surface,  $dS$  an element of  $S$  about a point  $P$  at a distance  $r$  from a fixed point  $O$ , and  $\phi$  the angle which the normal at  $P$  drawn inwards

makes with the radius vector  $OP$ , shew that the volume contained by the surface

$$= \frac{1}{3} \int r \cos \phi \, dS,$$

the summation being extended over the whole surface.

Taking the centre of an ellipsoid as the point  $O$ , apply this formula to find its volume interpreting geometrically the steps of the integration.

20. The centre of a variable circle moves along the arc of a fixed circle; its plane is normal to the fixed circle, and its radius equal to the distance of its centre from a fixed diameter; find the volume generated, and if the solid so formed revolve round the fixed diameter, shew that the volume swept through is to the volume of the solid as 5 to 2.

21. Find the value of  $\iiint x^2 \, dx \, dy \, dz$  over the volume of an ellipsoid.

$$\text{Result. } \frac{4\pi a^3 bc}{15}.$$

22. The centre of a regular hexagon moves along a diameter of a given circle (radius =  $a$ ), the plane of the hexagon being perpendicular to this diameter and its magnitude varying in such a manner that one of its diagonals always coincides with a chord of the circle; shew that the volume of the solid generated is  $2\sqrt{3} a^3$ . Shew also that the surface of the solid is

$$a^3 (2\pi + 3\sqrt{3}).$$

23. Determine the limits of integration in order to obtain the volume contained between the plane of  $(x, y)$  and the surface whose equation is

$$Ax^2 + Bxy + Cy^2 - Dz - F = 0.$$

24. State the limits of the integration to be used in applying the formula  $\iiint dx dy dz$  to find the volume of a closed surface of the second order whose equation is

$$ax^2 + by^2 + cz^2 + a'yz + b'xz + c'xy = 1.$$

25. State between what limits the integrations in

$$\iiint dx dy dz$$

must be performed, in order to obtain the volume contained between the conical surface whose equation is

$$z = a - \sqrt{(x^2 + y^2)},$$

and the planes whose equations are  $x = z$  and  $x = 0$ ; and find the volume by this or by any other method.

*Result.*  $\frac{2a^3}{9}$ .

26. State between what limits the integrations must be taken in order to find the volume of the solid contained between the two surfaces  $cz = mx^2 + ny^2$  and  $z = ax + by$ ; and find the volume when

$$m = n = a = b = 1.$$

27. A cavity is just large enough to allow of the complete revolution of a circular disc of radius  $c$ , whose centre describes a circle of the same radius  $c$ , while the plane of the disc is constantly parallel to a fixed plane, and perpendicular to that of the circle in which its centre moves. Shew that the volume of the cavity is

$$\frac{2c^3}{3} (3\pi + 8).$$

28. The axis of a right cone coincides with the generating line of a cylinder; the diameter of both cone and cylinder is equal to the common altitude; find the



surface and volume of each part into which the cone is divided by the cylinder.

*Results.*

$$\text{Surfaces, } \frac{4\pi\sqrt{5}-3\sqrt{15}}{6} a^2 \text{ and } \frac{2\pi\sqrt{5}+3\sqrt{15}}{6} a^2;$$

$$\text{Volumes, } \frac{8\pi+27\sqrt{3}-64}{9} a^3 \text{ and } \frac{64-27\sqrt{3}-2\pi}{9} a^3;$$

where  $a$  is the radius of the base of the cone or cylinder.

29. Find the volume of the cono-cuneus determined by

$$z^2 + \frac{a^2 y^2}{x^2} = c^2,$$

which is contained between the planes  $x = 0$  and  $x = a$ .

$$\text{Result. } \frac{\pi c^2 a}{2}.$$

30. A conoid is generated by a straight line which passes through the axis of  $z$  and is perpendicular to it. Two sections are made by parallel planes, both planes being parallel to the axis of  $z$ . Shew that the volume of the conoid included between the planes is equal to the product of the distance of the planes into half the sum of the areas of the sections made by the planes.

## CHAPTER IX.

## DIFFERENTIATION OF AN INTEGRAL WITH RESPECT TO ANY QUANTITY WHICH IT MAY INVOLVE.

211. It is sometimes necessary to differentiate an integral with respect to some quantity which it involves; this question we shall now consider.

Required the differential coefficient of  $\int_a^b \phi(x) dx$  with respect to  $b$ , supposing  $\phi(x)$  not to contain  $b$ , and  $a$  to be independent of  $b$ .

Let 
$$u = \int_a^b \phi(x) dx;$$

suppose  $b$  changed into  $b + \Delta b$ , in consequence of which  $u$  becomes  $u + \Delta u$ ; thus

$$u + \Delta u = \int_a^{b+\Delta b} \phi(x) dx;$$

therefore 
$$\Delta u = \int_a^{b+\Delta b} \phi(x) dx - \int_a^b \phi(x) dx$$

$$= \int_b^{b+\Delta b} \phi(x) dx.$$

Now, by Art. 40,

$$\int_b^{b+\Delta b} \phi(x) dx = \Delta b \phi(b + \theta \Delta b),$$

where  $\theta$  is some proper fraction; thus

$$\frac{\Delta u}{\Delta b} = \phi(b + \theta \Delta b).$$

Let  $\Delta b$  and  $\Delta u$  diminish without limit; thus

$$\frac{du}{db} = \phi(b).$$

212. Similarly, if we differentiate  $u$  with respect to  $a$ , supposing  $\phi(x)$  not to contain  $a$ , and  $b$  to be independent of  $a$ , we obtain

$$\frac{du}{da} = -\phi(a).$$

213. Suppose  $\phi(x)$  to contain a quantity  $c$ , and let it be required to find the differential coefficient of  $\int_a^b \phi(x) dx$  with respect to  $c$ , supposing  $a$  and  $b$  independent of  $c$ .

Instead of  $\phi(x)$  it will be convenient to write  $\phi(x, c)$ , so that the presence of the quantity  $c$  may be more clearly indicated; denote the integral by  $u$ , thus

$$u = \int_a^b \phi(x, c) dx.$$

Suppose  $c$  changed into  $c + \Delta c$ , in consequence of which  $u$  becomes  $u + \Delta u$ ; thus

$$u + \Delta u = \int_a^b \phi(x, c + \Delta c) dx;$$

therefore 
$$\Delta u = \int_a^b \phi(x, c + \Delta c) dx - \int_a^b \phi(x, c) dx$$

$$= \int_a^b \{\phi(x, c + \Delta c) - \phi(x, c)\} dx;$$

thus 
$$\frac{\Delta u}{\Delta c} = \int_a^b \frac{\phi(x, c + \Delta c) - \phi(x, c)}{\Delta c} dx.$$

Now by the nature of a differential coefficient we have

$$\frac{\phi(x, c + \Delta c) - \phi(x, c)}{\Delta c} = \frac{d\phi(x, c)}{dc} + \rho,$$

where  $\rho$  is a quantity which diminishes without limit when  $\Delta c$  does so. Thus we have

$$\frac{\Delta u}{\Delta c} = \int_a^b \frac{d\phi(x, c)}{dc} dx + \int_a^b \rho dx.$$

When  $\Delta c$  is diminished indefinitely, the second integral vanishes; for it is not greater than  $(b-a)\rho'$  where  $\rho'$  is the greatest value  $\rho$  can have, and  $\rho'$  ultimately vanishes. Hence proceeding to the limit, we have

$$\frac{du}{dc} = \int_a^b \frac{d\phi(x, c)}{dc} dx.$$

214. It should be noticed that the preceding article supposes that neither  $a$  nor  $b$  is infinite; if, for example,  $b$  were infinite, we could not assert that  $(b-a)\rho'$  would necessarily vanish in the limit.

215. We have shewn then in Art. 213 that

$$\frac{d}{dc} \int_a^b \phi(x, c) dx = \int_a^b \frac{d\phi(x, c)}{dc} dx \dots\dots\dots(1).$$

We will point out a useful application of this equation. Suppose that  $\psi(x, c)$  is the function of which  $\phi(x, c)$  is the differential coefficient with respect to  $x$ , and that  $\chi(x, c)$  is the function of which  $\frac{d\phi(x, c)}{dc}$  is the differential coefficient with respect to  $x$ ; thus (1) may be written

$$\frac{d\psi(b, c)}{dc} - \frac{d\psi(a, c)}{dc} = \chi(b, c) - \chi(a, c) \dots\dots\dots(2),$$

let us suppose that  $b$  does not occur in  $\phi(x, c)$ , and that  $a$  is also independent of  $b$ ; then (2) may be written

$$\frac{d\psi(b, c)}{dc} + C = \chi(b, c) \dots\dots\dots(3),$$

where  $C$  denotes terms which are independent of  $b$ , that is, are constant with respect to  $b$ . Hence as  $b$  may have

any value we please in (3), we may replace  $b$  by  $x$ , and write

$$\chi(x, c) = \frac{d\psi(x, c)}{dc} + C \dots\dots\dots(4).$$

This equation may be applied to find  $\chi(x, c)$ ; as the constant may be introduced if required, we may dispense with writing it, and put (4) in the form

$$\int \frac{d\phi(x, c)}{dc} dx = \frac{d}{dc} \int \phi(x, c) dx.$$

For example, let  $\phi(x, c) = \frac{1}{1 + c^2 x^2}$ ; then

$$\int \phi(x, c) dx = \int \frac{dx}{1 + c^2 x^2} = \frac{1}{c} \tan^{-1} cx,$$

thus

$$\begin{aligned} \frac{d}{dc} \left( \frac{1}{c} \tan^{-1} cx \right) &= \int \frac{d}{dc} \left( \frac{1}{1 + c^2 x^2} \right) dx \\ &= - \int \frac{2cx^2}{(1 + c^2 x^2)^2} dx. \end{aligned}$$

Thus from knowing the value of  $\int \frac{dx}{1 + c^2 x^2}$  we are able to deduce by differentiation the value of the more complex integral  $\int \frac{2cx^2}{(1 + c^2 x^2)^2} dx$ .

216. Required the differential coefficient of  $\int_a^b \phi(x, c) dx$  with respect to  $c$  when both  $b$  and  $a$  are functions of  $c$ . Denote the integral by  $u$ ; then  $\frac{du}{dc}$  consists of three terms, one arising from the fact that  $\phi(x, c)$  contains  $c$ , one from the fact that  $b$  contains  $c$ , and one from the fact that  $a$  contains  $c$ .

Hence by the preceding articles,

$$\begin{aligned} \frac{du}{dc} &= \int_a^b \frac{d\phi(x, c)}{dc} dx + \frac{du}{db} \frac{db}{dc} + \frac{du}{da} \frac{da}{dc} \\ &= \int_a^b \frac{d\phi(x, c)}{dc} dx + \phi(b, c) \frac{db}{dc} - \phi(a, c) \frac{da}{dc}. \end{aligned}$$

217. With the suppositions of the preceding article we may proceed to find  $\frac{d^2u}{dc^2}$ . By differentiating with respect to  $c$  the term  $\int_a^b \frac{d\phi(x, c)}{dc} dx$  we obtain

$$\int_a^b \frac{d^2\phi(x, c)}{dc^2} dx + \frac{d\phi(b, c)}{dc} \frac{db}{dc} - \frac{d\phi(a, c)}{dc} \frac{da}{dc}.$$

From the other terms in  $\frac{du}{dc}$  we obtain by differentiation


$$\begin{aligned} &\phi(b, c) \frac{d^2b}{dc^2} + \frac{d\phi(b, c)}{db} \left(\frac{db}{dc}\right)^2 + \frac{d\phi(b, c)}{dc} \frac{db}{dc} \\ &- \phi(a, c) \frac{d^2a}{dc^2} - \frac{d\phi(a, c)}{da} \left(\frac{da}{dc}\right)^2 - \frac{d\phi(a, c)}{dc} \frac{da}{dc}. \end{aligned}$$

Thus  $\frac{d^2u}{dc^2} = \int_a^b \frac{d^2\phi(x, c)}{dc^2} dx$

$$\begin{aligned} &+ \phi(b, c) \frac{d^2b}{dc^2} + \frac{d\phi(b, c)}{db} \left(\frac{db}{dc}\right)^2 + 2 \frac{d\phi(b, c)}{dc} \frac{db}{dc} \\ &- \phi(a, c) \frac{d^2a}{dc^2} - \frac{d\phi(a, c)}{da} \left(\frac{da}{dc}\right)^2 - 2 \frac{d\phi(a, c)}{dc} \frac{da}{dc}. \end{aligned}$$

Similarly  $\frac{d^3u}{dc^3}$  may be found and higher differential coefficients of  $u$  if required.

218. The following geometrical illustration may be given of Art. 216.



Let  $y = \phi(x, c)$  be the equation to the curve  $APQ$ , and  $y = \phi(x, c + \Delta c)$  the equation to the curve  $A'P'Q'$ .

$$\begin{aligned} \text{Let} \quad OM &= a, & ON &= b, \\ MM' &= \Delta a, & NN' &= \Delta b. \end{aligned}$$

Then  $u$  denotes the area  $PMNQ$ , and  $u + \Delta u$  denotes the area  $P'M'N'Q'$ . Hence

$$\Delta u = P'p q Q' + QNN'q - PMM'p,$$

$$\text{and} \quad \frac{\Delta u}{\Delta c} = \frac{P'p q Q'}{\Delta c} + \frac{QNN'q}{\Delta c} - \frac{PMM'p}{\Delta c}.$$

It may easily be seen that the limit of the first term is the limit of  $\int_a^b \frac{\phi(x, c + \Delta c) - \phi(x, c)}{\Delta c} dx$ ; that the limit of the

second term is the limit of  $\phi(b, c) \frac{\Delta b}{\Delta c}$ , and that the limit of the third term is the limit of  $\phi(a, c) \frac{\Delta a}{\Delta c}$ . This gives the result of Art. 216.

219. *Example.* Find a curve such that the area between the curve, the axis of  $x$ , and any ordinate, shall bear a constant ratio to the rectangle contained by that ordinate and the corresponding abscissa.

Suppose  $\phi(x)$  the ordinate of the curve to the abscissa  $x$ ; then  $\int_0^c \phi(x) dx$  expresses the area between the curve, the axis of  $x$ , and the ordinate  $\phi(c)$ : hence by supposition we must have

$$\int_0^c \phi(x) dx = \frac{c\phi(c)}{n},$$

where  $n$  is some constant. This is to hold for all values of  $c$ ; hence we may differentiate with respect to  $c$ ; thus

$$\phi(c) = \frac{\phi(c)}{n} + \frac{c\phi'(c)}{n};$$

therefore  $c\phi'(c) = (n-1)\phi(c)$ ,

and  $\frac{\phi'(c)}{\phi(c)} = \frac{n-1}{c}$ .

By integration  $\log \phi(c) = (n-1) \log c + \text{constant}$ ;

thus  $\phi(c) = Ac^{n-1}$ ,

and  $\phi(x) = Ax^{n-1}$ ,

which determines the required curve.

220. Find the form of  $\phi(x)$ , so that for all values of  $c$

$$\frac{\int_0^c x \{\phi(x)\}^2 dx}{\int_0^c \{\phi(x)\}^2 dx} = \frac{c}{n}.$$

By the supposition

$$\int_0^c x \{\phi(x)\}^2 dx = \frac{c}{n} \int_0^c \{\phi(x)\}^2 dx.$$



Differentiate with respect to  $c$ ; thus

$$c \{\phi(c)\}^2 = \frac{1}{n} \int_0^c \{\phi(x)\}^2 dx + \frac{c}{n} \{\phi(c)\}^2;$$

thus 
$$c \left(1 - \frac{1}{n}\right) \{\phi(c)\}^2 = \frac{1}{n} \int_0^c \{\phi(x)\}^2 dx.$$

Differentiate again with respect to  $c$ ;

thus 
$$\left(1 - \frac{1}{n}\right) \{\phi(c)\}^2 + 2c \left(1 - \frac{1}{n}\right) \phi(c) \phi'(c) = \frac{\{\phi(c)\}^2}{n};$$

hence 
$$\left(1 - \frac{2}{n}\right) \phi(c) + 2c \left(1 - \frac{1}{n}\right) \phi'(c) = 0;$$

therefore 
$$\frac{\phi'(c)}{\phi(c)} = \frac{2-n}{2(n-1)} \frac{1}{c}.$$

Integrate; thus

$$\log \phi(c) = \frac{2-n}{2(n-1)} \log c + \text{constant};$$

therefore 
$$\phi(c) = Ac^{\frac{2-n}{2(n-1)}},$$

where  $A$  is some constant; thus we have finally

$$\phi(x) = Ax^{\frac{2-n}{2(n-1)}}.$$

This is the solution of a problem in Analytical Statics, which may be enunciated thus: The distance of the centre of gravity of a segment of a solid of revolution from the vertex is always  $\frac{1}{n}$  th part of the height of the segment; find the generating curve. The required equation is  $y = \phi(x)$ .

221. Find the form of  $\phi(x)$  so that the integral  $\int_0^c \frac{\phi(x) dx}{\sqrt{c-x}}$  may be independent of  $c$ .

Denote the integral by  $u$ , and suppose  $x = cz$ ; thus

$$u = \int_0^c \frac{\phi(x) dx}{\sqrt{c-x}} = \int_0^1 \frac{\sqrt{c} \phi(cz) dz}{\sqrt{1-z}}.$$

Since  $u$  is to be independent of  $c$ , the differential coefficient of  $u$  with respect to  $c$  must vanish. Now

$$\begin{aligned}\frac{du}{dc} &= \int_0^1 \frac{\frac{\phi(cz)}{2\sqrt{c}} + z\sqrt{c}\phi'(cz)}{\sqrt{1-z}} dz \\ &= \int_0^c \frac{\phi(x) + 2x\phi'(x)}{2c\sqrt{c-x}} dx.\end{aligned}$$

This last integral then must vanish whatever  $c$  may be; hence we must have

$$\phi(x) + 2x\phi'(x) = 0;$$

therefore 
$$\frac{\phi'(x)}{\phi(x)} = -\frac{1}{2x};$$

therefore 
$$\log \phi(x) = -\frac{1}{2} \log x + \text{constant},$$

therefore 
$$\phi(x) = \frac{A}{\sqrt{x}}.$$

This is the solution of a problem in Dynamics, which may be thus enunciated. Find a curve, such that the time of falling down an arc of the curve from *any* point to the lowest point may be the same. If  $s$  denote the arc of the curve measured from the lowest point, then

$$\frac{ds}{dx} = \phi(x) \quad \text{and} \quad s = 2A\sqrt{x};$$

so that the curve is a cycloid.

## CHAPTER X.

## ELLIPTIC INTEGRALS.

222. THE integrals  $\int \frac{d\theta}{\sqrt{(1-c^2 \sin^2 \theta)}}$ ,  $\int \sqrt{(1-c^2 \sin^2 \theta)} d\theta$ , and  $\int \frac{d\theta}{(1+a \sin^2 \theta) \sqrt{(1-c^2 \sin^2 \theta)}}$ , are called *elliptic functions* or *elliptic integrals* of the first, second, and third order respectively; the first is denoted by  $F(c, \theta)$ , the second by  $E(c, \theta)$ , and the third by  $\Pi(c, a, \theta)$ . The integrals are all supposed to be taken between the limits 0 and  $\theta$ , so that they vanish with  $\theta$ ;  $\theta$  is called the *amplitude* of the function. The constant  $c$  is supposed less than unity; it is called the *modulus* of the function. The constant  $a$ , which occurs in the function of the third order is called the *parameter*. When the integrals are taken between the limits 0 and  $\frac{\pi}{2}$ , they are called *complete functions*; that is, the *amplitude* of a complete function is  $\frac{\pi}{2}$ .

223. The second elliptic integral expresses the length of a portion of the arc of an ellipse measured from the end of the minor axis, the excentricity of the ellipse being the *modulus* of the function. From this circumstance, and from the fact that the three integrals are connected by remarkable properties, the name *elliptic integrals* has been derived.

224. The subject of *elliptic integrals* is very extensive; we shall merely give a few of the simpler results, and refer the student for fuller investigations to Hymers's *Integral Calculus*, or to the writings of Legendre, Jacobi and Abel.

225. If  $\theta$  and  $\phi$  are connected by the equation

$$F(c, \theta) + F(c, \phi) = F(c, \mu),$$

where  $\mu$  is a constant; then will

$$\cos \theta \cos \phi - \sin \theta \sin \phi \sqrt{(1 - c^2 \sin^2 \mu)} = \cos \mu.$$

Consider  $\theta$  and  $\phi$  as functions of a new variable  $t$ , and differentiate the given equation; thus

$$\frac{1}{\sqrt{(1 - c^2 \sin^2 \theta)}} \frac{d\theta}{dt} + \frac{1}{\sqrt{(1 - c^2 \sin^2 \phi)}} \frac{d\phi}{dt} = 0 \dots\dots(1).$$

Now as  $t$  is a new arbitrary variable, we are at liberty to assume

$$\frac{d\theta}{dt} = \sqrt{(1 - c^2 \sin^2 \theta)},$$

thus from the equation (1)

$$\frac{d\phi}{dt} = -\sqrt{(1 - c^2 \sin^2 \phi)}.$$

Square these two equations and differentiate; thus

$$\frac{d^2\theta}{dt^2} = -c^2 \sin \theta \cos \theta, \quad \frac{d^2\phi}{dt^2} = -c^2 \sin \phi \cos \phi;$$

therefore, 
$$\frac{d^2(\theta \pm \phi)}{dt^2} = -\frac{c^2}{2} (\sin 2\theta \pm \sin 2\phi).$$

Let  $\theta + \phi = \psi$  and  $\theta - \phi = \chi$ ; thus

$$\frac{d^2\psi}{dt^2} = -c^2 \sin \psi \cos \chi, \quad \frac{d^2\chi}{dt^2} = -c^2 \sin \chi \cos \psi.$$

Also 
$$\frac{d\psi}{dt} \frac{d\chi}{dt} = \left(\frac{d\theta}{dt}\right)^2 - \left(\frac{d\phi}{dt}\right)^2 = -c^2 \sin \psi \sin \chi;$$

therefore 
$$\frac{\frac{d^2\psi}{dt^2}}{\frac{d\psi}{dt} \frac{d\chi}{dt}} = \cot \chi, \quad \frac{\frac{d^2\chi}{dt^2}}{\frac{d\psi}{dt} \frac{d\chi}{dt}} = \cot \psi;$$

therefore

$$\frac{d}{dt} \left( \log \frac{d\psi}{dt} \right) = \frac{d}{dt} \log \sin \chi, \quad \frac{d}{dt} \left( \log \frac{d\chi}{dt} \right) = \frac{d}{dt} \log \sin \psi;$$

therefore 
$$\log \frac{d\psi}{dt} = \log \sin \chi + \text{constant},$$

therefore 
$$\left. \begin{aligned} \frac{d\psi}{dt} &= A \sin \chi \\ \text{and similarly,} \quad \frac{d\chi}{dt} &= B \sin \psi \end{aligned} \right\} \dots\dots\dots (2),$$

where  $A$  and  $B$  are constants.

Hence 
$$A \sin \chi \frac{d\chi}{dt} = B \sin \psi \frac{d\psi}{dt},$$

therefore 
$$A \cos \chi = B \cos \psi + C \dots\dots\dots (3).$$

Now from the original given equation we see that if  $\phi = 0$

$$F(c, \theta) = F(c, \mu);$$

therefore then  $\theta = \mu$  and  $\chi = \psi = \mu$ ;

thus from (3) 
$$(A - B) \cos \mu = C;$$

thus 
$$A \cos (\theta - \phi) = B \cos (\theta + \phi) + (A - B) \cos \mu;$$

therefore

$$(A - B) \cos \theta \cos \phi + (A + B) \sin \theta \sin \phi = (A - B) \cos \mu \dots (4).$$

In (2) put for  $\frac{d\psi}{dt}$  its value

$$\sqrt{(1 - c^2 \sin^2 \theta)} - \sqrt{(1 - c^2 \sin^2 \phi)},$$

and for  $\frac{d\chi}{dt}$  its value

$$\sqrt{(1 - c^2 \sin^2 \theta) + \sqrt{(1 - c^2 \sin^2 \phi)}},$$

and then suppose  $\phi = 0$ ; thus

$$\sqrt{(1 - c^2 \sin^2 \mu) - 1} = A \sin \mu,$$

and

$$\sqrt{(1 - c^2 \sin^2 \mu) + 1} = B \sin \mu.$$

Substitute for  $A - B$  and  $A + B$  in (4);

thus  $\cos \theta \cos \phi - \sqrt{(1 - c^2 \sin^2 \mu) \sin \theta \sin \phi} = \cos \mu.$

226. The relation just found may be put in a different form. Clear the equation of radicals; thus

$$(\cos \theta \cos \phi - \cos \mu)^2 = (1 - c^2 \sin^2 \mu) \sin^2 \theta \sin^2 \phi;$$

therefore

$$\begin{aligned} \cos^2 \theta + \cos^2 \phi + \cos^2 \mu - 2 \cos \theta \cos \phi \cos \mu \\ = 1 - c^2 \sin^2 \mu \sin^2 \theta \sin^2 \phi. \end{aligned}$$

Add  $\cos^2 \phi \cos^2 \mu$  to both sides and transpose; thus

$$\begin{aligned} (\cos \theta - \cos \phi \cos \mu)^2 \\ = 1 - \cos^2 \phi - \cos^2 \mu + \cos^2 \phi \cos^2 \mu - c^2 \sin^2 \mu \sin^2 \theta \sin^2 \phi \\ = \sin^2 \phi \sin^2 \mu (1 - c^2 \sin^2 \theta); \end{aligned}$$

therefore  $\cos \theta = \cos \phi \cos \mu + \sin \phi \sin \mu \sqrt{(1 - c^2 \sin^2 \theta)}.$

The positive sign of the radical is taken, because when  $\theta = 0$ , we must have  $\phi = \mu.$

227. We shall now shew how an elliptic function of the first order may be connected with another having a different modulus.

Let  $F(c, \theta)$  denote the function; assume

$$\tan \theta = \frac{\sin 2\phi}{c + \cos 2\phi};$$

therefore 
$$\frac{1}{\cos^2 \theta} \frac{d\theta}{d\phi} = \frac{2(1+c \cos 2\phi)}{(c + \cos 2\phi)^2},$$

therefore 
$$\frac{d\theta}{d\phi} = \frac{2(1+c \cos 2\phi)}{1+2c \cos 2\phi + c^2}.$$

And 
$$1 - c^2 \sin^2 \theta = 1 - \frac{c^2 \sin^2 2\phi}{1+2c \cos 2\phi + c^2}$$

$$= \frac{1+2c \cos 2\phi + c^2 \cos^2 2\phi}{1+2c \cos 2\phi + c^2};$$

therefore

$$\int \frac{d\theta}{\sqrt{1-c^2 \sin^2 \theta}} = \int \frac{2(1+c \cos 2\phi)}{1+2c \cos 2\phi + c^2} \cdot \frac{\sqrt{(1+2c \cos 2\phi + c^2)}}{1+c \cos 2\phi} d\phi$$

$$= 2 \int \frac{d\phi}{\sqrt{(1+2c \cos 2\phi + c^2)}} = \frac{2}{1+c} \int \sqrt{\left\{1 - \frac{4c}{(1+c)^2} \sin^2 \phi\right\}}.$$

No constant is added because  $\phi$  vanishes with  $\theta$ . Thus

$$F(c, \theta) = \frac{2}{1+c} F(c_1, \phi), \text{ where}$$

$$c_1^2 = \frac{4c}{(1+c)^2} \text{ and } \tan \theta = \frac{\sin 2\phi}{c + \cos 2\phi}.$$

The last relation may be written thus

$$c \sin \theta = \sin (2\phi - \theta).$$

We may notice that  $c_1$  is greater than  $c$  for

$$\frac{c_1^2}{c^2} = \frac{4}{c(1+c)^2},$$

and since  $c$  is less than unity, 4 is greater than  $c(1+c)^2$ .

If  $\phi = \frac{\pi}{2}$ , then  $\theta = \pi$ ; thus

$$\frac{2}{1+c} F\left(c_1, \frac{\pi}{2}\right) = F(c, \pi) = 2 F\left(c, \frac{\pi}{2}\right).$$

228. We will give one more proposition in this subject, by establishing a relation among Elliptic Functions of the second order, analogous to that proved in Art. 225 for functions of the first order.

If  $\cos \theta \cos \phi - \sin \theta \sin \phi \sqrt{1 - c^2 \sin^2 \mu} = \cos \mu$ ,  
then will

$$E(c, \theta) + E(c, \phi) - E(c, \mu) = c^2 \sin \theta \sin \phi \sin \mu.$$

By virtue of the given equation connecting the amplitudes,  $\phi$  is a function of  $\theta$ ; thus we may assume

$$E(c, \theta) + E(c, \phi) - E(c, \mu) = f(\theta).$$

Differentiate; thus

$$\begin{aligned} f'(\theta) &= \sqrt{1 - c^2 \sin^2 \theta} + \sqrt{1 - c^2 \sin^2 \phi} \frac{d\phi}{d\theta} \\ &= \frac{\cos \theta - \cos \phi \cos \mu}{\sin \phi \sin \mu} + \frac{\cos \phi - \cos \theta \cos \mu}{\sin \theta \sin \mu} \frac{d\phi}{d\theta} \\ &\hspace{15em} \text{(by Art. 226),} \\ &= \frac{d\{\sin^2 \theta + \sin^2 \phi + 2 \cos \theta \cos \phi \cos \mu\}}{d\theta} \times \frac{1}{2 \sin \theta \sin \phi \sin \mu}. \end{aligned}$$

But  $\sin^2 \theta + \sin^2 \phi + 2 \cos \theta \cos \phi \cos \mu$

$$= 1 + \cos^2 \mu + c^2 \sin^2 \theta \sin^2 \phi \sin^2 \mu;$$

thus  $f'(\theta) = c^2 \frac{d(\sin \theta \sin \phi \sin \mu)}{d\theta}$ .

Therefore, by integration

$$f(\theta) = c^2 \sin \theta \sin \phi \sin \mu.$$

No constant is added, because  $f(\theta)$  obviously vanishes with  $\theta$ .




## CHAPTER XI.

## CHANGE OF THE VARIABLES IN A MULTIPLE INTEGRAL.

229. WE have seen in Art. 62 that the double integral  $\int_a^b \int_a^b \phi(x, y) dx dy$  is equal to  $\int_a^b \int_a^b \phi(x, y) dy dx$  when the *limits are constant*, that is, a change in the *order* of integration produces no change in the limits for the two integrations. But when the limits of the first integration are functions of the other variable, this statement no longer holds, as we have seen in several examples in the seventh and eighth chapters. We add here a few additional examples.

230. Change the order of integration in

$$\int_0^a \int_0^{\sqrt{a^2-x^2}} \phi(x, y) dx dy.$$



The limits of the integration with respect to  $y$  here are  $y=0$  and  $y=\sqrt{a^2-x^2}$ ; that is, we may consider the


integral extending from the axis of  $x$  to the boundary of a circle, having its centre at the origin, and radius equal to  $a$ . Then the integration with respect to  $x$  extends from the axis of  $y$  to the extreme point  $A$  of the quadrant. Thus if we consider  $z = \phi(x, y)$  as the equation to a surface, the above double integral represents the volume of that solid which is contained between the surface, the plane of  $(x, y)$ , and a line moving perpendicularly to this plane round the boundary  $OAPBO$ .

It is then obvious from the figure that if the integration with respect to  $x$  is performed first, the limits will be  $x=0$  and  $x=\sqrt{a^2-y^2}$ , and then the limits for  $y$  will be  $y=0$  and  $y=a$ . Thus the transformed integral is

$$\int_0^a \int_0^{\sqrt{a^2-y^2}} \phi(x, y) dy dx.$$

231. Change the order of integration in

$$\int_0^{\frac{\pi}{2}} \int_0^{2a \cos \theta} \phi(r, \theta) r d\theta dr.$$




Let  $OA = 2a$ , and describe a semicircle on  $OA$  as diameter. Let  $POX = \theta$ , then  $OP = 2a \cos \theta$ . Thus the double integral may be considered as the limit of a summation of values of  $\phi(r, \theta) r \Delta\theta \Delta r$  over all the area of the semicircle. Hence when the order is changed we must integrate for  $\theta$  from 0 to  $\cos^{-1} \frac{r}{2a}$ , and for  $r$  from 0 to  $2a$ .

Thus the transformed integral is

$$\int_0^{2a} \int_0^{\cos^{-1} \frac{r}{2a}} \phi(r, \theta) r dr d\theta.$$

232. Change the order of integration in

$$\int_0^{2a} \int_{\frac{x^2}{4a}}^{3a-x} \phi(x, y) dx dy.$$



The integration for  $y$  is taken from  $y = \frac{x^2}{4a}$  to  $y = 3a - x$ .

The equation  $y = \frac{x^2}{4a}$  belongs to a parabola  $OLD$ , and  $y = 3a - x$  to a straight line  $BLC$ , which passes through  $L$ , the extremity of the latus rectum of the parabola.

Thus the integration may be considered as extending over the area  $OLBSO$ . Now let the order of integration be changed; we shall have to consider separately the spaces  $OLS$  and  $BLS$ . For the space  $OLS$  we must integrate from  $x=0$  to  $x=2\sqrt{(ay)}$ , and then from  $y=0$  to  $y=a$ ; and for the space  $BLS$  we must integrate from  $x=0$  to  $x=3a-y$ , and then from  $y=a$  to  $y=3a$ . Thus the transformed integral is

$$\int_0^a \int_0^{2\sqrt{(ay)}} \phi(x, y) dy dx + \int_a^{3a} \int_0^{3a-y} \phi(x, y) dy dx.$$

233. Change the order of integration in

$$\int_0^1 \int_x^{x(2-x)} \phi(x, y) dx dy.$$

Here the integration with respect to  $y$  is taken from  $y = x$  to  $y = x(2 - x)$ . The equation  $y = x$  represents a straight line, and  $y = x(2 - x)$  represents a parabola. The reader will find on examining a figure, that the transformed integral is

$$\int_0^1 \int_{1-\sqrt{1-y}}^y \phi(x, y) dy dx.$$

234. Change the order of integration in

$$\int_0^a \int_{\sqrt{a^2-x^2}}^{x+2a} \phi(x, y) dx dy.$$


Here the integration with respect to  $y$  is taken from  $y = \sqrt{a^2 - x^2}$  to  $y = x + 2a$ . The equation  $y = \sqrt{a^2 - x^2}$  represents a circle, and  $y = x + 2a$  represents a straight line. The reader will find on examining a figure, that when the integration with respect to  $x$  is performed first, the integral must be separated into three portions; the transformed integral is

$$\begin{aligned} \int_0^a \int_{\sqrt{a^2-y^2}}^a \phi(x, y) dy dx + \int_a^{2a} \int_0^a \phi(x, y) dy dx \\ + \int_{2a}^{3a} \int_{y-2a}^a \phi(x, y) dy dx. \end{aligned}$$

235. Change the order of integration in

$$\int_0^a \int_0^{\frac{b}{b+x}} \phi(x, y) dx dy.$$

Here the integration with respect to  $y$  is taken from  $y = 0$  to  $y = \frac{b}{b+x}$ . The equation  $y = \frac{b}{b+x}$  represents an hyperbola; let  $BDE$  be this hyperbola, and let  $OA = a$ . Then the integration may be considered as extending over the



space  $OBDA$ . Let the order of the integration be changed; we shall then have to consider separately the spaces  $OADC$  and  $CDB$ . For the space  $OADC$  we must integrate from  $x=0$  to  $x=a$ , and then from  $y=0$  to  $y=\frac{b}{b+a}$ . For the space  $CDB$  we must integrate from  $x=0$  to  $x=\frac{b(1-y)}{y}$ , and then from  $y=\frac{b}{b+a}$  to  $y=1$ . Thus the transformed integral is

$$\int_0^{\frac{b}{b+a}} \int_0^a \phi(x, y) dy dx + \int_{\frac{b}{b+a}}^1 \int_0^{\frac{b(1-y)}{y}} \phi(x, y) dy dx.$$

236. Change the order of integration in

$$\int_0^h \int_{\lambda x}^{c-\mu x} \phi(x, y) dx dy,$$

where  $h = \frac{c}{\lambda + \mu}$ . The transformed integral is

$$\int_0^{\lambda h} \int_{\frac{y}{\lambda}}^{\frac{c-y}{\mu}} \phi(x, y) dy dx + \int_{\lambda h}^c \int_0^{\frac{c-y}{\mu}} \phi(x, y) dy dx.$$

237. Change the order of integration in

$$\int_0^a \int_0^x \int_0^y \phi(x, y, z) dx dy dz.$$

The integration here may be considered to be extended throughout a pyramid, the bounding planes of which are given by the equations

$$z = 0, \quad z = y, \quad y = x, \quad x = a.$$

The integral may be transformed in different ways, and thus we obtain

$$\int_0^a \int_y^a \int_0^y \phi(x, y, z) dy dx dz,$$

or 
$$\int_0^a \int_0^y \int_y^a \phi(x, y, z) dy dz dx,$$

or 
$$\int_0^a \int_x^a \int_y^a \phi(x, y, z) dz dy dx,$$

or 
$$\int_0^a \int_0^x \int_x^a \phi(x, y, z) dx dz dy,$$

or 
$$\int_0^a \int_x^a \int_x^z \phi(x, y, z) dz dx dy.$$

These transformations may be verified by putting for  $\phi(x, y, z)$  some simple function, so that the integrals can be actually obtained; for example, if we replace  $\phi(x, y, z)$  by unity, we find  $\frac{a^3}{6}$  as the value of any one of the six forms.

238. These examples will sufficiently illustrate the subject; it is impossible to lay down any simple rules for the discovery of the limits of the transformed integral. It is not absolutely necessary to draw figures as we have done, for the figures convey no information which could not be obtained by reflection on the different values which the variables must have, in order to make the integration extend over the range indicated by the given limits. But the figures materially assist in arriving speedily and correctly at the result.

We now proceed to the problem which is the main object of the present chapter, namely, the change of the variables in a *multiple* integral. We begin with the case of a *double* integral.

239. The problem to be solved is the following. Required to transform the double integral  $\iint V dx dy$ , where  $V$  is a function of  $x$  and  $y$ , to another double integral in which the variables are  $u$  and  $v$ , the old and new variables being connected by the equations

$$\phi_1(x, y, u, v) = 0 \dots \dots \quad \phi_2(x, y, u, v) = 0 \dots \dots (1).$$

We suppose that the original integral is to be taken between known limits of  $y$  and  $x$ ; as we integrate with respect to  $y$  first, the limits of  $y$  may be functions of  $x$ . Of course while integrating with respect to  $y$  we regard  $x$  as constant.

We first transform the integral with respect to  $y$  into an integral with respect to  $v$ . This is theoretically very simple; from equations (1) eliminate  $u$  and obtain  $y$  as a function of  $x$  and  $v$ , say

$$y = \psi(x, v) \dots \dots \dots (2),$$

from which we get

$$dy = \psi'(x, v) dv,$$

where  $\psi'(x, v)$  means the differential coefficient of  $\psi(x, v)$  with respect to  $v$ .

Substitute then for  $y$  and  $dy$  in  $\int V dy$ , and we obtain  $\int V_1 \psi'(x, v) dv$ , where  $V_1$  is what  $V$  becomes when we put for  $y$  its value in  $V$ . Hence the original double integral becomes

$$\iint V_1 \psi'(x, v) dx dv.$$

Thus we have removed  $y$  and taken  $v$  instead. As the limiting values of  $y$  between which we had originally to

integrate are known, we shall from (2) know the limiting values of  $v$ , between which we ought to integrate. It will be observed, that in finding  $\frac{dy}{dv}$  from (2), we supposed  $x$  constant; this we do because, as already remarked, when we integrate the proposed expression with respect to  $y$  we must consider  $x$  constant.

The next step is to change the *order* of the above integration with respect to  $x$  and  $v$ , that is, to perform the integration with respect to  $x$  *first*. This is a subject which we have already examined; all we have to do is to determine the *new limits* properly. Thus supposing this point settled, we have changed the original expression into

$$\iint V_1 \psi'(x, v) dv dx.$$

It remains to remove  $x$  from this expression and replace it by  $u$ . We proceed precisely as before. From equations (1) eliminate  $y$ , and obtain  $x$  as a function of  $v$  and  $u$ , say

$$x = \chi(v, u) \dots \dots \dots (3),$$

from which we get

$$dx = \chi'(v, u) du,$$

where  $\chi'(v, u)$  means the differential coefficient of  $\chi(v, u)$  with respect to  $u$ .

Substitute then for  $x$  and  $dx$ , and the double integral becomes

$$\iint V' \psi'(x, v) \chi'(v, u) dv du,$$

where  $V'$  is what  $V_1$  becomes when we put for  $x$  its value in  $V_1$ . Thus the double integral now contains only  $u$  and  $v$ , since for the  $x$  which occurs in  $\psi'(x, v)$  we suppose its value substituted, namely,  $\chi(v, u)$ . Moreover since the limits between which the integration with respect to  $x$  was to be taken have been already settled, we know the limits between which the integration with respect to  $u$  must be taken.



We have thus given the complete *theoretical* solution of the problem; it only remains to add a *practical* method for determining  $\psi'(x, v)$  and  $\chi'(v, u)$ ; to this we proceed.

We observe that  $\psi'(x, v)$  or  $\frac{dy}{dv}$  is to be found from equations (1) by eliminating  $u$ , considering  $x$  constant; the following is exactly equivalent; from (1)

$$\frac{d\phi_1}{dy} \frac{dy}{dv} + \frac{d\phi_1}{du} \frac{du}{dv} + \frac{d\phi_1}{dv} = 0,$$

$$\frac{d\phi_2}{dy} \frac{dy}{dv} + \frac{d\phi_2}{du} \frac{du}{dv} + \frac{d\phi_2}{dv} = 0.$$

Eliminate  $\frac{du}{dv}$ ; thus

$$\frac{\frac{d\phi_1}{dy} \frac{dy}{dv} + \frac{d\phi_1}{dv}}{\frac{d\phi_1}{du}} = \frac{\frac{d\phi_2}{dy} \frac{dy}{dv} + \frac{d\phi_2}{dv}}{\frac{d\phi_2}{du}},$$

therefore

$$\frac{dy}{dv} = \frac{\frac{d\phi_1}{dv} \frac{d\phi_2}{du} - \frac{d\phi_1}{du} \frac{d\phi_2}{dv}}{\frac{d\phi_1}{du} \frac{d\phi_2}{dy} - \frac{d\phi_1}{dy} \frac{d\phi_2}{du}}.$$

This then is an equivalent for  $\psi'(x, v)$ .

Again,  $\chi'(v, u)$  or  $\frac{dx}{du}$  is to be found from equations (1) by eliminating  $y$ , regarding  $v$  as constant; the following is exactly equivalent; from (1)

$$\frac{d\phi_1}{dx} \frac{dx}{du} + \frac{d\phi_1}{dy} \frac{dy}{du} + \frac{d\phi_1}{du} = 0,$$

$$\frac{d\phi_2}{dx} \frac{dx}{du} + \frac{d\phi_2}{dy} \frac{dy}{du} + \frac{d\phi_2}{du} = 0.$$

From these equations by eliminating  $\frac{dy}{du}$  we find

$$\frac{dx}{du} = \frac{\frac{d\phi_1}{du} \frac{d\phi_2}{dy} - \frac{d\phi_1}{dy} \frac{d\phi_2}{du}}{\frac{d\phi_1}{dy} \frac{d\phi_2}{dx} - \frac{d\phi_1}{dx} \frac{d\phi_2}{dy}}.$$

This then is an equivalent for  $\chi'(v, u)$ .

Thus 
$$\psi'(x, v) \chi'(v, u) = \frac{\frac{d\phi_1}{dv} \frac{d\phi_2}{du} - \frac{d\phi_1}{du} \frac{d\phi_2}{dv}}{\frac{d\phi_1}{dy} \frac{d\phi_2}{dx} - \frac{d\phi_1}{dx} \frac{d\phi_2}{dy}}.$$

Hence the conclusion is that

$$\iint V dx dy = \iint V \frac{\frac{d\phi_1}{dv} \frac{d\phi_2}{du} - \frac{d\phi_1}{du} \frac{d\phi_2}{dv}}{\frac{d\phi_1}{dy} \frac{d\phi_2}{dx} - \frac{d\phi_1}{dx} \frac{d\phi_2}{dy}} dv du \dots \dots \dots (4),$$

where after the differentiations have been performed, we must put for  $x$  and  $y$  their values in terms of  $u$  and  $v$  to be found from (1); also the values of  $x$  and  $y$  must be substituted in  $V$ .

An important particular case is that in which  $x$  and  $y$  are given *explicitly* as functions of  $u$  and  $v$ ; the equations (1) then take the form

$$x - f_1(u, v) = 0, \quad y - f_2(u, v) = 0 \dots \dots \dots (5).$$

Here 
$$\frac{d\phi_1}{dx} = 1, \quad \frac{d\phi_1}{dy} = 0, \quad \frac{d\phi_2}{dx} = 0, \quad \frac{d\phi_2}{dy} = 1,$$

and the transformed integral becomes

$$\iint V \left( \frac{df_1}{du} \frac{df_2}{dv} - \frac{df_1}{dv} \frac{df_2}{du} \right) dv du,$$

where we must substitute for  $x$  and  $y$  their values from (5) in  $V$ .

Thus we may write

$$\iint V dx dy = \iint V \left( \frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du} \right) dv du \dots\dots (6).$$

The formulæ in (4) and (6) are those which are usually given; they contain a simple solution of the proposed problem in those cases where the limits of the new integrations are obvious. But in some examples the difficulty of determining the limits of the new integrations would be very great, and to ensure a correct result it would be necessary instead of using these formulæ, to carry on the process precisely in the manner indicated in the theory, by removing one of the old variables at a time.

240. The following is an example.

Required to transform  $\int_0^a \int_0^b V dx dy$ , having given

$$y + x = u, \quad y = uv.$$

From the given equations we have

$$x = u(1 - v), \quad y = uv;$$

thus  $\frac{dx}{du} = 1 - v, \quad \frac{dx}{dv} = -u, \quad \frac{dy}{du} = v, \quad \frac{dy}{dv} = u;$

therefore  $\frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du} = u(1 - v) + uv = u.$

Hence by equation (6) of Art. 239, we have

$$\int_0^a \int_0^b V dx dy = \iint V u dv du;$$

but we have not determined the limits of the integrations with respect to  $u$  and  $v$ , so that the result is of little value. We will now solve this example by following the steps indicated in the theory given above.

From the given equations connecting the old and new variables we eliminate  $u$ ; thus we have

$$y = \frac{vx}{1-v}, \quad \text{therefore } \frac{dy}{dv} = \frac{x}{(1-v)^2};$$

to the limits  $y = 0$  and  $y = b$ , correspond respectively  $v = 0$  and  $v = \frac{b}{b+x}$ ; thus

$$\int_0^a \int_0^b V \, dx \, dy = \int_0^a \int_0^{\frac{b}{b+x}} V_1 x (1-v)^{-2} \, dx \, dv.$$

We have now to change the *order* of integration in

$$\int_0^a \int_0^{\frac{b}{b+x}} V_1 x (1-v)^{-2} \, dx \, dv.$$

This question has been solved in Art. 235; hence we obtain

$$\begin{aligned} \int_0^a \int_0^b V \, dx \, dy &= \int_0^a \int_0^{\frac{b}{b+x}} V_1 x (1-v)^{-2} \, dx \, dv \\ &= \int_0^{\frac{b}{b+a}} \int_0^a V_1 x (1-v)^{-2} \, dv \, dx + \int_{\frac{b}{b+a}}^1 \int_0^{\frac{b(1-v)}{v}} V_1 x (1-v)^{-2} \, dv \, dx. \end{aligned}$$

We have now to change  $x$  for  $u$  where

$$x = u(1-v), \quad \frac{dx}{du} = 1-v;$$

thus we obtain

$$\int_0^{\frac{b}{b+a}} \int_0^{1-v} V' u \, dv \, du + \int_{\frac{b}{a+b}}^1 \int_0^{\frac{b}{v}} V' u \, dv \, du,$$

since to the limits 0 and  $a$  for  $x$  correspond respectively 0 and  $\frac{a}{1-v}$  for  $u$ , and to the limits 0 and  $\frac{b(1-v)}{v}$  for  $x$  correspond respectively 0 and  $\frac{b}{v}$  for  $u$ .

If  $a = b$  the transformed integral becomes

$$\int_0^{\frac{1}{2}} \int_0^{\frac{a}{1-v}} V'u \, dv \, du + \int_{\frac{1}{2}}^1 \int_0^{\frac{a}{v}} V'u \, dv \, du.$$

If  $a$  is made infinite, these two terms combine into the single expression

$$\int_0^1 \int_0^{\infty} V'u \, dv \, du.$$

241. *Second Example.* Required to transform

$$\int_0^c \int_0^{c-x} V \, dx \, dy,$$

having given  $y + x = u$ ,  $y = uv$ .

Perform the whole operation as before; so that we put

$$y = \frac{vx}{1-v} \quad \text{and} \quad \frac{dy}{dv} = \frac{x}{(1-v)^2}.$$

When  $y = 0$  we have  $v = 0$ , and when  $y = c - x$  we have  $v = \frac{c-x}{c}$ . Thus the integral is transformed into

$$\int_0^c \int_0^{\frac{c-x}{c}} V_1 x (1-v)^{-2} \, dx \, dv.$$

Now change the order of integration; thus we obtain

$$\int_0^1 \int_0^{c(1-v)} V_1 x (1-v)^{-2} \, dv \, dx.$$

Now put  $x = u(1-v)$  and  $\frac{dx}{du} = 1-v$ ; the limits of  $u$  will be 0 and  $c$ . Hence we have finally for the transformed integral

$$\int_0^1 \int_0^c V'u \, dv \, du.$$

242. *Third Example.* Transform  $\iint V dx dy$  to a double integral with the variables  $r$  and  $\theta$ , supposing

$$x = r \cos \theta, \quad y = r \sin \theta.$$

We may put  $\theta$  for  $v$  and  $r$  for  $u$  in the general formulæ; thus

$$\frac{dx dy}{du dv} - \frac{dx dy}{dv du} = r \cos^2 \theta + r \sin^2 \theta = r;$$

and the transformed integral is

$$\iint V' r d\theta dr.$$

This is a transformation with which the student is probably already familiar; the limits must of course be so taken that every element which enters into the original integral shall also occur in the transformed integral.

A particular case of this example may be noticed. Suppose the integral to be

$$\iint \phi(ax + by) dx dy;$$

by the present transformation this becomes

$$\iint \phi \{kr \cos(\theta - \alpha)\} r d\theta dr,$$

where  $k \cos \alpha = a$  and  $k \sin \alpha = b$ . Now put  $\theta - \alpha = \theta'$ , so that the integral becomes

$$\iint \phi(kr \cos \theta') r d\theta' dr;$$

then suppose  $r \cos \theta' = x'$  and  $r \sin \theta' = y'$  and the integral may be again changed to

$$\iint \phi(kx') dx' dy'.$$

Thus suppressing the accents we may write

$$\iint \phi(ax + by) dx dy = \iint \phi(kx) dx dy,$$

where  $k = \sqrt{a^2 + b^2}$ . The limits will generally be different in the two integrals; those on the right hand side must be determined by special examination, corresponding to given limits on the left hand side.

243. *Fourth Example.* Transform  $\int_0^c \int_0^x V dx dy$ , having given

$$x = au + bv, \quad y = bu + av, \quad a > b.$$

Eliminate  $u$ , thus  $ay - bx = (a^2 - b^2)v$ , and the first transformation gives

$$\frac{a^2 - b^2}{a} \int_0^c \int_{\frac{bx}{a^2 - b^2}}^{\frac{x}{a+b}} V_1 dx dv,$$

where  $V_1$  is what  $V$  becomes when we put  $\frac{bx}{a} + \frac{a^2 - b^2}{a}v$  for  $y$ . Next change the order of integration; this gives

$$\frac{a^2 - b^2}{a} \int_0^c \int_{(a+b)v}^{\frac{c}{a+b}} V_1 dv dx + \frac{a^2 - b^2}{a} \int_{-\frac{bc}{a^2 - b^2}}^0 \int_{-\frac{a^2 - b^2}{b}v}^c V_1 dv dx.$$

We have now to change from  $x$  to  $u$  by means of the equation  $x = au + bv$ , which gives  $\frac{dx}{du} = a$ ; the limits of  $u$  corresponding to the known limits of  $x$  are easily ascertained.

Thus we have finally for the transformed integral

$$(a^2 - b^2) \int_0^{\frac{c}{a+b}} \int_v^{\frac{c-bv}{a}} V' dv du + (a^2 - b^2) \int_{-\frac{bc}{a^2 - b^2}}^0 \int_{-\frac{av}{b}}^{\frac{c-bv}{a}} V' dv du.$$

The correctness of the transformation may be verified by supposing  $V$  to be some simple function of  $x$  and  $y$ ; for

example, if  $V$  be unity, the value of the original or of the transformed integral is  $\frac{c^3}{2}$ .

244. *Fifth Example.* The area of a surface is given by the integral

$$\iint dx dy \sqrt{\left\{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2\right\}} \quad (\text{Art. 170});$$

required to transform it into an integral with respect to  $\theta$  and  $\phi$ , having given

$$z = r \cos \theta, \quad x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi.$$

From the known equation to the surface  $z$  is given in terms of  $x$  and  $y$ ; hence by substituting we have an equation which gives  $r$  in terms of  $\theta$  and  $\phi$ .

We will first find the transformation for  $dx dy$ :

$$\frac{dx}{d\theta} = \frac{dr}{d\theta} \sin \theta \cos \phi + r \cos \theta \cos \phi,$$

$$\frac{dx}{d\phi} = \frac{dr}{d\phi} \sin \theta \cos \phi - r \sin \theta \sin \phi,$$

$$\frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta \sin \phi + r \cos \theta \sin \phi,$$

$$\frac{dy}{d\phi} = \frac{dr}{d\phi} \sin \theta \sin \phi + r \sin \theta \cos \phi.$$

Hence  $\frac{dx}{d\theta} \frac{dy}{d\phi} - \frac{dx}{d\phi} \frac{dy}{d\theta} = r \sin \theta \left( r \cos \theta + \frac{dr}{d\theta} \sin \theta \right);$

thus  $dx dy$  will be replaced by

$$r \sin \theta \left( r \cos \theta + \frac{dr}{d\theta} \sin \theta \right) d\phi d\theta.$$

We have next to transform

$$\sqrt{\left\{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2\right\}}.$$



We have 
$$\frac{dz}{d\theta} = \frac{dz}{dx} \frac{dx}{d\theta} + \frac{dz}{dy} \frac{dy}{d\theta},$$

$$\frac{dz}{d\phi} = \frac{dz}{dx} \frac{dx}{d\phi} + \frac{dz}{dy} \frac{dy}{d\phi}.$$

Also 
$$\frac{dz}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta,$$

$$\frac{dz}{d\phi} = \frac{dr}{d\phi} \cos \theta.$$

Thus  $\frac{dz}{dx}$  is a fraction of which the numerator is

$$\frac{dz}{d\theta} \frac{dy}{d\phi} - \frac{dz}{d\phi} \frac{dy}{d\theta},$$

that is, 
$$\left( \frac{dr}{d\theta} \cos \theta - r \sin \theta \right) \left( \frac{dr}{d\phi} \sin \theta \sin \phi + r \sin \theta \cos \phi \right)$$

$$- \frac{dr}{d\phi} \cos \theta \left( \frac{dr}{d\theta} \sin \theta \sin \phi + r \cos \theta \sin \phi \right),$$

that is,

$$- r \sin \phi \frac{dr}{d\phi} + r \sin \theta \cos \theta \cos \phi \frac{dr}{d\theta} - r^2 \sin^2 \theta \cos \phi,$$

and the denominator is

$$\frac{dx}{d\theta} \frac{dy}{d\phi} - \frac{dx}{d\phi} \frac{dy}{d\theta},$$

the value of which was found before; thus

$$\frac{dz}{dx} = \frac{r \sin \theta \cos \theta \cos \phi \frac{dr}{d\theta} - r \sin \phi \frac{dr}{d\phi} - r^2 \sin^2 \theta \cos \phi}{r \sin \theta \left( r \cos \theta + \sin \theta \frac{dr}{d\theta} \right)}.$$

Similarly

$$\frac{dz}{dy} = \frac{r \cos \phi \frac{dr}{d\phi} + r \sin \theta \cos \theta \sin \phi \frac{dr}{d\theta} - r^2 \sin^2 \theta \sin \phi}{r \sin \theta \left( r \cos \theta + \sin \theta \frac{dr}{d\theta} \right)};$$

therefore

$$1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2 = \frac{r^4 \sin^2 \theta + r^2 \left(\frac{dr}{d\phi}\right)^2 + r^2 \sin^2 \theta \left(\frac{dr}{d\theta}\right)^2}{r^2 \sin^2 \theta \left(r \cos \theta + \sin \theta \frac{dr}{d\theta}\right)^2},$$

and finally the transformed integral is

$$\iint \sqrt{\left\{r^2 \sin^2 \theta + \left(\frac{dr}{d\phi}\right)^2 + \sin^2 \theta \left(\frac{dr}{d\theta}\right)^2\right\}} r d\phi d\theta.$$

245. There will be no difficulty now in the transformation of a triple integral. Suppose that  $V$  is a function of  $x, y, z$ , and that  $\iiint V dx dy dz$  is to be transformed into a triple integral with respect to three new variables  $u, v, w$ , which are connected with  $x, y, z$  by three equations. From the investigation of Art. 239, we may anticipate that the result will take its simplest form when the old variables are given explicitly in terms of the new. Suppose then

$$x = f_1(u, v, w), \quad y = f_2(u, v, w), \quad z = f_3(u, v, w) \dots (1).$$

We first transform the integral with respect to  $z$  into an integral with respect to  $w$ . During the integration for  $z$  we regard  $x$  and  $y$  as constants; theoretically then we should from (1) express  $z$  as a function of  $x, y$ , and  $w$ , by eliminating  $u$  and  $v$ ; we should then find the differential coefficient of  $z$  with respect to  $w$  regarding  $x$  and  $y$  as constants. But we may obtain the required result by differentiating equations (1) as they stand,

thus 
$$\frac{df_1}{du} \frac{du}{dw} + \frac{df_1}{dv} \frac{dv}{dw} + \frac{df_1}{dw} = 0,$$

$$\frac{df_2}{du} \frac{du}{dw} + \frac{df_2}{dv} \frac{dv}{dw} + \frac{df_2}{dw} = 0,$$

$$\frac{df_3}{du} \frac{du}{dw} + \frac{df_3}{dv} \frac{dv}{dw} + \frac{df_3}{dw} = \frac{dz}{dw}.$$

Eliminate  $\frac{du}{dw}$  and  $\frac{dv}{dw}$ ; thus we find

$$\frac{dz}{dw} = \frac{N}{\frac{df_1}{du} \frac{df_2}{dv} - \frac{df_2}{du} \frac{df_1}{dv}},$$

$$\text{where } N = \frac{df_3}{dw} \left( \frac{df_1}{du} \frac{df_2}{dv} - \frac{df_2}{du} \frac{df_1}{dv} \right) + \frac{df_1}{dw} \left( \frac{df_3}{du} \frac{df_2}{dv} - \frac{df_3}{du} \frac{df_2}{dv} \right) \\ + \frac{df_2}{dw} \left( \frac{df_3}{du} \frac{df_1}{dv} - \frac{df_3}{du} \frac{df_1}{dv} \right).$$

Hence the integral is transformed into

$$\iiint V_1 \frac{N}{\frac{df_1}{du} \frac{df_2}{dv} - \frac{df_2}{du} \frac{df_1}{dv}} dx dy dw,$$

where  $V_1$  indicates what  $V$  becomes when for  $z$  its value in terms of  $x$ ,  $y$  and  $w$  is substituted. We must also determine the limits of  $w$  from the known limits of  $z$ . Next we may change the order of integration for  $y$  and  $w$ , and then proceed as before to remove  $y$  and introduce  $v$ . Then again we should change the order of integration for  $w$  and  $x$  and then for  $v$  and  $x$ , and finally remove  $x$  and introduce  $u$ . And in examples it might be advisable to go through the process step by step, in order to obtain the limits of the transformed integral.

We may however more simply ascertain the final formula thus. Transform the integral with respect to  $z$  into an integral with respect to  $w$  as above; then twice change the order of integration, so that we have

$$\iiint V_1 \frac{N}{\frac{df_1}{du} \frac{df_2}{dv} - \frac{df_2}{du} \frac{df_1}{dv}} dw dx dy.$$

Now we have to transform the double integral with respect to  $x$  and  $y$  into a double integral with respect to  $u$  and  $v$  by means of the first two of equations (1). Hence we know by Art. 239 that the symbol  $dx dy$  will be replaced by

$$\left(\frac{df_1}{du} \frac{df_2}{dv} - \frac{df_2}{du} \frac{df_1}{dv}\right) dv du;$$

and the integral is finally transformed into

$$\iiint V' N dw dv du,$$

where  $V'$  is what  $V$  becomes when for  $x, y,$  and  $z,$  their values in terms of  $u, v,$  and  $w$  are substituted.

The student will now have no difficulty in investigating the more complex case, in which the old and new variables are connected by equations of the form

$$\phi_1(x, y, z, u, v, w) = 0,$$

$$\phi_2(x, y, z, u, v, w) = 0,$$

$$\phi_3(x, y, z, u, v, w) = 0.$$

Here it will be found that

$$\frac{dz}{dw} = \frac{N_1}{D_1}, \quad \frac{dy}{dv} = \frac{N_2}{D_2}, \quad \frac{dx}{du} = \frac{N_3}{D_3};$$

also that  $N_2 = D_1,$  and  $N_3 = D_2.$


Thus  $\iiint V dx dy dz = \iiint V' \frac{N_1}{D_3} du dv dw,$

where

$$N_1 = \frac{d\phi_1}{dw} \left(\frac{d\phi_2}{du} \frac{d\phi_3}{dv} - \frac{d\phi_3}{du} \frac{d\phi_2}{dv}\right) + \frac{d\phi_2}{dw} \left(\frac{d\phi_3}{du} \frac{d\phi_1}{dv} - \frac{d\phi_1}{du} \frac{d\phi_3}{dv}\right) \\ + \frac{d\phi_3}{dw} \left(\frac{d\phi_1}{du} \frac{d\phi_2}{dv} - \frac{d\phi_2}{du} \frac{d\phi_1}{dv}\right),$$

and  $-D_3$  is equal to a similar expression with  $x, y, z$  instead of  $u, v, w$  respectively.

246. It may be instructive to illustrate these transformations geometrically. We begin with the double integral.



Let  $\iint V dx dy$  be a double integral, which is to be taken for all the values of  $x$  and  $y$  comprised within the boundary  $ABCD$ . Suppose the variables  $x$  and  $y$  connected with two new variables  $u$  and  $v$  by the equations

$$x = f_1(u, v), \quad y = f_2(u, v) \dots \dots \dots (1).$$

From these equations let  $u$  and  $v$  be found in terms of  $x$  and  $y$ , so that we may write

$$u = F_1(x, y), \quad v = F_2(x, y) \dots \dots \dots (2).$$

Now by ascribing any constant value to  $u$  the first equation of (2) may be considered as representing a curve, and by giving in succession different constant values to  $u$ , we have a series of such curves. Let then  $APQC$  be a curve, at every point of which  $F_1(x, y)$  has a certain constant value  $u$ ; and let  $A'SRC'$  be a curve, at every point of which  $F_1(x, y)$  has a certain constant value  $u + \delta u$ . Similarly let  $BPSD$  be a curve, at every point of which  $F_2(x, y)$  has a certain constant value  $v$ ; and let  $B'QRD'$  be a curve, at every point of which

$F_2(x, y)$  has a certain constant value  $v + \delta v$ . Let  $x, y$  now denote the co-ordinates of  $P$ ; we shall proceed to express the co-ordinates of  $Q, S$ , and  $R$ .

The co-ordinates of  $Q$  are found from those of  $P$ , by changing  $v$  into  $v + \delta v$ ; hence by (1) they are ultimately, when  $\delta v$  is indefinitely small,

$$x + \frac{dx}{dv} \delta v, \text{ and } y + \frac{dy}{dv} \delta v.$$

Similarly the co-ordinates of  $S$  are found from those of  $P$  by changing  $u$  into  $u + \delta u$ ; hence by (1) they are ultimately

$$x + \frac{dx}{du} \delta u, \text{ and } y + \frac{dy}{du} \delta u.$$

The co-ordinates of  $R$  are found from those of  $P$  by changing both  $u$  into  $u + \delta u$  and  $v$  into  $v + \delta v$ ; hence by (1) they are ultimately

$$x + \frac{dx}{du} \delta u + \frac{dx}{dv} \delta v, \text{ and } y + \frac{dy}{du} \delta u + \frac{dy}{dv} \delta v.$$

These results shew that  $P, Q, R, S$  are ultimately situated at the angular points of a parallelogram, and the area of this parallelogram may be taken without error in the limit for the area of the curvilinear figure  $PQRS$ . The expression for the area of the triangle  $PQR$  in terms of the co-ordinates of its angular points is known, (see Plane Co-ordinate Geometry, Art. 11), and the area of the parallelogram is double that of the triangle. Hence we have ultimately for the area of  $PQRS$  the expression


$$\pm \left( \frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du} \right) \delta u \delta v.$$

Thus it is obvious that the integral  $\iint V dx dy$  may be replaced by

$$\pm \iint V' \left( \frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du} \right) du dv;$$

the ambiguity of sign would disappear in an example in which the limits of integration were known. In finding the value of the transformed integral, we may suppose that we first integrate with respect to  $v$ , so that  $u$  is kept constant; this amounts to taking all the elements such as  $PQRS$ , which form a strip such as  $AA'C'C$ . Then the integration with respect to  $u$  amounts to taking all such strips as  $AA'C'C$  which are contained within the assigned boundary  $ABCD$ .

247. We proceed to illustrate geometrically the transformation of a triple integral.



Let  $\iiint V dx dy dz$  be a triple integral, which is to be taken for all values of  $x$ ,  $y$ , and  $z$  comprised between certain assigned limits. Suppose the variables  $x$ ,  $y$ , and  $z$  connected with three new variables  $u$ ,  $v$ ,  $w$  by the equations

$$x = f_1(u, v, w), \quad y = f_2(u, v, w), \quad z = f_3(u, v, w) \dots (1).$$

From these equations let  $u$ ,  $v$ , and  $w$  be found in terms of  $x$ ,  $y$ , and  $z$ , so that we may write

$$u = F_1(x, y, z), \quad v = F_2(x, y, z), \quad w = F_3(x, y, z) \dots (2).$$

Now by ascribing any constant value to  $u$ , the first equation of (2) may be considered as representing a surface, and by giving in succession different constant values to  $u$  we have a series of such surfaces. Suppose there to be a surface at every point of which  $F_1(x, y, z)$  has the constant value  $u$ , and let the four points  $P, B, D, C$ , be in that surface; also suppose there to be a surface at every point of which  $F_1(x, y, z)$  has the constant value  $u + \delta u$ , and let the four points  $A, F, G, E$  be in that surface. Similarly suppose  $P, A, E, C$  to be in a surface at every point of which  $F_2(x, y, z)$  has the constant value  $v$ , and  $B, D, G, F$  to be in a surface at every point of which  $F_2(x, y, z)$  has the constant value  $v + \delta v$ . Lastly suppose  $P, A, F, B$  to be in a surface at every point of which  $F_3(x, y, z)$  has the constant value  $w$ , and  $C, D, G, E$  to be in a surface at every point of which  $F_3(x, y, z)$  has the constant value  $w + \delta w$ .

Let  $x, y, z$  now denote the co-ordinates of  $P$ ; we shall proceed to express the co-ordinates of the other points. The co-ordinates of  $A$  are found from those of  $P$  by changing  $u$  into  $u + \delta u$ ; hence by (1) they are ultimately when  $\delta u$  is indefinitely small,

$$x + \frac{dx}{du} \delta u, \quad y + \frac{dy}{du} \delta u, \quad z + \frac{dz}{du} \delta u.$$

The co-ordinates of  $B$  are found from those of  $P$  by changing  $v$  into  $v + \delta v$ ; hence by (1) they are ultimately

$$x + \frac{dx}{dv} \delta v, \quad y + \frac{dy}{dv} \delta v, \quad z + \frac{dz}{dv} \delta v.$$

Similarly the co-ordinates of  $C$  are ultimately

$$x + \frac{dx}{dw} \delta w, \quad y + \frac{dy}{dw} \delta w, \quad z + \frac{dz}{dw} \delta w.$$

The co-ordinates of  $D$  are found from those of  $P$  by changing  $v$  into  $v + \delta v$ , and  $w$  into  $w + \delta w$ ; hence by (1) they are ultimately

$$x + \frac{dx}{dv} \delta v + \frac{dx}{dw} \delta w, \quad y + \frac{dy}{dv} \delta v + \frac{dy}{dw} \delta w, \quad z + \frac{dz}{dv} \delta v + \frac{dz}{dw} \delta w.$$

Similarly the co-ordinates of  $E, F$  and  $G$  may be found.



These results shew that  $P, A, B, C, D, E, F, G$  are ultimately situated at the angular points of a parallelepiped; and the volume of this parallelepiped may be taken without error in the limit for the volume of the solid bounded by the six surfaces which we have referred to. Now by a known theorem the volume of a tetrahedron can be expressed in terms of the co-ordinates of its angular points, and the volume of the parallelepiped  $PG$  is six times that of the tetrahedron  $ABPC$ . Hence finally we have for the volume of the parallelepiped

$$\pm \left\{ \frac{dx}{du} \left( \frac{dy}{dv} \frac{dz}{dw} - \frac{dy}{dw} \frac{dz}{dv} \right) + \frac{dy}{du} \left( \frac{dz}{dv} \frac{dx}{dw} - \frac{dz}{dw} \frac{dx}{dv} \right) \right. \\ \left. + \frac{dz}{du} \left( \frac{dx}{dv} \frac{dy}{dw} - \frac{dx}{dw} \frac{dy}{dv} \right) \right\} \delta u \delta v \delta w = \pm N \delta u \delta v \delta w \text{ say.}$$

Hence the triple integral is transformed into

$$\pm \iiint V' N du dv dw;$$

the ambiguity in sign would disappear in an example where the limits of integration were known.

248. We have now given the theory of the transformation of double and triple integrals; the essential point in our investigation is, that we have shewn how to remove the old variables and replace them by the new variables *one at a time*. We recommend the student to pay attention to this point, as we conceive that the theory of the subject is thus made clear and simple, and at the same time the limits of the transformed integral can be more easily ascertained. We do not lay any stress on the geometrical *illustrations* in the two preceding articles; they require much more development before they can be accepted as rigid *demonstrations*.

249. Before leaving the subject we will briefly indicate the method formerly used in solving the problem. This method we have not brought prominently forward, partly because it gives no assistance in determining the new limits,

and partly on account of its obscurity; the latter defect has been frequently noticed by writers on the subject.

Suppose  $\iint V dx dy$  is to be transformed into an integral with respect to two new variables  $u$  and  $v$  of which the old variables are known functions.

Let the variables undergo infinitesimal changes: thus

$$dx = \frac{dx}{du} du + \frac{dx}{dv} dv \dots \dots \dots (1),$$

$$dy = \frac{dy}{du} du + \frac{dy}{dv} dv \dots \dots \dots (2).$$

Now in the original expression  $V dx dy$  in forming  $dx$  we suppose  $y$  constant, that is  $dy = 0$ ; hence (2) becomes

$$0 = \frac{dy}{du} du + \frac{dy}{dv} dv \dots \dots \dots (3),$$

find  $dv$  from this and substitute it in (1); thus

$$dx = \frac{\frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du}}{\frac{dy}{dv}} du \dots \dots \dots (4).$$

Again, in forming  $dy$  in  $V dx dy$  we suppose  $x$  constant, that is,  $dx = 0$ ; hence by (4) we must suppose  $du = 0$ ; thus from (2)

$$dy = \frac{dy}{dv} dv \dots \dots \dots (5).$$

From (4) and (5)

$$dx dy = \left( \frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du} \right) du dv;$$

and  $\iint V dx dy$  becomes

$$\iint V' \left( \frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du} \right) du dv.$$

With respect to the limits of integration we can only give the general direction, that the new limits must be so taken as to include every element which was included by the old limits.

250. Similarly in transforming a triple integral

$$\iiint V \, dx \, dy \, dz$$

the process was as follows. Let the new variables be  $u, v, w$ ; in forming  $dz$  we must suppose  $x$  and  $y$  constant; thus we have

$$dz = \frac{dz}{du} du + \frac{dz}{dv} dv + \frac{dz}{dw} dw,$$

$$0 = \frac{dx}{du} du + \frac{dx}{dv} dv + \frac{dx}{dw} dw,$$

$$0 = \frac{dy}{du} du + \frac{dy}{dv} dv + \frac{dy}{dw} dw,$$

thus 
$$dz = \frac{N dw}{\frac{dx}{du} \frac{dy}{dv} - \frac{dx}{dv} \frac{dy}{du}} \dots \dots \dots (1),$$

where  $N$  has the same value as in Art. 247.

Next in forming  $dy$  we have to regard  $x$  and  $z$  as constant; hence by (1) we must regard  $w$  as constant; thus we have

$$dy = \frac{dy}{du} du + \frac{dy}{dv} dv,$$

$$0 = \frac{dx}{du} du + \frac{dx}{dv} dv;$$

therefore 
$$dy = \frac{\left(\frac{dy}{dv} \frac{dx}{du} - \frac{dy}{du} \frac{dx}{dv}\right) dv}{\frac{dx}{du}} \dots \dots \dots (2).$$

And lastly in forming  $dx$  we suppose  $y$  and  $z$  constant, that is, by (1) and (2) we suppose  $w$  and  $v$  constant; thus

$$dx = \frac{dx}{du} du \dots \dots \dots (3).$$

From (1), (2) and (3)

$$dx dy dz = N du dv dw.$$

251. The student who wishes to extend his knowledge of this subject may be assisted by the following references. Lacroix, *Calcul Diff. et Intégral*, Vol. II. p. 205; also the references to the older authorities will be found in page XI of the table prefixed to this volume. De Morgan, *Diff. and Integral Calculus*, p. 392. Moigno, *Calcul Diff. et Intégral*, Vol. II. p. 214; Ostrogradsky, *Mémoires de l'Académie de St Pétersbourg*, Sixième Série, 1838, p. 401. Catalan, *Mémoires Couronnés par l'Académie... de Bruxelles*, Vol. XIV. p. 1. Boole, *Cambridge Mathematical Journal*, Vol. IV. p. 20. Cauchy, *Exercices d'Analyse et de Physique Mathématique*, Vol. IV. p. 128. De Morgan, *Transactions of the Cambridge Phil. Society*, Vol. IX. p. [133.]

EXAMPLES.

1. Shew that if  $x = a \sin \theta \sin \phi$  and  $y = b \cos \theta \sin \phi$  the double integral  $\iint dx dy$  is transformed into

$$\pm \iint ab \sin \phi \cos \phi d\phi d\theta.$$

2. If  $x = u \sin \alpha + v \cos \alpha$  and  $y = u \cos \alpha - v \sin \alpha$ , prove that

$$\iint f(x, y) \frac{dx dy}{\sqrt{(1-x^2-y^2)}} = \iint f_1(u, v) \frac{du dv}{\sqrt{(1-u^2-v^2)}}.$$

3. Prove that

$$\int_0^{\infty} \int_0^{\infty} \phi(a^2x^2 + b^2y^2) dx dy = \frac{\pi}{4ab} \int_0^{\infty} \phi(x) dx.$$

4. Transform  $\iint V dx dy$ , where  $y = xu$  and  $x = \frac{v}{1+u}$ .

If the limits of  $y$  be 0 and  $x$  and the limits of  $x$  be 0 and  $a$ , find the limits in the transformed integral.

$$\text{Result. } \int_0^1 \int_0^{a(1+u)} V' v (1+u)^{-2} du dv.$$

5. Transform  $\iint e^{-(x^2+2xy \cos \alpha + y^2)} dx dy$  from rectangular to polar co-ordinates, and thence shew that if the limits both of  $x$  and  $y$  be zero and infinity, the value of the integral will be  $\frac{\alpha}{2 \sin \alpha}$ .

6. Prove by transforming the expression from rectangular to polar co-ordinates that the value of the definite integral

$$\int_0^{\infty} \int_0^{\infty} e^{-(x^2+2x^2y^2 \cos \alpha + y^4)} dx dy$$

is equal to  $\frac{1}{2} \sqrt{\pi} F\left(\sin \frac{\alpha}{2}\right)$  where  $F\left(\sin \frac{\alpha}{2}\right)$  denotes a complete elliptic function of the first order, of which  $\sin \frac{\alpha}{2}$  is the modulus.

7. Apply the transformation from rectangular to polar co-ordinates in double integrals to shew that

$$\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{a dx dy}{(x^2 + y^2 + a^2)^{\frac{3}{2}} (x^2 + y^2 + a'^2)^{\frac{3}{2}}} = \frac{2\pi}{a+a'}.$$

8. Transform the double integral  $\iint f(x, y) dx dy$  into one

in which  $r$  and  $\theta$  shall be the independent variables, having given

$$x = r \cos \theta + a \sin \theta, \quad y = r \sin \theta + a \cos \theta.$$

*Result.*

$$\iint f(r \cos \theta + a \sin \theta, r \sin \theta + a \cos \theta) (a \sin 2\theta - r) d\theta dr.$$

9. Transform  $\iint e^{-x^2-y^2} dx dy$  into a double integral where  $r$  and  $t$  are the independent variables where  $\frac{y}{x} = t$  and  $r^2 = x^2 + y^2$ ; and if the limits of  $x$  and  $y$  be each 0 and  $\infty$ , find the limits of  $r$  and  $t$ .

$$\text{Result. } \int_0^\infty \int_0^\infty \frac{e^{-r^2} r dr dt}{1+t^2}.$$

10. If  $x$  and  $y$  are given as functions of  $r$  and  $\theta$ , transform the integral  $\iiint dx dy dz$  into another where  $r$ ,  $\theta$  and  $z$  are the variables; and if  $x = r \cos \theta$  and  $y = r \sin \theta$ , find the volume included by the four surfaces whose equations are  $r = a$ ,  $z = 0$ ,  $\theta = 0$ , and  $z = mr \cos \theta$ .

$$\text{Result. The volume} = \int_0^{\frac{\pi}{2}} \int_0^a r^2 m \cos \theta d\theta dr = \frac{ma^3}{3}.$$

11. If  $\alpha x = yz$ ,  $\beta y = zx$ ,  $\gamma z = xy$ , shew that

$$\iiint f(\alpha, \beta, \gamma) d\alpha d\beta d\gamma = 4 \iiint f\left(\frac{yz}{x}, \frac{zx}{y}, \frac{xy}{z}\right) dx dy dz.$$

12. Transform  $\iiint V dx_1 dx_2 dx_3 dx_4$  to  $r, \theta, \phi$  and  $\psi$  when

$$\begin{aligned} x_1 &= r \sin \theta \cos \phi, & x_3 &= r \cos \theta \cos \psi, \\ x_2 &= r \sin \theta \sin \phi, & x_4 &= r \cos \theta \sin \psi. \end{aligned}$$

$$\text{Result. } \iiint V r^3 \sin \theta \cos \theta dr d\theta d\phi d\psi.$$

13. Find the elementary area included between the curves  $\phi(x, y) = u$ ,  $\psi(x, y) = v$ , and the curves obtained by giving to the parameters  $u$  and  $v$  indefinitely small increments.

Find the area included between a parabola and the tangents at the extremities of the latus rectum by dividing the area by a series of parabolas which touch these tangents and by a series of lines drawn from the intersection of the tangents.

14. Transform the triple integral  $\iiint f(x, y, z) dx dy dz$  into one in which  $r, y, z$  are the independent variables, having given  $\psi(x, y, z, r) = 0$ ; and change the variables in the above integral from  $x, y, z$  to  $r, \theta, \phi$ , having given

$$\psi(x, y, z, r) = 0, \quad \psi_1(y, z, r, \theta) = 0, \quad \psi_2(z, r, \theta, \phi) = 0.$$

$$\text{Result.} \quad - \iiint \frac{\frac{d\psi}{dr} \frac{d\psi_1}{d\theta} \frac{d\psi_2}{d\phi}}{\frac{d\psi}{dx} \frac{d\psi_1}{dy} \frac{d\psi_2}{dz}} f_1(r, \theta, \phi) dr d\theta d\phi.$$

15. Transform the double integral

$$\iint dx dy \sqrt{\left\{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2\right\}}$$

in which  $x, y, z$  are connected by the equation  $x^2 + y^2 + z^2 = 1$ , to an integral in terms of  $\theta$  and  $\phi$ , having these relations

$$\begin{aligned} x &= \sin \phi \sqrt{1 - m^2 \sin^2 \theta}, & y &= \cos \theta \cos \phi, \\ z &= \sin \theta \sqrt{1 - n^2 \sin^2 \phi}, & m^2 + n^2 &= 1. \end{aligned}$$

Hence prove that

$$\int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \frac{m^2 \cos^2 \theta + n^2 \cos^2 \phi}{\sqrt{1 - m^2 \sin^2 \theta} \sqrt{1 - n^2 \sin^2 \phi}} d\theta d\phi = \frac{\pi}{2}.$$

16. Transform the integral  $\iiint dx dy dz$  to  $r, \theta, \phi$  where

$$x = r \sin \phi \sqrt{1 - n^2 \cos^2 \theta}, \quad y = r \cos \phi \sin \theta, \\ z = r \cos \theta \sqrt{\cos^2 \phi + n^2 \sin^2 \phi}.$$

*Result.* 
$$\iiint \frac{r^2 \{ (n^2 - 1) \cos^2 \phi - n^2 \sin^2 \theta \} dr d\theta d\phi}{\sqrt{1 - n^2 \cos^2 \theta} \sqrt{\cos^2 \phi + n^2 \sin^2 \phi}}.$$

17. Transform the expression  $\iint \frac{r^3}{3} \sin \theta d\theta d\phi$  for a volume to rectangular co-ordinates.

*Result.*  $\frac{1}{3} \iint (z - px - qy) dx dy$ ; this should be interpreted geometrically.

18. Prove that

$$\left\{ \int_0^\infty e^{-x^{2n}} dx \right\}^2 = \frac{1}{2} \int_0^\infty e^{-x^n} dx \int_0^\infty \frac{dx}{(1 + x^{2n})^{\frac{1}{n}}}.$$

(See Arts. 263 and 66; and transform as in Art. 242.)

19. If  $x_1 = r \cos \theta_1,$

$$x_2 = r \sin \theta_1 \cos \theta_2,$$

$$x_3 = r \sin \theta_1 \sin \theta_2 \cos \theta_3,$$

.....

$$x_{n-1} = r \sin \theta_1 \sin \theta_2 \dots \sin \theta_{n-2} \cos \theta_{n-1},$$

$$x_n = r \sin \theta_1 \sin \theta_2 \dots \sin \theta_{n-2} \sin \theta_{n-1},$$

shew that  $\iiint \dots V dx_1 dx_2 \dots dx_n$

$$= \pm \iiint \dots V' r^{n-1} (\sin \theta_1)^{n-2} (\sin \theta_2)^{n-3} \\ \dots \sin \theta_{n-2} dr d\theta_1 d\theta_2 \dots d\theta_{n-1},$$

where  $V$  is any function of  $x_1, x_2, \dots, x_n$ , and  $V'$  what this function becomes when the variables are changed.



## CHAPTER XII.

## DEFINITE INTEGRALS.

252. WHEN the indefinite integral of a function is known, we can immediately obtain the value of the *definite integral* corresponding to any assigned limits of the variable. Sometimes however we are able by special methods to assign the value of a *definite* integral when we cannot express the indefinite integral in a finite form; sometimes without actually finding the value of a definite integral we can shew that it possesses important properties. In some cases in which the indefinite integral of a function can be found, the definite integral between certain limits may have a value which is worthy of notice, on account of the simple form in which it may be expressed. We shall in the present chapter give examples of these general statements.

253. Suppose  $f(x)$  and  $F(x)$  rational algebraical functions of  $x$ , and  $f(x)$  of lower dimensions than  $F(x)$ , and suppose the equation  $F(x) = 0$  to have no real roots; it is required to find the value of

$$\int_{-\infty}^{\infty} \frac{f(x)}{F(x)} dx.$$

It will be seen that under the above suppositions, the expression to be integrated never becomes infinite for real values of  $x$ .

Let  $\alpha + \beta \sqrt{-1}$  and  $\alpha - \beta \sqrt{-1}$  represent a pair of the imaginary roots of  $F(x) = 0$ ; then the corresponding quadratic

fraction of the series into which  $\frac{f(x)}{F'(x)}$  can be decomposed, may be represented by

$$\frac{2A(x-\alpha) + 2B\beta}{(x-\alpha)^2 + \beta^2},$$

the constants  $A$  and  $B$  being found from the equation

$$A - B\sqrt{-1} = \frac{f\{\alpha + \beta\sqrt{-1}\}}{F'\{\alpha + \beta\sqrt{-1}\}} \quad (\text{Art. 21}).$$

Now 
$$\int \frac{2B\beta dx}{(x-\alpha)^2 + \beta^2} = 2B \tan^{-1} \frac{x-\alpha}{\beta},$$

therefore 
$$\int_{-\infty}^{\infty} \frac{2B\beta dx}{(x-\alpha)^2 + \beta^2} = 2B\pi.$$

Also 
$$\int_{-\infty}^{\infty} \frac{(x-\alpha) dx}{(x-\alpha)^2 + \beta^2} = \int_{-\infty}^{\infty} \frac{tdt}{t^2 + \beta^2},$$

and it is obvious that the latter integral between the assigned limits is zero, for the negative part is numerically equal to the positive part. Thus  $2B\pi$  represents the part of the integral corresponding to the pair of imaginary roots under consideration.

If then we suppose  $F(x)$  to be of  $2n$  dimensions, and  $B_1, B_2, \dots, B_n$  to be the  $n$  terms of which we have taken  $B$  as the type, we have

$$\int_{-\infty}^{\infty} \frac{f(x)}{F'(x)} dx = 2\pi \{B_1 + B_2 + \dots + B_n\}.$$

254. As an example of the preceding article we take

$$\int_{-\infty}^{\infty} \frac{x^{2m} dx}{1 + x^{2n}},$$

where  $m$  and  $n$  are positive integers, and  $m$  less than  $n$ . Here

$$A - B\sqrt{-1} = \frac{1}{2n \{\alpha + \beta\sqrt{-1}\}^{2n-2m-1}},$$

and it is known from the theory of equations that the values of  $\alpha + \beta \sqrt{-1}$  are obtained from the expression

$$\cos \frac{(2r+1)\pi}{2n} + \sqrt{-1} \sin \frac{(2r+1)\pi}{2n},$$

by giving to  $r$  successively the values 0, 1, 2, ..... up to  $n-1$ .

Thus, by Demoiivre's theorem,

$$\{\alpha + \beta \sqrt{-1}\}^{2n-2m-1} = \cos \phi + \sqrt{-1} \sin \phi,$$

where

$$\phi = (2n - 2m - 1) \frac{(2r+1)\pi}{2n} = (2r+1)\pi - (2r+1) \frac{(2m+1)\pi}{2n};$$

so that

$$\cos \phi + \sqrt{-1} \sin \phi = -\cos (2r+1)\theta + \sqrt{-1} \sin (2r+1)\theta,$$

where

$$\theta = \frac{2m+1}{2n} \pi.$$

Hence

$$\begin{aligned} A - B\sqrt{-1} &= \frac{1}{2n - \cos (2r+1)\theta + \sqrt{-1} \sin (2r+1)\theta} \\ &= -\frac{\cos (2r+1)\theta + \sqrt{-1} \sin (2r+1)\theta}{2n}; \end{aligned}$$

$$\text{therefore } B = \frac{\sin (2r+1)\theta}{2n}.$$

Hence

$$\int_{-\infty}^{\infty} \frac{x^{2m} dx}{1+x^{2n}} = \frac{\pi}{n} \{\sin \theta + \sin 3\theta + \sin 5\theta + \dots + \sin (2n-1)\theta\}.$$

The sum of the series of sines is shewn in works on Trigonometry to be  $\frac{\sin^2 n\theta}{\sin \theta}$ , and in the present case  $n\theta = \frac{2m+1}{2}\pi$ , so that  $\sin^2 n\theta = 1$ . Therefore

$$\int_{-\infty}^{\infty} \frac{x^{2m} dx}{1+x^{2n}} = \frac{\pi}{n \sin \frac{2m+1}{2n} \pi}.$$

It is obvious that  $\int_0^{\infty} \frac{x^{2m} dx}{1+x^{2n}}$  is half of the above result, that is,

$$\int_0^{\infty} \frac{x^{2m} dx}{1+x^{2n}} = \frac{\pi}{2n \sin \frac{2m+1}{2n} \pi}.$$

255. In the last formula of the preceding article put  $x^{2m} = y$ , and suppose  $\frac{2m+1}{2n} = k$ ; thus we obtain

$$\int_0^{\infty} \frac{y^{k-1} dy}{1+y} = \frac{\pi}{\sin k\pi}.$$

This result holds when  $k$  has any value comprised between 0 and 1; for the only restriction on the positive integers  $m$  and  $n$  is that  $m$  must be less than  $n$ , and therefore by properly choosing  $m$  and  $n$  we may make  $\frac{2m+1}{2n}$  equal to any assigned proper fraction.

In the last result put  $x^r$  for  $y$ , where  $r$  is any positive quantity; thus

$$\int_0^{\infty} \frac{rx^{kr-r} x^{r-1} dx}{1+x^r} = \frac{\pi}{\sin kr\pi},$$

that is, 
$$\int_0^{\infty} \frac{x^{kr-1} dx}{1+x^r} = \frac{\pi}{r \sin kr\pi}.$$

Let  $kr = s$ ; thus 
$$\int_0^{\infty} \frac{x^{s-1} dx}{1+x^r} = \frac{\pi}{r \sin \frac{s}{r} \pi}.$$

The only restriction on the positive quantities  $r$  and  $s$  is that  $s$  must be less than  $r$ .

*Eulerian Integrals.*

256. The definite integral

$$\int_0^1 x^{l-1} (1-x)^{m-1} dx$$

is called the *first Eulerian integral*; we shall denote it by the symbol  $B(l, m)$ .

The definite integral

$$\int_0^{\infty} e^{-x} x^{n-1} dx$$

is called the *second Eulerian integral*; it is denoted by the symbol  $\Gamma(n)$ .

We shall now give some of the properties of these integrals; the constants in these integrals which we have denoted by  $l, m, n$ , are supposed *positive* in all that follows.

257. In the first Eulerian integral put  $x = 1 - z$ ;

thus 
$$\int_0^1 x^{l-1} (1-x)^{m-1} dx = \int_0^1 z^{m-1} (1-z)^{l-1} dz;$$

this shews that the constants  $l$  and  $m$  may be interchanged without altering the value of the integral; that is,

$$B(l, m) = B(m, l).$$

Again in the first Eulerian integral put  $x = \frac{y}{1+y}$ ; thus

$$\int_0^1 x^{l-1} (1-x)^{m-1} dx = \int_0^{\infty} \frac{y^{l-1} dy}{(1+y)^{l+m}}.$$

In the same integral put  $x = \frac{1}{1+y}$ ; thus

$$\int_0^1 x^{l-1} (1-x)^{m-1} dx = \int_0^{\infty} \frac{y^{m-1} dy}{(1+y)^{l+m}}.$$

258. Let  $e^{-x} = y$ , so that  $x = \log \frac{1}{y}$ ; then we have

$$\int_0^{\infty} e^{-x} x^{n-1} dx = \int_0^1 \left(\log \frac{1}{y}\right)^{n-1} dy,$$

which consequently gives another form of  $\Gamma(n)$ .

259. We have by integration by parts

$$\int e^{-x} x^n dx = -e^{-x} x^n + n \int e^{-x} x^{n-1} dx;$$

and  $e^{-x} x^n$  vanishes when  $x=0$ , and also when  $x=\infty$ . (See *Dif. Cal.* Art. 153); thus

$$\int_0^{\infty} e^{-x} x^n dx = n \int_0^{\infty} e^{-x} x^{n-1} dx;$$

that is  $\Gamma(n+1) = n\Gamma(n)$ ..... (1).

Since  $\int e^{-x} dx = -e^{-x}$  we have  $\int_0^{\infty} e^{-x} dx = 1$ ; that is

$$\Gamma(1) = 1$$
 ..... (2).

From (1) and (2) we see that if  $n$  be an integer

$$\Gamma(n+1) = \underline{n}.$$

When  $n$  is not an integer we may by repeated use of equation (1) make the value of  $\Gamma(n)$  where  $n$  is greater than unity depend on that of  $\Gamma(m)$  where  $m$  is less than unity.

260. By assuming  $kx = z$  we have

$$\int_0^{\infty} e^{-kx} x^{n-1} dx = \frac{1}{k^n} \int_0^{\infty} e^{-z} z^{n-1} dz = \frac{\Gamma(n)}{k^n}.$$

261. We shall now prove an important equation which connects the two Eulerian integrals.

Integrate the double integral  $\int_0^\infty \int_0^\infty x^{l+m-1} y^{m-1} e^{-(l+y)x} dy dx$  first with respect to  $x$ ; we thus obtain, by Art. 260,

$$\Gamma(l+m) \int_0^\infty \frac{y^{m-1} dy}{(1+y)^{l+m}}.$$

Again integrate the same double integral first with respect to  $y$ ; we thus obtain

$$\Gamma(m) \int_0^\infty \frac{e^{-x} x^{l+m-1}}{x^m} dx,$$

that is,

$$\Gamma(m) \int_0^\infty e^{-x} x^{l-1} dx,$$

that is,

$$\Gamma(m) \Gamma(l).$$

Hence

$$\int_0^\infty \frac{y^{m-1} dy}{(1+y)^{l+m}} = \frac{\Gamma(l) \Gamma(m)}{\Gamma(l+m)}.$$

Hence, by Art. 257,

$$B(l, m) = \frac{\Gamma(l) \Gamma(m)}{\Gamma(l+m)}.$$

262. In the result of the preceding article, suppose  $l+m=1$ ; thus, if  $m$  is less than unity,

$$\int_0^\infty \frac{y^{m-1} dy}{1+y} = \Gamma(m) \Gamma(1-m),$$

since  $\Gamma(1) = 1$ . Hence, by Art. 255, if  $m$  is less than unity

$$\Gamma(m) \Gamma(1-m) = \frac{\pi}{\sin m\pi}.$$

263. Put  $m = \frac{1}{2}$  in the last result; then

$$\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) = \pi,$$

therefore

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$$

We will give another proof of the last result.

Let  $u = \int_0^{\infty} e^{-x^2} dx$ ; then it is obvious that  $u$  also

$$= \int_0^{\infty} e^{-y^2} dy;$$

thus

$$\begin{aligned} u^2 &= \int_0^{\infty} e^{-x^2} dx \times \int_0^{\infty} e^{-y^2} dy \\ &= \int_0^{\infty} \int_0^{\infty} e^{-x^2-y^2} dx dy \quad (\text{Art. 66}). \end{aligned}$$

This double integral is shewn in Art. 204 to be

$$= \frac{1}{4} \int_0^{2\pi} \int_0^{\infty} e^{-r^2} r d\theta dr = \frac{\pi}{4},$$

therefore

$$u = \frac{\sqrt{\pi}}{2}.$$

Now

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{\infty} e^{-x} x^{-\frac{1}{2}} dx; \text{ put } x = y^2,$$

thus

$$\Gamma\left(\frac{1}{2}\right) = 2 \int_0^{\infty} e^{-y^2} dy = 2u = \sqrt{\pi}.$$

264. We shall now give an expression for  $\Gamma(n)$  that will afford another proof of the result in Art. 262. We know that the limit of  $\frac{x^h - 1}{h}$  when  $h$  is indefinitely diminished is  $\log x$ ; hence

$$\left(\log \frac{1}{x}\right)^{n-1} = \text{limit of } \left(\frac{1-x^h}{h}\right)^{n-1};$$

so we may write

$$\left(\log \frac{1}{x}\right)^{n-1} = \left(\frac{1-x^h}{h}\right)^{n-1} + y,$$

where  $y$  is a quantity that diminishes without limit when  $h$  does so.



Put  $h = \frac{1}{r}$ , then, by Art. 258,

$$\Gamma(n) = r^{n-1} \int_0^1 (1 - x^r)^{n-1} dx + \int_0^1 y dx.$$

In the first integral put  $x = z^r$ ; thus

$$\Gamma(n) - \int_0^1 y dx = r^n \int_0^1 z^{r-1} (1 - z)^{n-1} dz.$$

We have it in our power to suppose  $r$  an integer; then the integral on the right hand side, by Art. 33, is

$$\frac{1 \cdot 2 \cdot 3 \dots r}{n(n+1) \dots (n+r-1)} r^{n-1}.$$

Let  $r$  increase indefinitely, then  $y$  vanishes and we have

$$\Gamma(n) = \text{limit of } \frac{1 \cdot 2 \cdot 3 \dots r}{n(n+1) \dots (n+r-1)} r^{n-1}.$$

265. From the result of the preceding article we have

$$\frac{\{\Gamma(n)\}^2}{\Gamma(n-m)\Gamma(n+m)} = \left\{1 - \frac{m^2}{n^2}\right\} \left\{1 - \frac{m^2}{(n+1)^2}\right\} \left\{1 - \frac{m^2}{(n+2)^2}\right\} \dots$$

A particular case of this is obtained by supposing  $n = 1$ ; thus

$$\frac{1}{\Gamma(1-m)\Gamma(1+m)} = \left(1 - \frac{m^2}{1^2}\right) \left(1 - \frac{m^2}{2^2}\right) \left(1 - \frac{m^2}{3^2}\right) \dots;$$

the expression on the right hand side is known to be equal to  $\frac{\sin m\pi}{m\pi}$ ; thus

$$\Gamma(1-m)\Gamma(1+m) = \frac{m\pi}{\sin m\pi},$$

therefore  $\Gamma(m)\Gamma(1-m) = \frac{\pi}{\sin m\pi}$  (Art. 259).

266. We shall now establish the following equation,  $n$  being an integer,

$$\Gamma\left(\frac{1}{n}\right) \Gamma\left(\frac{2}{n}\right) \Gamma\left(\frac{3}{n}\right) \dots \Gamma\left(\frac{n-1}{n}\right) = (2\pi)^{\frac{n-1}{2}} n^{-\frac{1}{2}}.$$

First suppose  $n$  odd; in Art. 262 put for  $m$  successively  $\frac{1}{n}, \frac{2}{n}, \frac{3}{n}, \dots$  up to  $\frac{n-1}{2n}$ , and multiply; thus

$$\begin{aligned} \Gamma\left(\frac{1}{n}\right) \Gamma\left(\frac{2}{n}\right) \Gamma\left(\frac{3}{n}\right) \dots \Gamma\left(\frac{n-1}{n}\right) &= \frac{\pi^{\frac{n-1}{2}}}{\sin \frac{\pi}{n} \sin \frac{2\pi}{n} \dots \sin \frac{(n-1)\pi}{2n}} \\ &= (2\pi)^{\frac{n-1}{2}} n^{-\frac{1}{2}}. \end{aligned}$$

(See Hymers's *Theory of Equations*.)

Next suppose  $n$  even; in this case put for  $m$  successively  $\frac{1}{n}, \frac{2}{n}, \dots$  up to  $\frac{n-2}{2n}$ , and form the product as before; then multiply the left hand member by  $\Gamma\left(\frac{1}{2}\right)$  and the right hand member by the equivalent  $\sqrt{\pi}$ ; then we obtain the same result as before.

267. A still more general formula is

$$\begin{aligned} \Gamma(x) \Gamma\left(x + \frac{1}{n}\right) \Gamma\left(x + \frac{2}{n}\right) \dots \Gamma\left(x + \frac{n-1}{n}\right) \\ = \Gamma(nx) (2\pi)^{\frac{n-1}{2}} n^{\frac{1}{2}-nx}, \end{aligned}$$

which we shall now prove. Let  $\phi(x)$  denote

$$\frac{n^{nx} \Gamma(x) \Gamma\left(x + \frac{1}{n}\right) \dots \Gamma\left(x + \frac{n-1}{n}\right)}{n! \Gamma(nx)};$$

we have then to shew that  $\phi(x) = (2\pi)^{\frac{n-1}{2}} n^{-\frac{1}{2}}$ .

We have

$$\begin{aligned}\phi(x+1) &= \frac{n^{nx+n} \Gamma(x+1) \Gamma\left(x+1+\frac{1}{n}\right) \dots \Gamma\left(x+1+\frac{n-1}{n}\right)}{n \Gamma(nx+n)} \\ &= \frac{n^n x \left(x+\frac{1}{n}\right) \left(x+\frac{2}{n}\right) \dots \left(x+\frac{n-1}{n}\right)}{(nx+n-1)(nx+n-2) \dots nx} \phi(x) \\ &= \phi(x).\end{aligned}$$

Similarly  $\phi(x+2) = \phi(x+1) = \phi(x)$ ; and by proceeding thus we have

$$\phi(x) = \phi(x+m),$$

where  $m$  may be as great as we please. Hence  $\phi(x)$  is equal to the limit of  $\phi(\mu)$  when  $\mu$  is infinite; thus  $\phi(x)$  must be independent of  $x$ , that is, must have the same value whatever  $x$  may be; hence  $\phi(x)$  must have the same value as it has when  $x = \frac{1}{n}$ ; thus the theorem follows by the preceding article. This theorem is ascribed to Gauss; a more rigid proof is given in Legendre's *Exercices de Calcul Intégral*, Vol. II. p. 23; see also the *Journal de l'Ecole Polytechnique*, Vol. XVI. p. 212.

268. Many definite integrals may be expressed in terms of the *Gamma-function*; we shall give some examples.

The integral  $\int_0^\infty e^{-a^2 x^2} dx$  becomes by putting  $y$  for  $a^2 x^2$

$$\int_0^\infty \frac{e^{-y} dy}{2a \sqrt{y}}, \text{ that is, } \frac{1}{2a} \Gamma\left(\frac{1}{2}\right), \text{ or } \frac{\sqrt{\pi}}{2a}.$$

Again, in  $\int_0^1 \frac{x^{l-1} (1-x)^{m-1} dx}{(x+a)^{l+m}}$  put  $\frac{x}{x+a} = \frac{y}{1+a}$ ; thus we obtain

$$\frac{1}{a^m (1+a)^l} \int_0^1 y^{l-1} (1-y)^{m-1} dy, \text{ that is } \frac{1}{a^m (1+a)^l} \frac{\Gamma(l) \Gamma(m)}{\Gamma(l+m)}.$$

Again, in  $\int_0^1 x^{l-1} (1-x^2)^{m-1} dx$  put  $x^2 = y$ ; thus we obtain

$$\frac{1}{2} \int_0^1 y^{\frac{l}{2}-1} (1-y)^{m-1} dy, \text{ that is } \frac{\Gamma\left(\frac{l}{2}\right) \Gamma(m)}{2\Gamma\left(\frac{l}{2}+m\right)}.$$

Thus 
$$\int_0^{\frac{\pi}{2}} \sin^p \theta \cos^q \theta d\theta = \int_0^1 x^p (1-x^2)^{\frac{q-1}{2}} dx$$

$$= \int_0^1 x^{p+1-1} (1-x^2)^{\frac{q+1}{2}-1} dx = \frac{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{q+1}{2}\right)}{2\Gamma\left(\frac{p+q}{2}+1\right)}.$$

Again, in  $\int_0^1 \frac{x^{l-1} (1-x)^{m-1} dx}{\{ax+b(1-x)\}^{l+m}}$  put  $x = \frac{by}{a(1-y)+by}$ ; thus we obtain

$$\frac{1}{a^l b^m} \int_0^1 y^{l-1} (1-y)^{m-1} dy, \text{ that is, } \frac{\Gamma(l) \Gamma(m)}{a^l b^m \Gamma(l+m)}.$$

269. In  $\int_0^a x^{l-1} (a-x)^{m-1} dx$  put  $x = ay$ ; thus we obtain

$$a^{l+m-1} \int_0^1 y^{l-1} (1-y)^{m-1} dy, \text{ that is, } a^{l+m-1} \frac{\Gamma(l) \Gamma(m)}{\Gamma(l+m)}.$$

270. It is required to find the value of the multiple integral

$$\iiint \dots x^{l-1} y^{m-1} z^{n-1} \dots dx dy dz \dots$$

the integral being so taken as to give to the variables all *positive* values consistent with the condition that  $x+y+z+\dots$  is not greater than unity.

We will suppose that there are three variables, and consequently that the integral is a triple integral; the method

adopted will be seen to be applicable for any number of variables.

We must first integrate for one of the variables, suppose  $z$ ; the limits then will be 0 and  $1 - x - y$ ; thus between these limits

$$\int z^{n-1} dz = \frac{(1-x-y)^n}{n} = \frac{\Gamma(n)}{\Gamma(n+1)} (1-x-y)^n.$$

Next integrate with respect to one of the remaining variables, suppose  $y$ ; the limits will be 0 and  $1 - x$ ; and between these limits, by Art. 269,

$$\int y^{m-1} (1-x-y)^n dy = \frac{(1-x)^{m+n} \Gamma(m) \Gamma(n+1)}{\Gamma(m+n+1)}.$$

Lastly integrate with respect to  $x$  between the limits 0 and 1; thus between these limits

$$\int x^{l-1} (1-x)^{m+n} dx = \frac{\Gamma(l) \Gamma(m+n+1)}{\Gamma(l+m+n+1)}.$$

Hence the final result is

$$\frac{\Gamma(n)}{\Gamma(n+1)} \frac{\Gamma(m) \Gamma(n+1)}{\Gamma(m+n+1)} \frac{\Gamma(l) \Gamma(m+n+1)}{\Gamma(l+m+n+1)},$$

that is, 
$$\frac{\Gamma(l) \Gamma(m) \Gamma(n)}{\Gamma(l+m+n+1)}.$$

271. It is required to find the value of the multiple integral

$$\iiint \dots \xi^{l-1} \eta^{m-1} \zeta^{n-1} \dots d\xi d\eta d\zeta \dots$$

the integral being so taken as to give to the variables all positive values consistent with the condition that

$$\left(\frac{\xi}{\alpha}\right)^p + \left(\frac{\eta}{\beta}\right)^q + \left(\frac{\zeta}{\gamma}\right)^r + \dots$$

is not greater than unity.

Assume  $x = \left(\frac{\xi}{\alpha}\right)^p$ ,  $y = \left(\frac{\eta}{\beta}\right)^q$ ,  $z = \left(\frac{\zeta}{\gamma}\right)^r, \dots$

Then the integral becomes

$$\frac{\alpha^l \beta^m \gamma^n \dots}{p q r \dots} \iiint \dots x^{\frac{l}{p}-1} y^{\frac{m}{q}-1} z^{\frac{n}{r}-1} \dots dx dy dz \dots$$

with the condition that  $x + y + z + \dots$  is not greater than unity. The value of the integral is, therefore, by the preceding article

$$\frac{\alpha^l \beta^m \gamma^n \dots}{p q r \dots} \frac{\Gamma\left(\frac{l}{p}\right) \Gamma\left(\frac{m}{q}\right) \Gamma\left(\frac{n}{r}\right) \dots}{\Gamma\left(\frac{l}{p} + \frac{m}{q} + \frac{n}{r} + \dots + 1\right)}.$$

272. As a simple case of the preceding article we may suppose  $p, q, r, \dots$  to be each unity, and  $\alpha, \beta, \gamma, \dots$  each equal to a constant  $h$ ; thus the condition is that  $\xi + \eta + \zeta + \dots$  is not to be greater than  $h$ . The value of the integral

$$\iiint \dots \xi^{l-1} \eta^{m-1} \zeta^{n-1} \dots d\xi d\eta d\zeta \dots$$

then is  $h^{l+m+n+\dots} \frac{\Gamma(l) \Gamma(m) \Gamma(n) \dots}{\Gamma(l+m+n+\dots+1)}$ ,

which we may denote by

$$N h^{l+m+n+\dots}.$$

Similarly if the integral is to be taken so that the sum of the variables shall not exceed  $h + \Delta h$ , we obtain for the result

$$N (h + \Delta h)^{l+m+n+\dots}.$$

Hence we conclude that the value of the integral extended over all such positive values of the variables as make the sum of the variables lie between  $h$  and  $h + \Delta h$  is

$$N \{(h + \Delta h)^{l+m+n+\dots} - h^{l+m+n+\dots}\},$$

and when  $\Delta h$  is indefinitely diminished, this becomes

$$N(l+m+n+\dots) h^{l+m+n+\dots-1} \Delta h,$$

that is 
$$\frac{\Gamma(l)\Gamma(m)\Gamma(n)\dots}{\Gamma(l+m+n+\dots)} h^{l+m+n+\dots-1} \Delta h.$$

273. It is required to find the value of the multiple integral

$$\iiint \dots x^{l-1} y^{m-1} z^{n-1} \dots f(x+y+z+\dots) dx dy dz \dots$$

the integral being so taken as to give to the variables all *positive* values consistent with the condition that  $x+y+z+\dots$  is not greater than  $c$ .

We will suppose for simplicity that there are three variables. By the preceding article, that part of the integral which arises from supposing the sum of the variables to lie between  $h$  and  $h+\Delta h$  is

$$\frac{\Gamma(l)\Gamma(m)\Gamma(n)}{\Gamma(l+m+n)} f(h) h^{l+m+n-1} \Delta h.$$

Hence the whole integral is

$$\frac{\Gamma(l)\Gamma(m)\Gamma(n)}{\Gamma(l+m+n)} \int_0^c f(h) h^{l+m+n-1} dh.$$

274. Similarly the value of

$$\iiint \xi^{l-1} \eta^{m-1} \zeta^{n-1} f\left\{\left(\frac{\xi}{\alpha}\right)^p + \left(\frac{\eta}{\beta}\right)^q + \left(\frac{\zeta}{\gamma}\right)^r\right\} d\xi d\eta d\zeta$$

for all positive values of the variables, such that

$$\left(\frac{\xi}{\alpha}\right)^p + \left(\frac{\eta}{\beta}\right)^q + \left(\frac{\zeta}{\gamma}\right)^r$$

is not greater than  $c$ , is

$$\frac{\alpha^l \beta^m \gamma^n}{p q r} \frac{\Gamma\left(\frac{l}{p}\right) \Gamma\left(\frac{m}{q}\right) \Gamma\left(\frac{n}{r}\right)}{\Gamma\left(\frac{l}{p} + \frac{m}{q} + \frac{n}{r}\right)} \int_0^1 f(h) h^{\frac{l}{p} + \frac{m}{q} + \frac{n}{r} - 1} dh.$$

The result of this and the preceding article may be extended to the case of any number of variables.

275. It is required to find the value of the multiple integral

$$\iiint \dots f(a_1 x_1 + a_2 x_2 + \dots + a_n x_n) dx_1 dx_2 \dots dx_n,$$

the integral being so taken as to give to the variables *all* values consistent with the condition that  $x_1^2 + x_2^2 + \dots + x_n^2$  is not greater than unity.

By successive application of a transformation for a double integral given in Art. 242, the multiple integral may be reduced to

$$\iiint \dots f(kx_1) dx_1 dx_2 \dots dx_n,$$

where  $k = \sqrt{(a_1^2 + a_2^2 + \dots + a_n^2)}$ .

We have first then to find the value of the multiple integral  $\iiint \dots dx_2 dx_3 \dots dx_n$ , the variables being supposed to have all values consistent with the condition that  $x_2^2 + x_3^2 + \dots + x_n^2$  is not greater than  $1 - x_1^2$ . First suppose that the variables are to have only *positive* values; then we obtain the value of the integral by supposing in Art. 271, that each of the quantities  $l, m, \dots$  is unity, that each of the quantities  $p, q, \dots$  is equal to 2, and that each of the quantities  $\alpha, \beta, \dots$  is equal to  $\sqrt{(1 - x_1^2)}$ . Thus the result is

$$\frac{\left\{\Gamma\left(\frac{1}{2}\right)\right\}^{n-1}}{2^{n-1} \Gamma\left(\frac{n-1}{2} + 1\right)} (1 - x_1^2)^{\frac{n-1}{2}}.$$



But if the variables may have negative as well as positive values, this result must be multiplied by  $2^{n-1}$ . Thus we get

$$\frac{\pi^{\frac{n-1}{2}} (1-x_1^2)^{\frac{n-1}{2}}}{\Gamma\left(\frac{n-1}{2} + 1\right)}.$$

Hence, finally, since the limits of  $x_1$  will be  $-1$  and  $1$ , the multiple integral is equal to

$$\frac{\pi^{\frac{n-1}{2}}}{\Gamma\left(\frac{n-1}{2} + 1\right)} \int_{-1}^1 f(kx_1) (1-x_1^2)^{\frac{n-1}{2}} dx_1.$$

This agrees with the result given by Professor Boole in the *Cambridge Mathematical Journal*, Vol. III. p. 280, as it may be found by integrating his equation (15) by parts.

276. It is required to find the value of the multiple integral

$$\iiint \dots \frac{f(a_1x_1 + a_2x_2 + \dots + a_nx_n)}{\sqrt{(1-x_1^2-x_2^2-\dots-x_n^2)}} dx_1 dx_2 \dots dx_n,$$

the integral being so taken as to give to the variables *all* values consistent with the condition that  $x_1^2 + x_2^2 + \dots + x_n^2$  is not greater than unity.

As in the preceding article the integral may be transformed into

$$\iiint \dots \frac{f(kx_1)}{\sqrt{(1-x_1^2-x_2^2-\dots-x_n^2)}} dx_1 dx_2 \dots dx_n.$$

First integrate with respect to the variables  $x_2, x_3, \dots, x_n$ , the limits being given by the condition that  $x_2^2 + x_3^2 + \dots + x_n^2$  is not greater than  $1-x_1^2$ . Now if the variables were restricted to positive values, the integral

$$\iiint \dots \frac{dx_2 dx_3 \dots dx_n}{\sqrt{(1-x_1^2-x_2^2-\dots-x_n^2)}}$$

by Art. 274 would be equal to

$$\frac{1}{2^{n-1}} \frac{\{\Gamma(\frac{1}{2})\}^{n-1}}{\Gamma(\frac{n-1}{2})} \int_0^{1-x_1^2} (1-x_1^2-h)^{-\frac{1}{2}} h^{\frac{n-1}{2}-1} dh,$$

that is, to

$$\frac{1}{2^{n-1}} \frac{\{\Gamma(\frac{1}{2})\}^{n-1}}{\Gamma(\frac{n-1}{2})} (1-x_1^2)^{\frac{n}{2}-1} \frac{\Gamma(\frac{1}{2}) \Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})}, \quad (\text{Art. 269}),$$

that is, to 
$$\frac{1}{2^{n-1}} \frac{\{\Gamma(\frac{1}{2})\}^n}{\Gamma(\frac{n}{2})} (1-x_1^2)^{\frac{n}{2}-1}.$$

But if the variables may have negative as well as positive values, this result must be multiplied by  $2^{n-1}$ . Thus we get

$$\frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} (1-x_1^2)^{\frac{n}{2}-1}.$$

Hence finally, since the limits of  $x_1$  are  $-1$  and  $1$ , the multiple integral is equal to

$$\frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_{-1}^1 f(kx_1) (1-x_1^2)^{\frac{n}{2}-1} dx_1.$$

277. Many methods have been used for exhibiting in simple terms an approximate value of  $\Gamma(n+1)$  when  $n$  is very large: we give one of them.

The product  $e^{-x} x^n$  vanishes when  $x=0$  and when  $x=\infty$ ; and it may be shewn that it has only one maximum value, namely when  $x=n$ . We may therefore assume

$$e^{-x} x^n = e^{-n} n^n e^{-t^2} \dots\dots\dots(1),$$

where  $t$  is a variable which must lie between the limits  $-\infty$  and  $+\infty$ .

Thus 
$$\int_0^\infty e^{-x} x^n dx = e^{-n} n^n \int_{-\infty}^\infty e^{-t^2} \frac{dx}{dt} dt \dots\dots\dots(2).$$

Take the logarithms of both members of (1); thus

$$x - n \log x = n - n \log n + t^2 \dots\dots\dots(3);$$

put  $x = n + u$ ; thus

$$u - n \log (n + u) = t^2 - n \log n \dots\dots\dots(4).$$

But by Taylor's Theorem

$$\log (n + u) = \log n + \frac{u}{n} - \frac{u^2}{2 (n + \theta u)^2},$$

where  $\theta$  is a proper fraction; thus (4) becomes

$$\frac{nu^2}{2 (n + \theta u)^2} = t^2;$$

therefore

$$\frac{\sqrt{(n)} u}{\sqrt{(2)} (n + \theta u)} = t \dots\dots\dots(5);$$

therefore

$$u = \frac{\sqrt{(2)} nt}{\sqrt{(n)} - \theta t \sqrt{2}} \dots\dots\dots(6).$$

But from (3)

$$\begin{aligned} \frac{dx}{dt} &= \frac{2xt}{x-n} = 2t + \frac{2nt}{u} \\ &= \sqrt{(2n)} + 2 (1 - \theta) t, \end{aligned} \quad \text{by (6).}$$

Hence (2) becomes

$$\int_0^\infty e^{-x} x^n dx = e^{-n} n^n \int_{-\infty}^\infty e^{-t^2} \{ \sqrt{(2n)} + 2 (1 - \theta) t \} dt;$$

and  $\int_{-\infty}^\infty e^{-t^2} dt = \sqrt{(\pi)}$ ; thus

$$\int_0^\infty e^{-x} x^n dx = e^{-n} n^n \sqrt{(2n\pi)} \left\{ 1 + \frac{2}{\sqrt{(2n\pi)}} \int_{-\infty}^\infty e^{-t^2} (1 - \theta) t dt \right\} (7).$$

But since  $1 - \theta$  is positive and less than unity, the numerical value of  $\int_{-\infty}^\infty e^{-t^2} (1 - \theta) t dt$  is less than  $\int_0^\infty e^{-t^2} t dt$ , that is, less than  $\frac{1}{2}$ . Hence we conclude from (7) that as  $n$  is increased indefinitely, the ratio of  $\Gamma(n + 1)$  to  $e^{-n} n^n \sqrt{(2n\pi)}$  approaches unity as its limit.

We may observe that the right sign is taken in (5), for if  $u$  be positive  $t$  must be positive; and  $\frac{dx}{dt}$  is *always* to be positive, and the value of  $\frac{dx}{dt}$  would be negative when  $t$  is negative if we used the other sign.

(See Liouville's *Journal de Mathématiques*, Vol. x. p. 464, and Vol. xvii. p. 448).

*Definite Integrals obtained by differentiating or integrating with respect to constants.*

278. We shall now give some examples in which definite integrals are obtained by means of differentiation with respect to a constant. (See Art. 213.)

To find the value of  $\int_0^{\infty} e^{-a^2x^2} \cos 2rx \, dx$ . Call the definite integral  $u$ ; then

$$\frac{du}{dr} = -2 \int_0^{\infty} xe^{-a^2x^2} \sin 2rx \, dx.$$

Integrate the right-hand term by parts; thus we find

$$\frac{du}{dr} = -\frac{2ru}{a^2};$$

therefore 
$$\frac{d \log u}{dr} = -\frac{2r}{a^2};$$

therefore 
$$\log u = -\frac{r^2}{a^2} + \text{constant},$$

therefore 
$$u = Ae^{-\frac{r^2}{a^2}},$$

where  $A$  is a quantity which is constant with respect to  $r$ , that is, it does not contain  $r$ . To determine  $A$  we may suppose  $r = 0$ ; thus  $u$  becomes  $\int_0^{\infty} e^{-a^2x^2} \, dx$ , that is,  $\frac{\sqrt{\pi}}{2a}$ , (Art. 268).

Hence  $A = \frac{\sqrt{\pi}}{2a}$ , and

$$\int_0^{\infty} e^{-a^2x^2} \cos 2rx \, dx = \frac{\sqrt{\pi}}{2a} e^{-\frac{r^2}{a^2}}.$$

279. We have stated in Art. 214, that when one of the limits of integration is *infinite* the process of differentiation with respect to a constant *may* be unsafe; in the present case however it is easy to justify it; we have to shew that  $\int_0^{\infty} e^{-a^2x^2} \rho \, dx$  vanishes where  $\rho$  is ultimately indefinitely small; it is obvious that this quantity is numerically less than  $\rho_1 \int_0^{\infty} e^{-a^2x^2} \, dx$  where  $\rho_1$  is the greatest value of  $\rho$ , that is less than  $\frac{\sqrt{\pi}}{2a} \rho_1$ ; but this vanishes since  $\rho_1$  does. Similar considerations apply to the succeeding cases.

280. To find the value of  $\int_0^{\infty} e^{-kx} \frac{\sin rx \, dx}{x}$ . Denote it by  $u$ , then

$$\frac{du}{dr} = \int_0^{\infty} e^{-kx} \cos rx \, dx.$$

But  $\int_0^{\infty} e^{-kx} \cos rx \, dx = e^{-kx} \frac{r \sin rx - k \cos rx}{k^2 + r^2}$ ;

therefore  $\int_0^{\infty} e^{-kx} \cos rx \, dx = \frac{k}{k^2 + r^2}$ ;

thus  $\frac{du}{dr} = \frac{k}{k^2 + r^2}$ ;

therefore  $u = \tan^{-1} \frac{r}{k}$ .

No constant is required because  $u$  vanishes with  $r$ . This result holds for any positive value of  $k$ ; if we suppose  $k$  to diminish without limit, we obtain

$$\int_0^{\infty} \frac{\sin rx}{x} \, dx = \frac{\pi}{2}$$

if  $r$  be positive; if  $r$  be negative the result should be  $-\frac{\pi}{2}$ .

281. To find the value of  $\int_0^{\infty} e^{-(x^2 + \frac{a^2}{x^2})} dx$ . Denote it by  $u$ , then

$$\frac{du}{da} = -2a \int_0^{\infty} e^{-(x^2 + \frac{a^2}{x^2})} \frac{dx}{x^3};$$

assume  $x = \frac{a}{z}$ , then the limits of  $z$  are  $\infty$  and 0; and we obtain

$$\frac{du}{da} = -2u;$$

therefore  $\frac{d \log u}{da} = -2;$

therefore  $\log u = -2a + \text{constant};$

therefore  $u = Ae^{-2a}.$

To determine  $A$  we may suppose  $a = 0$ ; then  $u = \frac{\sqrt{\pi}}{2};$

therefore  $A = \frac{\sqrt{\pi}}{2};$  thus

$$\int_0^{\infty} e^{-(x^2 + \frac{a^2}{x^2})} dx = \frac{\sqrt{\pi}}{2} e^{-2a}.$$

282. We may also apply the principle of *integration* with respect to a constant in order to determine some definite integrals; the principle may be established thus.

Let  $u = \int_a^b \phi(x, c) dx,$

then  $\int_a^{\beta} u dc = \int_a^{\beta} \int_a^b \phi(x, c) dc dx$   
 $= \int_a^b \int_a^{\beta} \phi(x, c) dx dc;$

since when the limits are constant, the order of integration is indifferent, (Art. 62). We shall now give some examples of this method.

283. We know that  $\int_0^{\infty} e^{-kx} dx = \frac{1}{k}$ .

Integrate both sides with respect to  $k$  between the limits  $a$  and  $b$ ; thus

$$\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} dx = \log \frac{b}{a}.$$

It should be noticed that  $\int_0^{\infty} \frac{e^{-ax} dx}{x}$  and  $\int_0^{\infty} \frac{e^{-bx} dx}{x}$  are both infinite; for  $\int_0^c \frac{e^{-ax} dx}{x}$  is greater than  $e^{-ca} \int_0^c \frac{dx}{x}$ , and  $\int_0^c \frac{dx}{x}$  is infinite. But this is not inconsistent with the assertion that  $\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} dx$  is finite, and without finding the value of this integral it is easy to shew that it must be finite. For it is equal to the sum of  $\int_0^c \frac{\phi(x) dx}{x}$  and  $\int_c^{\infty} \frac{\phi(x) dx}{x}$  where  $\phi(x) = e^{-ax} - e^{-bx}$ ; the second of these integrals is finite, for it is less than  $\frac{1}{c} \int_c^{\infty} \phi(x) dx$ , that is, less than  $\frac{1}{c} \left( \frac{e^{-ac}}{a} - \frac{e^{-bc}}{b} \right)$ .

We have then only to examine  $\int_0^c \frac{\phi(x)}{x} dx$ .

Now by Maclaurin's Theorem

$$\phi(x) = (b-a)x + \frac{x^2}{2} \phi''(x\theta),$$

where  $\theta$  is some fraction; thus  $\frac{\phi(x)}{x}$  is less than  $b-a + \frac{Ax}{2}$ , where  $A$  is the greatest value which  $\phi''(x)$  can assume for values of  $x$  less than  $c$ . Hence

$$\int_0^c \frac{\phi(x)}{x} dx \text{ is less than } (b-a)c + \frac{Ac^2}{4},$$

and is therefore finite.

284. We know that

$$\int_0^{\infty} e^{-kx} \cos rx dx = \frac{k}{k^2 + r^2}.$$

Integrate both sides with respect to  $k$  between the limits  $a$  and  $b$ ; thus

$$\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} \cos rx \, dx = \frac{1}{2} \log \frac{b^2 + r^2}{a^2 + r^2}.$$

285. Let  $\int_0^{\infty} \frac{\sin rx}{x} dx$  be denoted by  $A$ , and  $\int_0^{\infty} \frac{\cos rx}{1+x^2} dx$  by  $B$ ; we shall now determine the values of  $A$  and  $B$ ; the former has already been determined by another method in Art. 280.

In the integral  $A$  put  $y$  for  $rx$ ; thus

$$A = \int_0^{\infty} \frac{\sin y \, dy}{y};$$

this shews that  $A$  is independent of  $r$ .

We have 
$$\frac{dB}{dr} = - \int_0^{\infty} \frac{x \sin rx \, dx}{1+x^2},$$

and 
$$\int_0^r B dr = \int_0^{\infty} \frac{\sin rx \, dx}{x(1+x^2)};$$

thus 
$$\int_0^r B dr - \frac{dB}{dr} = \int_0^{\infty} \frac{1+x^2}{x} \frac{\sin rx}{1+x^2} dx = A;$$

hence 
$$\int_0^r B dr - \frac{dB}{dr} - A = 0 \dots\dots\dots (1).$$

Multiply by  $e^{-r}$  and integrate; we obtain since  $A$  is constant with respect to  $r$

$$e^{-r} \left\{ \int_0^r B dr + B - A \right\} = \text{constant}.$$

Now whatever be the value of  $r$ , it is obvious that the integrals represented by  $A$ ,  $B$ , and  $\int_0^r B dr$  are *finite*; hence the constant in the last equation must be zero, for the left-hand member vanishes when  $r$  is infinite.



Thus 
$$\int_0^r Bdr + B - A = 0 \dots \dots \dots (2).$$

From (1) and (2) 
$$\frac{dB}{dr} = -B;$$

therefore 
$$B = Ce^{-r},$$

where  $C$  is some constant. And from (2)

$$A = Ce^{-r} - C(e^{-r} - 1) = C;$$

therefore 
$$B = Ae^{-r} \dots \dots \dots (3).$$

Now when  $r$  is indefinitely diminished,  $B$  becomes  $\int_0^\infty \frac{dx}{1+x^2}$ , that is  $\frac{\pi}{2}$ ; hence from (3)

$$A = \frac{\pi}{2} \text{ and } B = \frac{\pi}{2} e^{-r}.$$

We have supposed  $r$  positive; it is obvious that if  $r$  be negative,  $B$  has the same value as if  $r$  were positive, and  $A$  has its sign changed; that is, if  $r$  be negative  $B = \frac{\pi}{2} e^r$  and  $A = -\frac{\pi}{2}$ . (*Transactions of the Royal Irish Academy*, Vol. XIX. p. 277.)

286. From  $\int_0^\infty \frac{\cos rx dx}{1+x^2} = \frac{\pi}{2} e^{-r}$ , we obtain by differentiation with respect to  $r$ ,

$$\int_0^\infty \frac{x \sin rx dx}{1+x^2} = \frac{\pi}{2} e^{-r}.$$

And from the same integral by integrating with respect to  $r$  between the limits 0 and  $c$ , we have

$$\int_0^\infty \frac{\sin cx dx}{x(1+x^2)} = \frac{\pi}{2} (1 - e^{-c}).$$

*Definite Integrals obtained by Expansion.*

287. If we expand  $\log \{1 - ae^{x\sqrt{(-1)}}\}$  and  $\log \{1 - ae^{-x\sqrt{(-1)}}\}$  and add, we obtain

$$\log (1 - 2a \cos x + a^2) \\ = -2 \left( a \cos x + \frac{a^2}{2} \cos 2x + \frac{a^3}{3} \cos 3x + \dots \right),$$

the series being convergent if  $a$  is less than unity. Integrate both sides with respect to  $x$  between the limits 0 and  $\pi$ ; thus

$$\int_0^\pi \log (1 - 2a \cos x + a^2) dx = 0, \quad a \text{ being less than 1.}$$

If  $a$  is greater than 1, since

$$\log (1 - 2a \cos x + a^2) = \log a^2 + \log \left( 1 - \frac{2}{a} \cos x + \frac{1}{a^2} \right),$$

we have

$$\int_0^\pi \log (1 - 2a \cos x + a^2) dx = \pi \log a^2 = 2\pi \log a.$$

288. By integration by parts we have

$$\int \log (1 - 2a \cos x + a^2) dx \\ = x \log (1 - 2a \cos x + a^2) - 2a \int \frac{x \sin x dx}{1 - 2a \cos x + a^2}.$$

Hence, if  $a$  be less than 1,

$$\int_0^\pi \frac{x \sin x dx}{1 - 2a \cos x + a^2} = \frac{\pi}{2a} \log (1 + a)^2, \quad \text{that is, } \frac{\pi}{a} \log (1 + a);$$

if  $a$  be greater than 1, the result is

$$\frac{\pi}{a} \log (1 + a) - \frac{\pi}{a} \log a, \quad \text{that is, } \frac{\pi}{a} \log \left( 1 + \frac{1}{a} \right).$$

289. In like manner we have, if  $r$  be an integer

$$\int_0^\pi \cos rx \log (1 - 2a \cos x + a^2) dx = -\frac{\pi}{r} a^r, \quad \text{or } -\frac{\pi}{r} a^{-r},$$

according as  $a$  is less or greater than unity.

290. Integrate by parts the integral in the preceding article; thus we find

$$\int_0^{\pi} \frac{\sin x \sin rx \, dx}{1 - 2a \cos x + a^2} = \frac{\pi}{2} a^{r-1} \text{ or } \frac{\pi}{2} a^{-(r+1)},$$

according as  $a$  is less or greater than unity.

291. Similarly from the known expansion

$$\begin{aligned} \frac{1 - a^2}{1 - 2a \cos x + a^2} \\ = 1 + 2a \cos x + 2a^2 \cos 2x + 2a^3 \cos 3x + \dots, \end{aligned}$$

where  $a$  is less than 1, we may deduce some definite integrals; thus if  $r$  is an integer

$$\int_0^{\pi} \frac{\cos rx \, dx}{1 - 2a \cos x + a^2} = \frac{\pi a^r}{1 - a^2},$$

for every term that we have to integrate vanishes with the assigned limits, except  $2a^r \int_0^{\pi} \cos^2 rx \, dx$ .

292. To find the value of  $\int_0^{\infty} \frac{1}{1+x^2} \frac{dx}{1-2a \cos cx + a^2}$ .

The term  $\frac{1}{1-2a \cos cx + a^2}$  may be expanded as in Art. 291; then each term may be integrated by Art. 286, and the results summed. Thus we shall obtain

$$\frac{\pi}{2} \cdot \frac{1}{1-a^2} \frac{1+ae^{-c}}{1-ae^{-c}}.$$

293. Similarly,

$$\int_0^{\infty} \log(1 - 2a \cos cx + a^2) \frac{dx}{1+x^2} = \pi \log(1 - ae^{-c}).$$

294. It is also known from Trigonometry that

$$\frac{\sin cx}{1 - 2a \cos cx + a^2} = \sin cx + a \sin 2cx + a^2 \sin 3cx + \dots,$$

$a$  being less than 1. Hence by Art. 286, we obtain

$$\int_0^\infty \frac{x \sin cx \, dx}{(1 + x^2)(1 - 2a \cos cx + a^2)} = \frac{\pi}{2(e^c - a)}.$$

This also follows from Art. 293, by differentiating with respect to  $c$ .

295. To find  $\int_0^1 \frac{\log x}{1-x} dx$ .

By expanding  $(1-x)^{-1}$ , we find for the integral a series of which the type is

$$\int_0^1 x^n \log x \, dx.$$

By integration by parts this is seen to be equal to  $-\frac{1}{(1+n)^2}$ . Hence the result is

$$-\left\{1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots\right\},$$

that is, by a known formula,  $-\frac{\pi^2}{6}$ .

296. To find  $\int_0^\pi \frac{x \sin x \, dx}{1 + (\cos x)^2}$ .

Expand the factor  $\{1 + (\cos x)^2\}^{-1}$ , and we find for the integral a series of which the type is

$$(-1)^n \int_0^\pi x \sin x (\cos x)^{2n} \, dx.$$

By integration by parts this may be shewn to be equal to  $\frac{(-1)^n \pi}{2n+1}$ .

Hence the result is

$$\pi \left\{ 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots \right\},$$

that is, by a known formula,  $\frac{\pi^2}{4}$ .

297. Let  $v$  denote  $e^{x\sqrt{-1}}$ , that is,  $\cos x + \sqrt{-1} \sin x$ ; then if  $f$  denote any function, we have by Taylor's Theorem,

$$\begin{aligned} f(a+v) + f(a+v^{-1}) \\ = 2 \left\{ f(a) + f'(a) \cos x + \frac{f''(a)}{1.2} \cos 2x + \dots \right\}. \end{aligned}$$

And

$$\frac{1-c^2}{1-2c \cos x + c^2} = 1 + 2c \cos x + 2c^2 \cos 2x + 2c^3 \cos 3x + \dots$$

Therefore

$$\begin{aligned} \int_0^\pi \frac{f(a+v) + f(a+v^{-1})}{1-2c \cos x + c^2} dx &= \frac{2\pi}{1-c^2} \left\{ f(a) + cf'(a) + \frac{c^2}{1.2} f''(a) + \dots \right\} \\ &= \frac{2\pi}{1-c^2} f(a+c). \end{aligned}$$

In this result it must be remembered that  $c$  is to be less than unity, and the functions  $f(a+v)$  and  $f(a+v^{-1})$  must be such that Taylor's Theorem holds for their expansions.

In a similar way it may be shewn that

$$\int_0^\pi \frac{f(a+v) - f(a+v^{-1})}{1-2c \cos x + c^2} \sin x dx = \frac{\pi \sqrt{-1}}{c} \{f(a+c) - f(a)\}$$

$$\begin{aligned} \text{and } \int_0^\pi \frac{1-c \cos x}{1-2c \cos x + c^2} \{f(a+v) + f(a+v^{-1})\} dx \\ = \pi \{f(a+c) + f(a)\}. \end{aligned}$$

*Substitution of imaginary values for Constants.*

298. Definite integrals are sometimes deduced from known integrals by substituting impossible values for some of the constants which occur. This process cannot be considered demonstrative, but will serve at least to suggest the forms which can be examined, and perhaps verified by other methods, (see De Morgan's *Differential and Integral Calculus*, p. 630). We will give some examples of it.

We have 
$$\int_0^{\infty} e^{-px} x^{n-1} dx = p^{-n} \Gamma(n).$$

For  $p$  put  $a + b\sqrt{-1}$ , and suppose  $r = \sqrt{a^2 + b^2}$  and  $\tan \theta = \frac{b}{a}$ , so that  $p = r \{ \cos \theta + \sqrt{-1} \sin \theta \}$ ; thus

$$\int_0^{\infty} e^{-\{a+b\sqrt{-1}\}x} x^{n-1} dx = r^{-n} \{ \cos n\theta - \sqrt{-1} \sin n\theta \} \Gamma(n).$$

Thus by separating the possible and impossible parts we have

$$\int_0^{\infty} e^{-ax} x^{n-1} \cos bx dx = \frac{\Gamma(n) \cos \left( n \tan^{-1} \frac{b}{a} \right)}{(a^2 + b^2)^{\frac{n}{2}}}.$$

$$\int_0^{\infty} e^{-ax} x^{n-1} \sin bx dx = \frac{\Gamma(n) \sin \left( n \tan^{-1} \frac{b}{a} \right)}{(a^2 + b^2)^{\frac{n}{2}}}.$$

For modes of verification see De Morgan, p. 630.

299. In the formula

$$\int_0^{\infty} e^{-a^2x^2} dx = \frac{\sqrt{\pi}}{2a}$$

change  $a$  into  $\frac{1 + \sqrt{-1}}{\sqrt{2}} c$ ; thus

$$\int_0^{\infty} e^{-c^2x^2\sqrt{-1}} dx = \frac{1 - \sqrt{-1}}{2c} \frac{\sqrt{\pi}}{\sqrt{2}};$$

therefore

$$\int_0^{\infty} (\cos c^2 x^2 - \sqrt{-1} \sin c^2 x^2) dx = \frac{1 - \sqrt{-1}}{2c} \frac{\sqrt{\pi}}{\sqrt{2}};$$

therefore 
$$\int_0^{\infty} \cos c^2 x^2 dx = \frac{\sqrt{\pi}}{2c\sqrt{2}},$$

and 
$$\int_0^{\infty} \sin c^2 x^2 dx = \frac{\sqrt{\pi}}{2c\sqrt{2}}.$$

If we write  $y$  for  $c^2 x^2$ , these become

$$\int_0^{\infty} \frac{\sin y dy}{\sqrt{y}} = \int_0^{\infty} \frac{\cos y dy}{\sqrt{y}} = \sqrt{\frac{\pi}{2}}.$$

300. In the integral  $\int_0^{\infty} e^{-(x^2 + \frac{a^2}{x^2})k} dx$ , suppose  $y = x\sqrt{k}$ ; thus the integral becomes  $\frac{1}{\sqrt{k}} \int_0^{\infty} e^{-(y^2 + \frac{k^2 a^2}{y^2})} dy$ , which is known by Art. 281. Thus

$$\int_0^{\infty} e^{-(x^2 + \frac{a^2}{x^2})k} dx = \frac{1}{\sqrt{k}} \frac{\sqrt{\pi}}{2} e^{-2ak}.$$

Now put  $\cos \theta + \sqrt{-1} \sin \theta$  for  $k$ ; thus the right-hand member becomes

$$\frac{1}{\cos \frac{\theta}{2} + \sqrt{-1} \sin \frac{\theta}{2}} \cdot \frac{\sqrt{\pi}}{2} e^{-2a\{\cos \theta + \sqrt{-1} \sin \theta\}},$$

that is,

$$\frac{\sqrt{\pi}}{2} \left\{ \cos \left( 2a \sin \theta + \frac{\theta}{2} \right) - \sqrt{-1} \sin \left( 2a \sin \theta + \frac{\theta}{2} \right) \right\} e^{-2a \cos \theta}.$$

Thus 
$$\int_0^{\infty} e^{-(x^2 + \frac{a^2}{x^2}) \cos \theta} \cos \left\{ \left( x^2 + \frac{a^2}{x^2} \right) \sin \theta \right\} dx$$

$$= \frac{\sqrt{\pi}}{2} e^{-2a \cos \theta} \cos \left( 2a \sin \theta + \frac{\theta}{2} \right),$$

and 
$$\int_0^{\infty} e^{-(x^2 + \frac{a^2}{x^2}) \cos \theta} \sin \left\{ \left( x^2 + \frac{a^2}{x^2} \right) \sin \theta \right\} dx$$

$$= \frac{\sqrt{\pi}}{2} e^{-2a \cos \theta} \sin \left( 2a \sin \theta + \frac{\theta}{2} \right).$$

EXAMPLES.

1. Evaluate  $\int_0^{\infty} \frac{(x^2 + a^2) dx}{x^4 + b^2 x^2 + b^4}$ . *Result.*  $\frac{(a^2 + b^2) \pi}{2b^3 \sqrt{3}}$

2. Evaluate  $\int_0^{\frac{1}{2}\pi} \cos(a \tan x) dx$ . *Result.*  $\frac{\pi}{2} e^{-a}$

3. Evaluate  $\int_0^1 x^{2n-1} e^{ax} dx$ . *Result.*  $\frac{1}{n}$

4.  $\int_0^{\frac{\pi}{2}} \frac{dx}{(a^2 \cos^2 x + b^2 \sin^2 x)^2} = \frac{\pi}{4} \left( \frac{1}{ab^3} + \frac{1}{a^3 b} \right)$ .

5. Prove  $\int_0^{\frac{\pi}{4}} \sqrt{\tan \phi} d\phi = \frac{1}{\sqrt{2}} \left[ \frac{\pi}{2} + \log \{ \sqrt{2} - 1 \} \right]$ .

6. Prove  $\int_0^{\frac{\pi}{4}} \sqrt{\cot \phi} d\phi = \frac{1}{\sqrt{2}} \left[ \frac{\pi}{2} + \log \{ \sqrt{2} + 1 \} \right]$ .

7. Find the limiting value of  $x e^{-x^2} \int_0^x e^{x^2} dx$  when  $x = \infty$ . *Result.*  $\frac{1}{2}$

8. Shew that  $\int_0^{\infty} \frac{\cos ax - \cos bx}{x} dx = \log \frac{b}{a}$ ,

9. If  $F\left(x, \frac{1}{x}\right)$  be any symmetrical function of  $x$  and  $\frac{1}{x}$ , then

$$\int_0^{\infty} \frac{dx}{x F\left(x, \frac{1}{x}\right)} = 2 \int_0^1 \frac{dx}{x F\left(x, \frac{1}{x}\right)}.$$



10. If  $F(x)$  be an algebraical polynomial of less than  $n$  dimensions

$$\int_b^a \frac{F(x) dx}{(x-c)^n} = \frac{1}{n-1} \frac{d^{n-1}}{dc^{n-1}} \left\{ F(c) \log \frac{a-c}{b-c} \right\}.$$

11. Prove that  $\int_0^{2\pi} e^{\cos \theta} \cos (\sin \theta) d\theta = 2\pi$ .

12. Prove that  $\int_0^{\frac{\pi}{2}} \frac{\sqrt{1-c} d\theta}{1-c \cos^n \theta} = \frac{\pi}{\sqrt{2n}}$  when  $c$  is indefinitely nearly equal to unity,  $n$  being a positive quantity.

13. Evaluate  $\int_0^{\pi} (a \cos \theta + b \sin \theta) \log (a \cos^2 \theta + b \sin^2 \theta) d\theta$ .

*Result.*  $2b \left\{ \log a - 2 + \frac{2\sqrt{b}}{\sqrt{a-b}} \cos^{-1} \frac{\sqrt{b}}{\sqrt{a}} \right\}$ ,  
supposing  $a$  greater than  $b$ .

14. Shew that

$$\int_0^{\infty} \log \frac{1+2n \cos ax + n^2}{1+2n \cos bx + n^2} \cdot \frac{dx}{x} = \log(1+n) \log \frac{a^2}{b^2},$$

or  $\log \left( 1 + \frac{1}{n} \right) \log \frac{a^2}{b^2}$ , according as  $n$  is greater or less than unity.

15. Find the value of

$$\int_0^{\infty} [e^{-\{a+\alpha\sqrt{-1}\}x} - e^{-\{b+\beta\sqrt{-1}\}x}] \frac{dx}{x},$$

where  $a$  and  $b$  are positive, but  $\alpha$  and  $\beta$  positive or negative; and shew that it is wholly real when  $\frac{\alpha}{a} = \frac{\beta}{b}$ .

16. Prove that  $\int_0^1 \cot^{-1}(1-x+x^2) dx = \frac{\pi}{2} - \log 2$ .

17. Prove that  $\int_0^{\infty} \frac{dx}{1+x^2} \log \left( x + \frac{1}{x} \right) = \pi \log 2$ .

18. From the value of  $\int_0^{\infty} \frac{\sin x}{x} dx$  deduce that of

$$\int_0^{\infty} \left( \frac{\sin x}{x} \right)^2 dx.$$

*Result.* The two integrals are equal.

19. Prove that  $\int_0^{\infty} \left( \frac{e^{-ax} - e^{-bx}}{x} \right)^2 dx = \log \frac{(2a)^{2a} (2b)^{2b}}{(a+b)^{2(a+b)}}$ .

20. Shew that

$$\int_0^{\frac{\pi}{4}} \tan^{-1} \{ m \sqrt{1 - \tan^2 x} \} dx = \frac{\pi}{2} \tan^{-1} m \sqrt{2} - \frac{\pi}{2} \cot^{-1} \frac{\sqrt{1+m^2}}{m}.$$

21. Shew that  $\int_0^{\infty} (e^{-\frac{a^2}{x^2}} - e^{-\frac{b^2}{x^2}}) dx = (b-a) \sqrt{\pi}$ .

(*Solutions of Senate-House Problems*, by O'Brien and Ellis, p. 44.)

22. Shew that  $\int_0^{\infty} \log \frac{e^x + 1}{e^x - 1} dx = \frac{\pi^2}{4}$ .

23. Prove that  $\int_0^1 \frac{x^m - x^n}{\log x} \cdot \frac{dx}{x} = \log \frac{m}{n}$ , and reconcile with this equation the result of transforming  $\int_0^1 \frac{x^{r-1} dx}{\log x}$  by making  $x^r = y$ .

24. Shew that  $\int_0^{\frac{\pi}{2}} \sin^n \theta d\theta = \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma \left( \frac{n+1}{2} \right)}{\Gamma \left( \frac{n+2}{2} \right)}$ .

25. Shew that  $\int_0^1 \frac{x^{l-1} (1-x)^{m-1} dx}{(b+cx)^{l+m}} = \frac{\Gamma(l) \Gamma(m)}{\Gamma(l+m)} \frac{1}{b^m (b+c)^l}$ .

26. Shew that  $\int_0^{\frac{\pi}{2}} \frac{\cos^{2l-1} \theta \sin^{2m-1} \theta d\theta}{(a \cos^2 \theta + b \sin^2 \theta)^{l+m}} = \frac{\Gamma(l) \Gamma(m)}{2\Gamma(l+m)} \frac{1}{a^l b^m}$ .

27. Shew that  $\int_0^{\frac{\pi}{2}} \frac{\tan^n \theta d\theta}{a \cos^2 \theta + b \sin^2 \theta} = \frac{\pi}{2 \cos \frac{1}{2} n\pi} \frac{1}{a^{\frac{1-n}{2}} b^{\frac{1+n}{2}}}$ ,  
 $n$  being less than unity.

28. Shew that  $\int_0^{\pi} \frac{\sin^{n-1} \theta d\theta}{(\alpha + \beta \cos \theta)^n} = \frac{\left\{ \Gamma\left(\frac{n}{2}\right) \right\}^2}{\Gamma(n)} \frac{2^{n-1}}{(\alpha^2 - \beta^2)^{\frac{n}{2}}}$ .

29. Shew that  $\int_0^1 \frac{x^{m-1} dx}{(1-x^n)^n} = \frac{\pi}{n \sin \frac{m\pi}{n}}$ .

30. Shew that  $\int_0^1 \frac{x^{n-1} dx}{(1+cx)(1-x)^n} = \frac{\pi}{(1+c)^n \sin n\pi}$ .

31. Shew that  $\int_0^{\infty} \frac{\sin ax \sin^2 cx}{x} dx = 0, \pm \frac{\pi}{4},$  or  $\pm \frac{\pi}{8},$  according to the values of  $a$  and  $c$ .

32. Trace the locus of the equation

$$y = \int_0^{\infty} \frac{\sin \theta \cos \theta x}{\theta} d\theta.$$

33. Trace the locus of the equation

$$\frac{y}{b} = \int_0^{\pi} \log \{1 - 2e^{-u} \cos \theta + e^{-2u}\} d\theta,$$

where  $u = \sin \frac{x}{a}$ .

34. Trace the locus of the equation

$$y = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{x \cos \theta d\theta}{\sqrt{(x^2 + 2x \cos \theta + 1)}},$$

in which the sign of the square root is always taken so as to make the quantity in the denominator positive.

35. Shew that

$$\int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \sin x \sin^{-1}(\sin x \sin y) \, dx \, dy = \frac{\pi^2}{4} - \frac{\pi}{2}.$$

36. Compare the results obtained from

$$\int_0^{\infty} \int_0^{\infty} \sin ax \, e^{-xy} \, dx \, dy,$$

by performing the integrations in different order.

37. Find the value of  $\int_0^{\infty} e^{-\frac{x^2}{a^2} - \frac{b^2}{x^2}} \, dx$ , and hence shew that

$$\int_0^{\infty} \left( \frac{x^2}{a^2} + \frac{a^2}{x^2} \right) e^{-\frac{x^2}{a^2} - \frac{a^2}{x^2}} \, dx = \frac{5a \sqrt{\pi}}{4e^2} = 5 \int_0^{\infty} \left( \frac{x^2}{a^2} - \frac{a^2}{x^2} \right) e^{-\frac{x^2}{a^2} - \frac{a^2}{x^2}} \, dx.$$

38. Shew that

$$\iint \frac{\sqrt{(1-x^2-y^2)}}{\sqrt{(1+x^2+y^2)}} \, dx \, dy = \frac{\pi}{4} \left( \frac{\pi}{2} - 1 \right),$$

the integral being extended over all the values of  $x$  and  $y$  which make  $x^2 + y^2$  not greater than unity.

39. Shew that

$$\iiint \dots \frac{dx \, dy \, dz \dots}{\sqrt{(1-x^2-y^2-z^2-\dots)}} = \frac{\pi^{\frac{n+1}{2}}}{2^n \Gamma\left\{\frac{n+1}{2}\right\}},$$

the number of variables being  $n$ , and the integration being extended over all values which make

$$x^2 + y^2 + z^2 + \dots$$

not greater than unity.

40. If  $A_0 + A_1x + A_2x^2 + \dots = F(x)$ ,

and  $a_0 + a_1x + a_2x^2 + \dots = f(x)$ ,

prove that  $A_0a_0 + A_1a_1x + A_2a_2x^2 + \dots$

$$= \frac{1}{2\pi} \int_0^{2\pi} \{F(u) + F(v)\} \{f(u) + f(v)\} d\theta - A_0a_0$$

where  $u = xe^{\theta \sqrt{-1}}$  and  $v = xe^{-\theta \sqrt{-1}}$ .

Apply the above formula to express the sum of the series

$$\frac{x^3}{1^2} - \frac{x^6}{3^2} + \frac{x^{10}}{5^2} - \dots$$

41. If the sum of the series  $a_0 + a_1x + a_2x^2 + \dots$  can be expressed in a finite form, then the sum of the series  $a_0^2 + a_1^2x^2 + a_2^2x^4 + \dots$  can be expressed by a definite integral. Prove this, and hence shew that the sum of the squares of the coefficients of the terms of the expansion of  $(1+x)^n$  when  $n$  is a positive whole number, may be expressed by

$$\frac{2^{2n+2}}{\pi} \int_0^{\frac{\pi}{2}} \cos^{2n} \theta \cos^2 n\theta d\theta - 1.$$

42. Prove that

$$\int_0^{\infty} \frac{\cos cx dx}{1+x^2} = \frac{\pi}{2} \left\{ \frac{e^c}{1+0^c} + \frac{e^{-c}}{1+0^c} \right\}.$$

43. Shew that

$$\int_0^{\frac{\pi}{2}} \phi(\sin 2x) \cos x dx = \int_0^{\frac{\pi}{2}} \phi(\cos^2 x) \cos x dx.$$

(Liouville's *Journal de Mathématiques*, Vol. XVIII. page 168.)

44. Shew that  $1 - \frac{x^3}{2^2} + \frac{x^4}{2^2 4^2} - \dots$

$$= \frac{2}{\pi} \int_0^{\frac{\pi}{2}} \cos(x \sin y) dy.$$

45. Prove that

$$\int_0^{\infty} x^{m-1} e^{-x^n} dx \int_0^{\infty} y^{n-m-1} e^{-y^n} dy = \frac{\pi}{n^2 \sin \frac{m\pi}{n}}.$$

(See Art. 66; and change the variable  $y$  to  $u$  where  $y = ux$ .)

46. Shew that

$$\begin{aligned} \int_{-\infty}^{\infty} e^{-(a^2 \cos 2\theta + \frac{a^2}{2x^2} \sin 2\theta)} \frac{\cos}{\sin} \left\{ x^2 \sin 2\theta + \frac{a^2}{2x^2} \cos 2\theta \right\} dx \\ = \pi^{\frac{1}{2}} e^{-a} \frac{\cos}{\sin} (\theta + a); \end{aligned}$$

$\theta$  being comprised between the limits  $\pm \frac{\pi}{4}$ .

## CHAPTER XIII.

## EXPANSION OF FUNCTIONS IN TRIGONOMETRICAL SERIES.

301. THE subject we are about to introduce is one of the most remarkable applications of the Integral Calculus, and although in an elementary work like the present, only an imperfect outline can be given of it, yet on account of the novelty of the methods, and the importance of the results, even such an outline may be of service to the student. For fuller information we may refer to the *Differential and Integral Calculus* of Professor De Morgan. The subject is also frequently considered in the writings of Poisson, for example, in his *Traité de Mécanique*, Vol. i. pp. 643—653; in his *Traité de la Chaleur*; and in different Memoirs in the *Journal de l'Ecole Polytechnique*. The student may also consult a Memoir by Professor Stokes, in the 8th Vol. of the *Cambridge Philosophical Transactions*, a Memoir by Sir W. Hamilton, in the 19th Vol. of the *Transactions of the Royal Irish Academy*, and a Memoir by Professor Boole, in the 21st Vol. of the same *Transactions*.

302. It is required to find the values of the  $m$  constants  $A_1, A_2, A_3, \dots, A_m$ , so that the expression

$$A_1 \sin x + A_2 \sin 2x + A_3 \sin 3x + \dots + A_m \sin mx$$

may coincide in value with an assigned function of  $x$  when  $x$  has the values  $\theta, 2\theta, 3\theta, \dots, m\theta$ , where  $\theta = \frac{\pi}{m+1}$ .

Let  $f(x)$  denote the assigned function of  $x$ , then we have

by hypothesis the following  $m$  equations from which the constants are to be determined,

$$\begin{aligned}
 f(\theta) &= A_1 \sin \theta + A_2 \sin 2\theta + A_3 \sin 3\theta + \dots + A_m \sin m\theta, \\
 f(2\theta) &= A_1 \sin 2\theta + A_2 \sin 4\theta + A_3 \sin 6\theta + \dots + A_m \sin 2m\theta, \\
 &\dots\dots\dots \\
 f(m\theta) &= A_1 \sin m\theta + A_2 \sin 2m\theta + A_3 \sin 3m\theta + \dots + A_m \sin mm\theta.
 \end{aligned}$$

Multiply the first of these equations by  $\sin r\theta$ , the second by  $\sin 2r\theta$ , ....., the last by  $\sin mr\theta$ ; then add the results. The coefficient of  $A_s$  on the second side will then be

$$\sin r\theta \sin s\theta + \sin 2r\theta \sin 2s\theta + \dots + \sin mr\theta \sin ms\theta;$$

we shall now shew that this coefficient is zero if  $s$  be different from  $r$ , and equal to  $\frac{1}{2}(m+1)$  when  $s$  is equal to  $r$ .

First suppose  $s$  different from  $r$ . Now twice the above coefficient is equal to the series

$$\cos (r-s)\theta + \cos 2(r-s)\theta + \dots + \cos m(r-s)\theta,$$

diminished by the series

$$\cos (r+s)\theta + \cos 2(r+s)\theta + \dots + \cos m(r+s)\theta.$$

The sum of the first series is known from Trigonometry to be equal to

$$\frac{\sin (2m+1) \frac{(r-s)\theta}{2} - \sin \frac{(r-s)\theta}{2}}{2 \sin \frac{(r-s)\theta}{2}},$$

that is to 
$$\frac{\sin \left\{ (r-s)\pi - \frac{(r-s)\theta}{2} \right\} - \sin \frac{(r-s)\theta}{2}}{2 \sin \frac{(r-s)\theta}{2}}.$$

This expression vanishes when  $r-s$  is an odd number, and is equal to  $-1$  when  $r-s$  is an even number.

The sum of the second series can be deduced from that of the first by changing the sign of  $s$ ; hence this sum vanishes



when  $r + s$  is an odd number, and is equal to  $-1$  when  $r + s$  is an even number.

Thus when  $s$  is different from  $r$ , the coefficient of  $A_s$  is zero.

When  $s$  is equal to  $r$ , the coefficient becomes

$$\sin^2 r\theta + \sin^2 2r\theta + \dots + \sin^2 mr\theta,$$

that is,  $\frac{m}{2} - \frac{1}{2} \{ \cos 2r\theta + \cos 4r\theta + \dots + \cos 2mr\theta \}$ .

And by the method already used it will be seen that the sum of the series of cosines is  $-1$ ; thus the coefficient of  $A_r$  is  $\frac{1}{2}(m+1)$ .

Hence we obtain

$$A_r = \frac{2}{m+1} \{ \sin r\theta f(\theta) + \sin 2r\theta f(2\theta) + \dots + \sin mr\theta f(m\theta) \},$$

and thus by giving to  $r$  in succession the different integral values from 1 to  $m$ , the constants are determined.

Now suppose  $m$  to increase indefinitely, then we have ultimately

$$A_r = \frac{2}{\pi} \int_0^\pi \sin rv f(v) dv.$$

And as  $f(x)$  now coincides in value with the expression

$$A_1 \sin x + A_2 \sin 2x + \dots$$

for an infinite number of equidistant values of  $x$  between 0 and  $\pi$ , we may write the result thus

$$f(x) = \frac{2}{\pi} \sum_1^\infty \sin nx \int_0^\pi \sin nv f(v) dv,$$

where the symbol  $\sum_1^\infty$  indicates a summation to be obtained by giving to  $n$  every positive integral value.

303. The theorem and demonstration of the preceding article are due to Lagrange; we have given this demonstra-

tion partly because of its historical interest, and partly because it affords an instructive view of the subject. We shall however not stop to examine the demonstration closely, but proceed at once to the mode of investigation adopted by Poisson.

304. The following expansion may be obtained by ordinary Trigonometrical methods,

$$\frac{1 - h^2}{1 - 2h \cos \frac{\pi(v-x)}{l} + h^2} = 1 + 2h \cos \frac{\pi(v-x)}{l} + 2h^2 \cos \frac{2\pi(v-x)}{l} + 2h^3 \cos \frac{3\pi(v-x)}{l} + \dots,$$

$h$  being less than unity, so that the series is convergent.

Multiply both sides by  $\phi(v)$ , and integrate with respect to  $v$  between the limits  $-l$  and  $l$ ; also make  $h$  approach to unity as its limit. On the right-hand side the different powers of  $h$  become ultimately unity. The numerator of the fraction on the left-hand side will ultimately vanish, and thus the integral would vanish *if the denominator of the fraction were never zero*. But *if  $x$  lies between  $-l$  and  $l$* , the term  $\cos \frac{\pi(v-x)}{l}$  will become equal to unity during the integration, and thus the denominator of the fraction will be  $(1-h)^2$ , and will tend towards zero as  $h$  approaches unity. Thus the integral will not necessarily vanish; we proceed to ascertain its value. Let  $v-x=z$  and  $h=1-g$ , thus

$$\int \frac{(1-h^2) \phi(v) dv}{1 - 2h \cos \frac{\pi(v-x)}{l} + h^2} = \int \frac{g(1+h) \phi(x+z) dz}{g^2 + 4h \sin^2 \frac{\pi z}{2l}}.$$

Now the only part of the integral which has any sensible value, is that which arises from very small positive or negative values of  $z$ ; thus we may put

$$\sin \frac{\pi z}{2l} = \frac{\pi z}{2l},$$

and  $\phi(x+z) = \phi(x)$ ;

and the integral becomes

$$\begin{aligned} g(1+h)\phi(x) \int \frac{dz}{g^2 + \frac{h\pi^2 z^2}{l^2}} &= 2g\phi(x) \int \frac{dz}{g^2 + \frac{\pi^2 z^2}{l^2}} \\ &= \frac{2l\phi(x)}{\pi} \tan^{-1} \frac{\pi z}{gl}. \end{aligned}$$

Suppose  $\alpha$  and  $-\beta$  to be the limits of  $z$ ; we thus get

$$\frac{2l\phi(x)}{\pi} \left\{ \tan^{-1} \frac{\pi\alpha}{gl} + \tan^{-1} \frac{\pi\beta}{gl} \right\}.$$

Hence, finally, when  $g$  is supposed to vanish, we have  $2l\phi(x)$ . Thus if  $x$  lies between  $-l$  and  $l$ ,

$$\phi(x) = \frac{1}{2l} \int_{-l}^l \phi(v) dv + \frac{1}{l} \sum_1^\infty \int_{-l}^l \phi(v) \cos \frac{n\pi(v-x)}{l} dv.$$

If  $x$  does not lie between  $-l$  and  $l$ , the left-hand member must be replaced by zero. If however  $x=l$  or  $-l$ , then the integral on the left-hand side has its sensible part when  $v$  is indefinitely near to  $l$  and  $-l$ ; we should then have to perform the above process in both cases, but the integral with respect to  $z$  would only extend in the former case from  $-\beta$  to 0, and in the latter from 0 to  $\alpha$ . Hence instead of  $2l\phi(l)$  on the left-hand side, we should have

$$l\phi(l) + l\phi(-l).$$

305. In the same way as the result in Art. 313 is found, we have, if we integrate between 0 and  $l$ ,

$$\phi(x) = \frac{1}{2l} \int_0^l \phi(v) dv + \frac{1}{l} \sum_1^\infty \int_0^l \phi(v) \cos \frac{n\pi(v-x)}{l} dv \dots (1);$$

this holds if  $x$  has any value between 0 and  $l$ ; but when  $x=0$  the left-hand member must be  $\frac{1}{2}\phi(0)$ , and when  $x=l$  the left-hand member must be  $\frac{1}{2}\phi(l)$ ; for all other values of  $x$  the left-hand member should be zero.

Similarly

$$0 = \frac{1}{2l} \int_0^l \phi(v) dv + \frac{1}{l} \sum_1^\infty \int_0^l \phi(v) \cos \frac{n\pi(v+x)}{l} dv \dots\dots\dots(2);$$

this holds for any value of  $x$  between 0 and  $l$ ; but when  $x = 0$  the left-hand member must be  $\frac{1}{2} \phi(0)$ , and when  $x = l$  the left-hand member must be  $\frac{1}{2} \phi(l)$ .

From (1) and (2) by addition

$$\phi(x) = \frac{1}{l} \int_0^l \phi(v) dv + \frac{2}{l} \sum_1^\infty \cos \frac{n\pi x}{l} \int_0^l \cos \frac{n\pi v}{l} \phi(v) dv \dots(3).$$

This holds for any value of  $x$  between 0 and  $l$ , both inclusive.

From (1) and (2) by subtraction

$$\phi(x) = \frac{2}{l} \sum_1^\infty \sin \frac{n\pi x}{l} \int_0^l \sin \frac{n\pi v}{l} \phi(v) dv \dots\dots\dots(4).$$

This holds for any value of  $x$  between 0 and  $l$  both exclusive; and when  $x = 0$  or  $l$ , the left-hand member should be zero.

Equation (4) coincides with Lagrange's Formula.

We will now give some examples.

306. Expand  $x$  in a series of sines. Take formula (4) of Art. 314, and suppose  $l = \pi$ ; then

$$\int_0^\pi v \sin nv dv = -\frac{v \cos nv}{n} + \frac{\sin nv}{n^2};$$

therefore  $\int_0^\pi v \sin nv dv = \frac{\pi}{n}$  if  $n$  be odd, and  $-\frac{\pi}{n}$  if  $n$  be even.

Thus

$$x = 2 \left\{ \sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x - \frac{1}{4} \sin 4x + \dots\dots \right\}.$$

This holds for values of  $x$  between 0 and  $\pi$ , and as both sides vanish with  $x$  it holds when  $x = 0$ ; and it is obvious that if it holds for any positive value of  $x$  it holds for the

corresponding negative value; hence it holds for values of  $x$  between  $-\pi$  and  $\pi$ , exclusive of these limiting values.

307. Expand  $\cos x$  in a series of sines. Take formula (4) of Art. 314 and suppose  $l = \pi$ ; then

$$\begin{aligned} \int \cos v \sin nv \, dv &= \frac{1}{2} \int \{ \sin (n+1)v + \sin (n-1)v \} \, dv \\ &= -\frac{1}{2} \left\{ \frac{\cos (n+1)v}{n+1} + \frac{\cos (n-1)v}{n-1} \right\}; \end{aligned}$$

therefore  $\int_0^\pi \cos v \sin nv \, dv = 0$  if  $n$  is odd,

$$= \frac{2n}{n^2-1} \text{ if } n \text{ is even;}$$

therefore

$$\cos x = \frac{2}{\pi} \left\{ \frac{4}{3} \sin 2x + \frac{8}{15} \sin 4x + \dots + \frac{1 + (-1)^n}{n^2-1} n \sin nx + \dots \right\}.$$

This holds from  $x=0$  to  $x=\pi$ , exclusive of these limiting values.

308. Expand  $x$  in a series of cosines.

Take formula (3) of Art. 314, and suppose  $l = \pi$ ; then

$$\int v \cos nv \, dv = \frac{v \sin nv}{n} + \frac{\cos nv}{n^2};$$

therefore  $\int_0^\pi v \cos nv \, dv = \text{zero}$  if  $n$  be even, and  $-\frac{2}{n^2}$  if  $n$  be odd; and

$$\int_0^\pi v \, dv = \frac{\pi^2}{2},$$

thus  $x = \frac{\pi}{2} - \frac{4}{\pi} \left\{ \cos x + \frac{1}{3^2} \cos 3x + \frac{1}{5^2} \cos 5x + \dots \right\}.$

This holds from  $x=0$  to  $x=\pi$  both inclusive.

If we put  $x = \frac{\pi}{2} - y$ , we obtain the following formula,

which holds for any value of  $y$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ , both inclusive,

$$y = \frac{4}{\pi} \left\{ \sin y - \frac{1}{3^2} \sin 3y + \frac{1}{5^2} \sin 5y - \dots \right\}.$$

309. Expand  $e^{ax} - e^{-ax}$  in a series of sines.

Here 
$$\int_0^\pi (e^{av} - e^{-av}) \sin nv \, dv = -\frac{n(e^{a\pi} - e^{-a\pi})}{a^2 + n^2} \cos n\pi.$$

Therefore 
$$\frac{\pi}{2} \frac{e^{ax} - e^{-ax}}{e^{a\pi} - e^{-a\pi}} = \frac{\sin x}{1^2 + a^2} - \frac{2 \sin 2x}{2^2 + a^2} + \frac{3 \sin 3x}{3^2 + a^2} - \dots$$

310. Expand  $e^{a(\pi-x)} + e^{-a(\pi-x)}$  in a series of cosines.

Here 
$$\int_0^\pi \{e^{a(\pi-v)} + e^{-a(\pi-v)}\} \cos nv \, dv = \frac{a(e^{a\pi} - e^{-a\pi})}{a^2 + n^2},$$

and 
$$\int_0^\pi \{e^{a(\pi-v)} + e^{-a(\pi-v)}\} \, dv = \frac{e^{a\pi} - e^{-a\pi}}{a}.$$

Therefore 
$$\frac{\pi}{2a} \frac{e^{a(\pi-x)} + e^{-a(\pi-x)}}{e^{a\pi} - e^{-a\pi}} = \frac{1}{2a^2} + \frac{\cos x}{1^2 + a^2} + \frac{\cos 2x}{2^2 + a^2} + \dots$$

311. We have shewn that the formula (3) of Art. 314 holds for any value of  $x$  between 0 and  $l$  both inclusive; it is easy to determine what the right-hand member is equal to when  $x$  lies beyond these limits. Suppose  $x$  positive, and between  $l$  and  $2l$ ; put  $x = 2l - x'$  so that  $x'$  is less than  $l$ , then

$$\cos \frac{n\pi x}{l} = \cos \left( 2n\pi - \frac{n\pi x'}{l} \right) = \cos \frac{n\pi x'}{l};$$

therefore the value of the right-hand member is  $\phi(x')$ . Next suppose  $x$  greater than  $2l$ ; and suppose it equal to  $2ml + x'$  where  $x'$  is less than  $2l$ ; then

$$\cos \frac{n\pi x}{l} = \cos \frac{n\pi x'}{l},$$

so that the value is the same as it would be if  $x'$  were put instead of  $x$ ; that is, the value is  $\phi(x')$  if  $x'$  be less than  $l$ , and  $\phi(2l-x')$  if  $x'$  be greater than  $l$ .

It is obvious that for any negative value of  $x$  the value is the same as for the corresponding positive value.

Similarly we may shew that if  $x$  is positive and  $= 2ml + x'$ , the value of the right-hand side of equation (4) of Art. 314 is the same as if  $x'$  were put instead of  $x$ , and is  $\phi(x')$  if  $x'$  be less than  $l$ , and  $-\phi(2l-x')$  if  $x'$  be greater than  $l$ . And for negative values of  $x$  the value is the same numerically as for the corresponding positive value, but with an opposite sign.

312. It may be observed that in the fundamental demonstration of Art. 313, we suppose that when  $h$  approaches unity as a limit, the expression

$$\int h^n \phi(v) \cos \frac{n\pi(v-x)}{l} dv$$

may be replaced by

$$\int \phi(v) \cos \frac{n\pi(v-x)}{l} dv,$$

however large  $n$  may be. We may shew that no error arises from this supposition, by proving that the latter integral vanishes when  $n$  is increased indefinitely. For

$$\begin{aligned} \int \phi(v) \cos \frac{n\pi(v-x)}{l} dv &= \frac{l\phi(v)}{\pi n} \sin \frac{n\pi(v-x)}{l} \\ &\quad - \frac{l}{\pi n} \int \phi'(v) \sin \frac{n\pi(v-x)}{l} dv, \end{aligned}$$

which shews that the integral on the left-hand side will vanish when  $n$  is infinite, at least if  $\phi'(v)$  be not infinite.

313. We have not yet alluded to one of the most remarkable points in connexion with the formulæ (3) and (4) of Art. 314. In these formulæ  $\phi(x)$  need not be a *continuous function*; for example, from  $x=0$  to  $x=a$  we might have  $\phi(x) = f_1(x)$ , then from  $x=a$  to  $x=b$  we might have  $\phi(x) = f_2(x)$ , then from  $x=b$  to  $x=c$  we might have  $\phi(x) = f_3(x)$ , then from  $x=c$  to  $x=l$  we might have

$\phi(x) = f_1(x)$ . The formula (3) for instance would still be true for all values of  $x$  between 0 and  $l$  inclusive, as is evident from the mode of demonstration, *except* for the values where the discontinuity occurs. When for example  $x = a$ , then the value of the right-hand member would not be  $f_1(a)$  or  $f_2(a)$  but  $\frac{1}{2} \{f_1(a) + f_2(a)\}$ . If therefore for  $x = a$  we have  $f_1(x) = f_2(x)$ , the formula holds also when  $x = a$ .

314. Find an expression which shall be equal to  $c$  when  $x$  lies between 0 and  $a$ , and equal to zero when  $x$  lies between  $a$  and  $l$ .

Take formula (3) of Art. 314. Here  $\phi(v) = c$  from  $v = 0$  to  $v = a$ , and then from  $v = a$  to  $v = l$  it is zero; thus

$$\int_0^l \cos \frac{n\pi v}{l} \phi(v) dv$$

becomes  $c \int_0^a \cos \frac{n\pi v}{l} dv = \frac{cl}{n\pi} \sin \frac{n\pi a}{l}$

therefore the required expression is

$$\frac{ca}{l} + \frac{2c}{\pi} \left\{ \sin \frac{\pi a}{l} \cos \frac{\pi x}{l} + \frac{1}{2} \sin \frac{2\pi a}{l} \cos \frac{2\pi x}{l} + \frac{1}{3} \sin \frac{3\pi a}{l} \cos \frac{3\pi x}{l} + \dots \right\},$$

this will give  $\frac{1}{2}c$  when  $x = a$ .

Or we may use formula (4) of Art. 314. Then

$$c \int_0^a \sin \frac{n\pi v}{l} dv = \frac{cl}{n\pi} \left( 1 - \cos \frac{n\pi a}{l} \right),$$

and we have for the required expression

$$\frac{2c}{\pi} \left\{ \text{vers} \frac{\pi a}{l} \sin \frac{\pi x}{l} + \frac{1}{2} \text{vers} \frac{2\pi a}{l} \sin \frac{2\pi x}{l} + \frac{1}{3} \text{vers} \frac{3\pi a}{l} \sin \frac{3\pi x}{l} + \dots \right\};$$

this gives 0 when  $x = 0$ , and  $\frac{1}{2}c$  when  $x = a$ .

315. Find an expression which shall be equal to  $kx$  from  $x = 0$  to  $x = \frac{l}{2}$ , and equal to  $k(l-x)$  from  $x = \frac{l}{2}$  to  $x = l$ .



Here

$$\begin{aligned} \int_0^l \phi(v) \cos \frac{n\pi v}{l} dv &= \int_0^{\frac{l}{2}} kv \cos \frac{n\pi v}{l} dv + \int_{\frac{l}{2}}^l k(l-v) \cos \frac{n\pi v}{l} dv \\ &= \frac{kl^2}{\pi} \left\{ \frac{1}{2n} \sin \frac{n\pi}{2} + \frac{1}{n^2\pi} \cos \frac{n\pi}{2} - \frac{1}{n^2\pi} \right\} + \frac{kl^2}{n\pi} \left( \sin n\pi - \sin \frac{n\pi}{2} \right) \\ &\quad - \frac{kl^2}{\pi} \left\{ \frac{1}{n} \sin n\pi - \frac{1}{2n} \sin \frac{n\pi}{2} + \frac{\cos n\pi}{n^2\pi} - \frac{\cos \frac{n\pi}{2}}{n^2\pi} \right\} \\ &= \frac{kl^2}{\pi^2 n^2} \{ 2 \cos \frac{n\pi}{2} - \cos n\pi - 1 \}. \end{aligned}$$

This is  $-\frac{4kl^2}{\pi^2 n^2}$  when  $n$  is of the form  $4r+2$ , and 0 in every other case, and

$$\int_0^l \phi(v) dv = k \int_0^{\frac{l}{2}} v dv + k \int_{\frac{l}{2}}^l (l-v) dv = \frac{kl^2}{4};$$

thus the required expression is

$$\frac{kl}{4} - \frac{8kl}{\pi^2} \left\{ \frac{1}{2^2} \cos \frac{2\pi x}{l} + \frac{1}{6^2} \cos \frac{6\pi x}{l} + \dots \right\}.$$

If we denote this by  $y$  then from  $x=0$  to  $x=\frac{1}{2}l$  both inclusive  $y=kx$ , then from  $x=\frac{1}{2}l$  to  $x=l$  both inclusive  $y=k(l-x)$ ; for values of  $x$  greater than  $l$  the values of  $y$  recur as shewn in Art. 320. Thus the value of  $y$  is the ordinate of the figure formed by measuring from the origin equal lengths along the axis of  $x$  to the right and left, and drawing on each base thus obtained the same isosceles triangle.

316. As another example we may propose the following: find a function  $\phi(x)$  which shall be equal to  $x$  from  $x=0$  to  $x=\alpha$ , then be equal to  $\alpha$  from  $x=\alpha$  to  $x=\pi-\alpha$ , and then be equal to  $\pi-x$  from  $x=\pi-\alpha$  to  $x=\pi$ .

The result is

$$\phi(x) = \frac{4}{\pi} \left\{ \sin \alpha \sin x + \frac{1}{3^2} \sin 3\alpha \sin 3x + \frac{1}{5^2} \sin 5\alpha \sin 5x + \dots \right\};$$

this is true from  $x=0$  to  $x=\pi$  both inclusive.

317. The student may verify the following examples.

If  $x$  be numerically less than  $a$  the expression

$$\frac{8a}{\pi^2} \sum_0^\infty \left\{ \frac{\cos (2n+1) \frac{\pi x}{2a}}{2n+1} \right\}^2$$

is equal to  $a - x$  if  $x$  be positive, and  $a + x$  if  $x$  be negative.

Prove that for values of  $x$  between  $-\pi$  and  $\pi$  inclusive

$$\frac{x^2}{4} = \frac{\pi^2}{12} - \cos x + \frac{\cos 2x}{2^2} - \frac{\cos 3x}{3^2} + \dots$$

This may be obtained from Art. 315 by integration; or from equation (3) of Art. 314.

318. In the formula

$$\phi(x) = \frac{1}{2l} \int_{-l}^l \phi(v) dv + \frac{1}{l} \sum_1^\infty \int_{-l}^l \cos \frac{n\pi(v-x)}{l} \phi(v) dv,$$

suppose  $l$  to increase without limit; then if  $\phi(v)$  be such that the term  $\frac{1}{2l} \int_{-l}^l \phi(v) dv$  vanishes with  $\frac{1}{l}$  we have

$$\phi(x) = \frac{1}{\pi} \int_0^\infty \int_{-\infty}^\infty \cos u(v-x) \phi(v) du dv.$$

This is called Fourier's Theorem.

## CHAPTER XIV.

APPLICATION OF THE INTEGRAL CALCULUS TO QUESTIONS  
OF MEAN VALUE AND PROBABILITY.

319. WE will here give a few simple examples of the application of the Integral Calculus to questions relating to *mean value* and to *probability*.

Let  $\phi(x)$  denote any function of  $x$ , and suppose  $x$  successively equal to  $a, a+h, a+2h, \dots a+(n-1)h$ . Then

$$\frac{\phi(a) + \phi(a+h) + \phi(a+2h) + \dots + \phi\{a+(n-1)h\}}{n}$$

may be said to be the *mean* or *average* of the  $n$  values which  $\phi(x)$  receives corresponding to the  $n$  values of  $x$ . Let

$$b - a = nh,$$

then the above *mean value* may be written thus

$$\frac{[\phi(a) + \phi(a+h) + \phi(a+2h) + \dots + \phi\{a+(n-1)h\}] h}{b - a}.$$

Suppose  $a$  and  $b$  to remain fixed and  $n$  to increase indefinitely; then the limit of the above expression is

$$\frac{\int_a^b \phi(x) dx}{b - a}.$$

This may accordingly be defined to be the *mean value* of  $\phi(x)$  when  $x$  varies continuously between  $a$  and  $b$ .

320. As an example we may take the following question; find the mean distance of all points within a circle from a fixed point on the circumference. By this enunciation we intend the following process to be performed. Let the area of a circle be divided into a large number  $n$  of equal small areas; form a fraction of which the numerator is the sum of the distances of these small areas from a fixed point on the circumference, and the denominator is  $n$ ; then find the limit of this fraction when  $n$  is infinite.

Suppose  $r_1, r_2, \dots, r_n$  to denote the respective distances of the small areas; then the fraction required is

$$\frac{1}{n} \{r_1 + r_2 + \dots + r_n\}.$$

Multiply both numerator and denominator by  $r \Delta\theta \Delta r$ , which represents the area of a small element (Art. 148), thus the fraction becomes

$$\frac{\{r_1 + r_2 + \dots + r_n\} r \Delta\theta \Delta r}{nr \Delta\theta \Delta r}.$$

The limit of the denominator will represent the area of the circle, that is  $\pi c^2$ , if  $c$  be the radius of the circle. The limit of the numerator will be, by the definitions of the Integral Calculus,  $\iint r^2 d\theta dr$ , the limits being so taken as to include all the elements of area within the boundary of the circle. Thus the result is

$$\frac{\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2c \cos \theta} r^2 d\theta dr}{\pi c^2}.$$

This will be found to give  $\frac{32c}{9\pi}$ .

321. The equation to a curve is  $r = c \sin \theta \cos \theta$ , find the *mean length* of all the radii vectores drawn at equal angular intervals in the first quadrant.

It easily follows, as in the last article, that the required *mean length* is

$$\frac{\int_0^{\frac{\pi}{2}} c \sin \theta \cos \theta d\theta}{\frac{\pi}{2}},$$

that is,  $\frac{c}{\pi}$ .

Again, suppose the portion of this curve which lies in the first quadrant to revolve round the initial line, and thus to generate a surface. Let radii vectores be drawn from the origin to different points of the surface *equally in all directions*: it is required to find the mean length of the radii vectores.

The only difficulty in this question lies in apprehending clearly what is meant by the words in Italics. Conceive a spherical surface having the origin as centre; then by equable angular distribution of the radii vectores, we mean that they are to be so drawn that the number of them which fall upon any portion of the spherical surface must be proportional to the area of that portion. Now the area of any portion of a sphere of radius  $a$  is found by integrating

$$a^2 \iint \sin \theta d\phi d\theta$$

within proper limits (Art. 175). Hence  $a^2 \sin \theta \Delta\phi \Delta\theta$  may be taken to denote an element of a spherical surface, and  $2\pi a^2$  is the area of half the surface of a sphere. Thus we shall have as the required result

$$\frac{\iint a^2 c \sin \theta \cos \theta \sin \theta d\phi d\theta}{2\pi a^2},$$

the limits being so taken as to extend the integrations over the entire surface considered.

Hence we obtain

$$\frac{\int_0^{2\pi} \int_0^{\frac{\pi}{2}} c \sin^2 \theta \cos \theta \, d\phi \, d\theta}{2\pi},$$

that is,  $\frac{c}{3}$ .

322. A large plane area is ruled with parallel equidistant lines; a thin rod, the length of which is less than the distance between two consecutive lines, is thrown at hazard on the area; find the probability that the rod will fall across one of the lines. Let  $2a$  be the distance between two consecutive lines and  $2c$  the length of the rod. It is easily seen that we do not alter the problem by supposing the centre of the rod constrained to fall upon a line drawn between consecutive lines of the given system and meeting them at right angles, for the proportion of the favourable cases to the whole number of cases remains the same after this limitation as before.

Let the centre of the rod be at a distance  $x$  from the nearer of the two selected parallels; then suppose the rod to revolve round its centre, and it is obvious that in this position of its centre the chance that it crosses the line is  $\frac{4\phi}{2\pi}$ , where

$$c \cos \phi = x.$$

And we may denote by  $\frac{\Delta x}{a}$  the chance that the centre of the rod falls between the distances  $x$  and  $x + \Delta x$  from the nearer of the two parallels. Thus the chance required will be denoted by the limit of the sum of such quantities as  $\frac{2\phi}{\pi} \frac{\Delta x}{a}$ , that is, it will be

$$\frac{2}{\pi a} \int \phi \, dx,$$

where  $\cos \phi = \frac{x}{c}$ .

The limits of  $x$  are 0 and  $c$ ; hence the result

$$\begin{aligned} &= \frac{2c}{\pi a} \int_0^{\frac{\pi}{2}} \phi \sin \phi \, d\phi \\ &= \frac{2c}{\pi a}. \end{aligned}$$

### EXAMPLES.

1. If  $r = f(\theta)$  and  $y = f\left(\frac{x}{a}\right)$  be the equations to two curves,  $f(\theta)$  being a function which vanishes for the values  $\theta_1, \theta_2$ , and is positive for all values between these limits, and if  $A$  be the area of the former between the limits  $\theta = \theta_1$  and  $\theta = \theta_2$ , and  $M$  the arithmetical mean of all the transverse sections of the solid generated by the revolution about the axis of  $x$  of the portion of the latter curve between the limits  $x = a\theta_1$  and  $x = a\theta_2$ , shew that

$$M = \frac{2\pi}{\theta_2 - \theta_1} A,$$

supposing  $\theta_2$  greater than  $\theta_1$ .

2. A ball is fired at random from a gun which is equally likely to be presented in any direction in space above the horizon; shew that the chance of its reaching more than  $\frac{1}{m}$ th of its greatest range is  $\sqrt{1 - \frac{1}{m}}$ .
3. From a point in the circumference of a circular field a projectile is thrown at random with a given velocity, which is such that the diameter of the field is equal to the greatest range of the projectile; find the chance of its falling within the field.
4. On a table a series of straight lines at equal distances from one another is drawn and a cube is thrown at

random on the table. Find the chance of its resting without covering any part of the lines.

5. Prove that the mean of all the radius-vectors of an ellipse, the focus being the origin, is equal to half the minor axis.
6. An indefinite number of equidistant parallel lines are drawn on a plane, and a rod whose length is equal to  $r$  times the perpendicular distance between two consecutive lines is thrown at random on the plane; find the probability of its falling upon  $n$  of the lines. If  $n = r = 1$ , shew that the probability is  $\frac{2}{\pi}$ .
7. Two arrows are sticking in a circular target; what is the chance that their distance is greater than the radius of the target?
8. Supposing the orbits of comets to be equably distributed through space, prove that their mean inclination to the plane of the ecliptic is the angle subtended by an arc equal to the radius.
9. A certain territory is bounded by two meridian circles and by two parallels of latitude which differ in longitude and latitude respectively by one degree, and is known to lie within certain limits of latitude; find the probable superficial area.
10. A line is taken of given length  $a$ , and two other lines are taken each less than the first line and laid down in it at hazard, any one position of either being as likely as any other. The lengths of these lines are  $b$  and  $b'$ ; it is required to find the probability that they shall not have a part exceeding  $c$  in common.

$$\text{Result } \frac{(a - b - b' + c)^2}{(a - b)(a - b')}.$$

*Camb. Phil. Transactions*, Vol. VIII. p. 386.



11. An indefinitely large plane area is ruled with parallel equidistant lines, the distance between consecutive lines being  $c$ . An ellipse whose major axis is less than  $c$  is thrown down on the area. Shew that the chance that the ellipse falls on one of the lines is  $\frac{l}{\pi c}$  where  $l$  denotes the perimeter of the ellipse.
12. A messenger  $M$  starts from  $A$  towards  $B$  (distance  $a$ ) at a rate of  $v$  miles per hour, but before he arrives at  $B$  a shower of rain commences at  $A$  and at all places occupying a certain distance  $z$  towards, but not reaching beyond,  $B$ , and moves at the rate of  $u$  miles an hour towards  $A$ ; if  $M$  be caught in this shower he will be obliged to stop until it is over; he is also to receive for his errand a number of shillings inversely proportional to the time occupied in it, at the rate of  $n$  shillings for one hour. Supposing the distance  $z$  to be unknown, as also the time at which the shower commenced, but all events to be equally probable, shew that the value of  $M$ 's expectation is, in shillings,

$$\frac{nv}{a} \left\{ \frac{1}{2} - \frac{u}{v} + \frac{u(u+v)}{v^2} \log \frac{u+v}{u} \right\}.$$

THE END.

# EDUCATIONAL MATHEMATICAL WORKS.

By I. TODHUNTER, M.A.

*Fellow and Assistant Tutor of St. John's College, Cambridge.*

1. A Treatise on the Differential Calculus :  
And the Elements of the Integral Calculus. With numerous  
Examples. Second Edition, revised. Crown 8vo. cloth, 10s. 6d.

“TO THE DIFFERENT CHAPTERS WILL BE FOUND APPENDED EXAMPLES  
SUFFICIENTLY NUMEROUS TO RENDER ANOTHER BOOK UNNECESSARY.  
The examples have been selected almost exclusively from the College  
and University Examination Papers.”—*Preface.*

“A TREATISE WHICH WILL TAKE ITS PLACE AMONGST OUR STANDARD  
EDUCATIONAL WORKS. THE EXPLANATIONS IN THE EARLY PARTS OF  
THE VOLUME ARE CLEAR AND CONVINCING, AND CANNOT FAIL TO  
INTEREST THE STUDENT.”—*English Journal of Education.*

“FOR THE GREAT BULK OF MATHEMATICAL STUDENTS—ESPECIALLY AT  
THE BEGINNING OF THEIR ACQUAINTANCE WITH THE CALCULUS—WE  
HAVE NOT SEEN A TREATISE SO WELL ADAPTED AS THE PRESENT.”—  
*Athenæum.*

2. A Treatise on the Integral Calculus,  
And its Applications; with numerous Examples. Crown 8vo.  
cloth, 10s. 6d.

MR. TODHUNTER'S CLASS-BOOKS,—CONTINUED.

3. A Treatise on Analytical Statics.

With numerous Examples. Crown 8vo. cloth, 10s. 6d.

"MR PRATT'S TREATISE ON MECHANICAL PHILOSOPHY is now out of print, and the present work may be regarded as a republication, with large additions, of that portion of it which treats of Statics."—PREFACE.

"We know of no other work which is so admirably adapted to the wants of a large class of Mathematical Students."—ATHENÆUM.

"One of the most satisfactory evidences of the advancement of Education in Cambridge, is the recent improvements of the University Class-Books. The perspicuous language, rigorous investigation, conscientious scrutiny of difficulties, and methodical treatment which characterise Mr Todhunter's works, entitle him to a large share of the honour of that improvement."—CIVIL ENGINEER'S AND ARCHITECT'S JOURNAL.

"The Treatise is an excellent one, fully sustaining the high character of its Author, and the various subjects which it embraces have received additional interest from the judicious observations and lucid explanations which Mr Todhunter has liberally interspersed throughout the Volume, tending not only to simplify processes, but to facilitate the acquisition of principles ..... The Treatise is admirably adapted either for School or Collegiate instruction, and will form a first-rate Text-book on Analytical Statics."—ENGLISH JOURNAL OF EDUCATION.

4. A Treatise on Plane Co-ordinate Geometry,

As applied to the Straight Line and the CONIC SECTIONS.

With numerous Examples. Crown 8vo. cloth, 10s. 6d.

"Mr Todhunter has laboured to make the treatment of the different portions of the subject simple and easy to be understood."—EDUCATIONAL TIMES.

5. A Treatise on Algebra,

For the use of Students at the Universities, and the Higher Classes in Schools. *Preparing.*

6. An Elementary Work on ALGEBRA,

For the use of Beginners.

*Preparing.*

## MATHEMATICAL CLASS-BOOKS.

### A Treatise on Dynamics.

With numerous Examples. By P. G. TAIT, M.A., Fellow of St. Peter's College, and Professor of Mathematics in Queen's College, Belfast, and W. J. STEELE, late Fellow of St. Peter's College. Crown 8vo. cloth, 10s. 6d.

*"This is a new Edition of that part of PRATT'S MECHANICAL PHILOSOPHY which treats of DYNAMICS, with large additions and improvements."*

*"A well-wrought book on the Cambridge-plan, with copious questions for exercise."*—ATHENÆUM, Feb. 23, 1856.

### An Elementary Treatise on Mechanics.

For the use of the Junior Classes at the University, and the Higher Classes in Schools. With a Collection of Examples. By S. PARKINSON, B.D., Fellow and Tutor of St. John's College, Cambridge. Crown 8vo. cloth, 9s. 6d.

*"He thoroughly understands his subject, he has gone to its root, and he has treated it like a man who is completely master of it. Its evolution in this book is the natural evolution of first principles, gradually and simply elucidated, yet eventually stated and applied in their most general forms. ....The language is clear.....The book should be in the hands of all who wish to make themselves master of the principles of Theoretical Mechanics."*—BRITISH QUARTERLY REVIEW.

*"We do not know of a better book."*—ATHENÆUM.

### Elementary Hydrostatics.

With numerous Examples. By J. B. PHEAR, M.A., Fellow and Assistant Tutor of Clare College, Cambridge, Author of "Elementary Mechanics." Crown 8vo. cloth, 5s. 6d.

*"An excellent introductory Book. The definitions are very clear; the descriptions and explanations are sufficiently full and intelligible; the investigations are simple and scientific. The examples greatly enhance its value."*—ENGLISH JOURNAL OF EDUCATION.

MATHEMATICAL CLASS-BOOKS.

By J. C. SNOWBALL, M. A.

*Fellow of St. John's College, Cambridge.*

*A New and Cheaper (THE NINTH) Edition of*

1. The Elements of Plane and Spherical Trigonometry.

Greatly improved and enlarged. Crown 8vo. bound in cloth, 7s. 6d.

This edition has been carefully revised by the Author, and some important alterations and additions have been introduced. A large addition has been made to the collection of *Examples for practice*.

*"A new Edition of an old favourite text-book, and an improvement on the seven that have preceded it in several respects. It has been carefully revised throughout; the methods for establishing the most important propositions are superior; more than 200 new examples—taken from recent Examination Papers—have been added; and to crown all, the price has been reduced. What more need be said to secure for it a welcome from those who wish to make themselves masters of the important subject of which it treats?"—ATHENÆUM.*

2. An Introduction to the Elements of Plane Trigonometry.

Designed for the use of Schools. Second Edition. 8vo. 5s.

3. The Cambridge Course of Elementary Natural Philosophy.

Being the Propositions in Mechanics and Hydrostatics in which persons who are not Candidates for Honors are examined for the Degree of B.A. To which are added numerous Examples and Problems from the University Examination Papers, and Original, with Hints for their Solutions. Fourth Edition. Crown 8vo. bound in cloth, 5s.

CAMBRIDGE, *November* 1856.

MACMILLAN & CO.'S  
PUBLICATIONS.

**ÆSCHYLI Eumenides.**

The Greek Text with English Notes: with an Introduction, containing an Analysis of C. O. Müller's Dissertations; and an English Metrical Translation. By BERNARD DRAKE, M.A., Fellow of King's College, Cambridge; Editor of "Demosthenes de Corona." 8vo. cloth, 7s. 6d.

**ANTHOLOGIA Latina Selecta. In 2 vols. Small 8vo.**

VOL. I.—Containing select Epigrams of Catullus, Virgil, Claudian, Ausonius, with others from the Anthologia Latina.

VOL. II.—Containing select Epigrams of Martial.

Edited with English notes, by J. E. B. MAYOR, M.A., Fellow and Classical Lecturer of St. John's College, Cambridge, Editor of *Juvenal*.

[*Preparing.*]

**ARISTOPHANES. The Greek Text revised, with a Commentary.** By W. G. CLARK, M.A., Fellow and Assistant Tutor of Trinity College.

[*Preparing.*]

**ARISTOTELES de Rhetorica. The Greek Text, with English Notes.** By A FELLOW OF TRINITY COLLEGE.

[*Preparing.*]

**ARISTOTLE on the Vital Principle. Translated from the Original Text, with Notes.** By CHARLES COLLIER, M.D., F.R.S., Fellow of the Royal College of Physicians. Crown 8vo. cloth, 8s. 6d.

**BEAMONT.—Catherine, the Egyptian Slave in 1852. A Tale of Eastern Life.** By W. J. BEAMONT, M.A., Fellow of Trinity College, Cambridge, late Principal of the English College, Jerusalem. Fcap. 8vo. 5s. 6d.

**BENGLII GNOMON NOVI TESTAMENTI, in quo ex Nativæ Verborum vi simplicitas, profunditas, concinnitas, salubritas ensuum celestium indicatur.** EDITIO TERTIA. Imperial 8vo. cloth, 18s.

**BIRKS.—The Difficulties of Belief in connexion with the Creation and the Fall.** By THOMAS RAWSON BIRKS, M.A., Rector of Kelshall, Formerly Fellow of Trinity College; Author of "The Life of the Rev. G. Bickersteth." Crown 8vo. cloth, 4s. 6d.

CONTENTS. Introduction.—On the Knowledge of God. Chap. I.—On the Power of God. II.—On the Nature of God. III.—On the Creation of Free Agents. IV.—On Temptation in Free Agents. V.—On the Creation and Fall of Angels. VI.—On the Creation and Fall of Man. VII.—On the Permission of Satanic Temptation. VIII. & IX.—On Original Sin.

**BOLTON.**—The Evidences of Christianity as Exhibited in the Writings of its Apologists down to Augustine. An Essay which obtained the Hulsean Prize for the Year 1852. By W. J. BOLTON, of Gonville and Caius College, Cambridge. 8vo. cloth, 6s.

**BRAVE WORDS for BRAVE SOLDIERS and SAILORS.**  
Tenth Thousand. 16mo. sewed, 2d.; or 10s. per 100.

[Printed for Distribution.

**BUTLER (Professor Archer).**—Sermons, Doctrinal and Practical. By the Rev. WILLIAM ARCHER BUTLER, M.A. late Professor of Moral Philosophy in the University of Dublin. Edited, with a Memoir of the Author's Life, by the Very Rev. THOMAS WOODWARD, M.A. Dean of Down. With Portrait. Third Edition. 8vo. cloth, 12s.

**BUTLER (Professor Archer).**—A Second Series of Sermons. Edited from the Author's Manuscripts, by J. A. JEREMIE, D.D. Regius Professor of Divinity in the University of Cambridge. 8vo. cloth, 10s. 6d.

**BUTLER (Professor Archer).**—Lectures on the History of Ancient Philosophy. By the Rev. W. ARCHER BUTLER, late Professor of Moral Philosophy in the University of Dublin. Edited, from the Author's Manuscripts, by WILLIAM HEZFWORTH THOMPSON, M.A. Regius Professor of Greek in the University of Cambridge. 2 vols. 8vo. cloth, 11. 5s.

**BUTLER (Professor Archer).**—Letters on Romanism, in Reply to Mr. NEWMAN's Essay on Development. Edited by the Very Rev. T. WOODWARD, Dean of Down. 8vo. cloth, 10s. 6d.

**BUTLER.**—A Hand Book to Butler's Analogy.

With a few Notes. By C. A. SWAINSON, M.A. Principal of Chichester Theological College, formerly Fellow and Tutor of Christ's College, Cambridge. Crown 8vo. 1s. 6d.

**CAMBRIDGE.**—Cambridge Theological Papers. Comprising those given at the Voluntary Theological and Crosse Scholarship Examinations. Edited, with References and Indices, by A. P. MOOR, M.A. of Trinity College, Cambridge, and Sub-warden of St. Augustine's College, Canterbury. 8vo. cloth, 7s. 6d.

**CAMBRIDGE PROBLEMS.**—Solutions of the Senate-House Riders for Four Years (1848 to 1851). By F. J. JAMESON, M.A. Fellow of Caius College, Cambridge. 8vo. cloth, 7s. 6d.

**CAMBRIDGE PROBLEMS.**—Solutions of Senate-House Problems for Four Years (1848 to 1851). By N. M. FERRERS, and J. S. JACKSON, Fellows of Caius College, Cambridge. 8vo. cloth, 15s. 6d.

**CAMBRIDGE PROBLEMS, 1854.**—Solutions of the Problems proposed in the Senate House Examination, January 1854. By the Moderators (W. WALTON, M.A. Trinity College, and C. F. MACKENZIE, M.A. Fellow of Caius College). In 8vo. cloth, 10s. 6d.

**CAMBRIDGE.**—**Cambridge Guide: Including Historical and Architectural Notices of the Public Buildings, and a concise Account of the Customs and Ceremonies of the University, with a Sketch of the Places most worthy of Note in the County.** A New Edition, with Engravings and a Map. 12mo. cloth, 5s.

**CAMBRIDGE FITZWILLIAM MUSEUM.**—**A Hand-Book to the Pictures in the Fitzwilliam Museum, Cambridge.** Crown 8vo. sewed, 1s. 6d.; or in cloth elegant, 2s. 6d.

**CAMBRIDGE.**—**Cambridge Mathematical Journal.** Vol. I. Second Edition, 8vo. cloth, 18s.

**CAMBRIDGE.**—**Cambridge and Dublin Mathematical Journal.** *The Complete Work, in Nine Vols.* 8vo. cloth, 7l. 4s.

ONLY A FEW COPIES OF THE COMPLETE WORK REMAIN ON HAND.

**CAMPBELL.**—**The Nature of the Atonement and its Relation to Remission of Sins and Eternal Life.** By JOHN McLEOD CAMPBELL. 8vo. cloth, 10s. 6d.

**COLENSO.**—**Ten Weeks in Natal.** A Journal of a First Tour of Visitation among the Colonists and Zulu Kafirs of Natal. By the Right Rev. JOHN WILLIAM COLENSO, D.D. Bishop of Natal, with a Map and Illustrations. Fcap. 8vo. cloth, 5s.

**COLENSO.**—**Village Sermons.** By the Right Rev. JOHN WILLIAM COLENSO, D.D. Bishop of Natal. **Second Edition.** Fcap. 8vo. cloth, 2s. 6d.

**COLENSO.**—**The Communion Service, from the Book of Common-Prayer, with Select Readings from the writings of the Rev. F. D. MAURICE, M.A.** Edited by the Right Rev. JOHN WILLIAM COLENSO, D.D., Bishop of Natal. *Fine Edition, rubricated and bound in morocco, antique style, 6s.; or in cloth, 2s. 6d. Common Paper, limp cloth, 1s.*

**COOPER.**—**A Geometrical Treatise on Conic Sections.** By the Rev. J. E. COOPER, M.A. late Fellow of St. John's College, Cambridge. [Preparing.]

**COTTON.**—**Sermons: chiefly connected with Public Events of 1854.** By G. E. LYNCH COTTON, M.A. Master of Marlborough College, formerly Fellow of Trinity College, Cambridge. Fcap. 8vo. cloth, 3s.

**DAVIES.**—**St. Paul and Modern Thought:** Remarks on some of the Views advanced in Professor Jowett's Commentary on St. Paul. By Rev. J. LL. DAVIES, M.A. Fellow of Trinity College, Cambridge, and Incumbent of St. Mark's, Whitechapel. 8vo. 2s. 6d.

**DEMOSTHENES.**—**Demosthenes de Corona.** The Greek Text, with English Notes. By BERNARD DRAKE, M.A. Fellow of King's Coll. Cambridge, Editor and Translator of the "Eumenides of Eschylus." Crown 8vo. cloth, 5s.



- DEMOSTHENES.**—Translation of Demosthenes on the Crown. By J. P. NORRIS, M.A. Fellow of Trinity College, Cambridge, and one of Her Majesty's Inspectors of Schools. Crown 8vo. cloth, 3s.
- DRAKE.**—Notes Explanatory and Critical on the Books of Jonah and Hosea. By WILLIAM DRAKE, M.A. late Fellow of St. John's College, Cambridge. 8vo. cloth, 9s.
- EVANS.**—Sonnets on the Death of the Duke of Wellington, by SEBASTIAN EVANS. 8vo. sewed, 1s.
- FERRAR.**—Lives of Nicholas Ferrar, Fellow of Clare Hall. By his BROTHER JOHN, and Dr. JEBB. Now first Edited, with Illustrations, by J. E. B. MAYOR, M.A., Fellow and Assistant Tutor of St. John's College, Cambridge. Fcap. 8vo. cloth, 7s. 6d.
- FROST.**—The First Three Sections of Newton's Principia. With Notes and Problems in illustration of the subject. By PERCIVAL FROST, M.A. late Fellow of St. John's College, Cambridge, and Mathematical Lecturer of Jesus College. Crown 8vo. cloth, 10s. 6d.
- FROST.**—Thucydides, Book VI. The Greek Text, and English Notes: with a Map of Syracuse. By PERCIVAL FROST, Jun. M.A. late Fellow of St. John's College, Cambridge. 8vo. cloth, 7s. 6d.
- GODFRAY.**—An Elementary Treatise on the Lunar Theory. With a brief Sketch of the History of the Problem up to the time of Newton. By HUGH GODFRAY, B.A. of St. John's College, Cambridge. 8vo. cloth. 5s. 6d.
- GRANT.**—Plane Astronomy. Including Explanations of Celestial Phenomena, and Descriptions of Astronomical Instruments. By A. R. GRANT, M.A., one of Her Majesty's Inspectors of Schools, late Fellow of Trinity College, Cambridge. 8vo. boards, 6s.
- HALLIFAX.**—Bishop Hallifax's Analysis of the Civil Law. In which a comparison is occasionally made between the Roman Law and those of England. A new Edition, with alterations and additions, being the heads of a Course of Lectures publicly delivered in the University of Cambridge, by J. W. GELDART, LL.D. 8vo. bds. 8s. 6d.; interleaved, 10s. 6d.; double interleaved, 12s. 6d.
- HAMILTON.**—On Truth and Error: Thoughts, in Prose and Verse, on the Principles of Truth, and the Causes and Effects of Error. By JOHN HAMILTON, (of St. Ernan's,) M.A. of St. John's College, Cambridge. Crown 8vo. cloth, 10s. 6d.
- HARE.**—Charges to the Clergy of the Archdeaconry of Lewes, delivered at the Ordinary Visitations from the Year 1840 to 1854. With Notes on the Principal Events affecting the Church during that period. By JULIUS CHARLES HARE, M.A. Archdeacon. With an Introduction, explanatory of his position in the Church with reference to the parties which divide it. 3 vols. 8vo. cloth, 11. 11s. 6d.

**HARE.**—Charges to the Clergy of the Archdeaconry of Lewes, delivered at the Ordinary Visitations in the Years 1843, 1845, 1846. By JULIUS CHARLES HARE, M.A. Archdeacon. Never before published. With an Introduction, explanatory of his position in the Church with reference to the parties which divide it. 6s. 6d.

\*.\* This is included in the three vols. of Charges.

**HARE.**—Miscellaneous Pamphlets on some of the Leading Questions agitated in the Church during the last Ten Years. 8vo. cloth, 12s.

**HARE.**—The Victory of Faith.

Second Edition. 8vo. cloth, 5s.

**HARE.**—The Mission of the Comforter.

Second Edition. With Notes. 8vo. cloth, 12s.

**HARE.**—Vindication of Luther from his English Assailants.

Second Edition. 8vo. cloth, 7s.

**HARE.**—The Contest with Rome.

With Notes, especially in answer to Dr. Newman's recent Lectures. Second Edition. 8vo. cloth, 10s. 6d.

\*.\* This is included in the three vols. of Charges.

**HARE.**—Parish Sermons.

Second Series. 8vo. cloth, 12s.

**HARE.**—Portions of the Psalms in English Verse.

Selected for Public Worship. 18mo. cloth, 2s. 6d.

**HARDWICK.**—Christ and other Masters.

An Historical Inquiry into some of the chief Parallels and Contrasts between Christianity and the Religious Systems of the Ancient World. With special reference to prevailing Difficulties and Objections. By CHARLES HARDWICK, M.A., Fellow of St Catherine's Hall, Divinity Lecturer at King's College, and Christian Advocate in the University of Cambridge. Pt. I. 8vo. cloth, 7s. 6d. [Part II. in the Press.]

**HARDWICK.**—A History of the Christian Church, during the

Middle Ages. By CHARLES HARDWICK, M.A. Fellow of St. Catherine's Hall, Divinity Lecturer in King's College, and Christian Advocate in the University of Cambridge. Author of "A History of the XXXIX. Articles." With Four Maps constructed for this Work by A. KEITH JOHNSTON. Crown 8vo. cloth, 10s. 6d.

**HARDWICK.**—A History of the Christian Church during the Reformation. By CHARLES HARDWICK, M.A. Crown 8vo. cloth,

10s. 6d.

\*.\* These two Books are part of a Series of Theological Manuals now in progress.

**HARDWICK.**—Twenty Sermons for Town Congregations. By

CHARLES HARDWICK, M.A. Crown 8vo. cloth, 6s. 6d.

**HARE.**—Two Sermons preached in Herstmonceux Church, on Septuagesima Sunday, February 4, 1855, being the Sunday after the Funeral of the Venerable Archdeacon Hare. By the Rev. H. VENN ELLIOTT, Perpetual Curate of St. Mary's, Brighton, late Fellow of Trinity College, Cambridge, and the Rev. J. N. SIMPKINSON, Rector of Brighton, Northampton, formerly Curate of Herstmonceux. 8vo. 1s. 6d.

**HEMMING.**—An Elementary Treatise on the Differential and Integral Calculus. For the Use of Colleges and Schools. By G. W. HEMMING, M.A. Fellow of St. John's College, Cambridge. **Second Edition**, with Corrections and Additions. 8vo. cloth, 8s.

**HERVEY.**—The Genealogies of our Lord and Saviour Jesus Christ, as contained in the Gospels of St. Matthew and St. Luke, reconciled with each other and with the Genealogy of the House of David, from Adam to the close of the Canon of the Old Testament, and shown to be in harmony with the true Chronology of the Times. By Lord ARTHUR HERVEY, M.A. Rector of Ickworth with Horringer. 8vo. cloth, 10s. 6d.

**HERVEY.**—The Inspiration of Holy Scripture.

Five Sermons preached before the University of Cambridge, in the month of December 1855. 8vo. cloth, 3s. 6d.

**HOWARD.**—The Book of Genesis, according to the Version of the LXX. Translated into English, with Notices of its Omissions and Insertions, and with Notes on the Passages in which it differs from our Authorized Version. By the Hon. HENRY E. J. HOWARD, D.D. Dean of Lichfield. Crown 8vo. cloth, 8s. 6d.

**HOWARD.**—The Books of Exodus and Leviticus.

Uniform with the above.

[In the Press.]

**HOWES.**—A History of the Christian Church during the First Six Centuries. By J. G. HOWES, M.A. Fellow of St. Peter's Coll. Camb.

[Nearly ready.]

\*.\* This is part of a Series of Theological Manuals now in progress.

**HUMPHREYS.**—Exercitationes Iambicæ; or, Progressive Exercises in Greek Iambic Verse. To which are prefixed the Rules of Greek Prosody, with copious Notes and Illustrations of the Exercises. By E. R. HUMPHREYS, LL.D. Head Master of the Cheltenham Grammar School. **Second Edition.** Fcap. cloth, 5s. 6d.

**INGLEBY.**—Outlines of Theoretical Logic.

Founded on the New Analytic of SIR WILLIAM HAMILTON. Designed for a Text-book in Schools and Colleges. By C. MANSFIELD INGLEBY, M.A. of Trinity College, Cambridge, Teacher of Logic in the Industrial Department of the Birmingham and Midland Institute. In fcap. 8vo. cloth, 8s. 6d.

[Just ready.]

**JEWELL.**—An Apology of the Church of England, and an Epistle to Seignor Scipio concerning the Council of Trent, translated from the original Latin, and illustrated with Notes, chiefly drawn from the Author's "Defence of the Apology." By A. T. RUSSELL. Fcp. 8vo. bds. 5s.

**JUSTIN MARTYR.**—S. *Justini Philosophi et Martyris Apologia Prima.* Edited, with a corrected Text, and English Introduction and explanatory Notes, by W. TROLLOPE, M.A. Pembroke College, Cambridge. 8vo. bds. 7s. 6d.

**JUVENAL.**—*Juvenal: chiefly from the Text of Jahn.*

With English Notes for the Use of Schools. By J. E. B. MAYOR, M.A. Fellow and Classical Lecturer of St. John's College, Cambridge. Crown 8vo. cloth, 10s. 6d.

**KENNEDY.**—*The Influence of Christianity upon International Law.* The Hulsean Prize Essay in the University of Cambridge for the year 1854. Crown 8vo. cloth, 4s.

**KINGSLEY.**—"Westward Ho!" or, the Voyages and Adventures of Sir Amyas Leigh, Knight of Burrough, in the County of Devon, in the Reign of Her Most Glorious Majesty Queen Elizabeth. By CHARLES KINGSLEY. **Second Edition.** 3 vols. crown 8vo. 1l. 11s. 6d.

**KINGSLEY.**—*The Heroes: or, Greek Fairy Tales for my Children.* With Eight Illustrations after Drawings by the Author. In 8vo. beautifully printed on tinted paper, and elegantly bound in cloth, with gilt leaves, 7s. 6d.

"If the public accept our recommendation, Mr. Kingsley's little book will run through many editions."—*Guardian*, March 12, 1856.

**KINGSLEY.**—*Glaucus; or, the Wonders of the Shore.*

**Third Edition,** corrected and enlarged. With a Frontispiece. Fcap. 8vo. elegantly bound in cloth, with gilt leaves, 3s. 6d.

**KINGSLEY.**—*Alexandria and Her Schools: being Four Lectures delivered at the Philosophical Institution, Edinburgh.* With a Preface. Crown 8vo. cloth, 5s.

**KINGSLEY.**—*Phaethon; or Loose Thoughts for Loose Thinkers.* **Second Edition.** Crown 8vo. boards, 2s.

**LATHAM.**—*Geometrical Problems in the Properties of Conic Sections.* By H. LATHAM, M.A. Fellow and Tutor of Trinity Hall. 8vo. sewed, 3s. 6d.

**LECTURES to Ladies on Practical Subjects.**

Delivered in London during the month of July, 1855, by the Rev. F. D. MAURICE, Professor TRENCH, Archdeacon ALLEN, J. S. BREWER, J. LL. DAVIES, CHARLES KINGSLEY, Dr. CHAMBERS, Dr. SIEVEKING, Dr. JOHNSON, TOM TAYLOR, Esq., and F. J. STEPHEN, Esq. **Second Edition.** Crown 8vo. cloth, 7s. 6d.

**LETTERS from Italy and Vienna.**

Small 8vo. cloth, 5s. 6d.

**LUSHINGTON.**—*La Nation Boutiquière: and other Poems,* chiefly Political. With a Preface. By the late HENRY LUSHINGTON, Chief Secretary to the Governor of Malta. **Points of War.** By FRANKLIN LUSHINGTON, Judge in the Supreme Courts of the Ionian Isles. In 1 vol. fcap. 8vo. cloth, 3s.

**MACKENZIE.—The Christian Clergy of the first Ten Centuries: their Influence on European Civilization.** By HENRY MACKENZIE.

B.A. formerly Scholar of Trinity College, Cambridge. Crown 8vo. cloth, 6s. 6d.

**MANSFIELD.—Paraguay, Brazil, and the Plate.**

With a Map, and numerous Woodcuts. Edited from the Author's MSS. With a Sketch of his Life. By the Rev. CHARLES KINGSLEY. Crown 8vo. cloth, 12s. 6d. [Just ready.]

**MANSFIELD.—On the Constitution of Salts.**

Edited from the Author's MS. by N. H. S. MASKELYNE, M.A. Wadham College, and Reader in Mineralogy in the University of Oxford. [In the Press.]

**M'COY.**—Preparing for Publication; to be completed in about Five Parts, price 5s. each, forming One Volume 8vo. of about 500 pages, with nearly 1,000 illustrations in the text, drawn and engraved by the Author,

**A Manual of the Genera of British Fossils.**

Comprising Systematic Descriptions of all the Classes, Orders, Families, and Genera of Fossil Animals found in the Strata of the British Isles; with figures of all the Generic Types. By FREDERICK M'COY, F.G.S., Hon. F.C.P.S., Professor of Natural History in the University of Melbourne, Author of "Characters of the Carboniferous Limestone Fossils of Ireland," "Synopsis of the Silurian Fossils of Ireland," one of the Authors of "Sedgwick and M'Coyle's British Palæozoic Rocks and Fossils," &c.

**M'COY.**—Preparing for Publication, in One Volume, crown 8vo. with numerous Illustrations,

**An Elementary Introduction to the Study of Palæontology.**

With numerous Figures illustrative of Structural Details.

\*.\* This little Work is intended to supply all that elementary information on the Structure of Fossil Animals, with reference to the most nearly allied existing types, illustrated explanation of technical terms, &c. which the beginner may require, but which would be out of place in the Author's systematic volume on the Genera.

**M'COY.—Contributions to British Palæontology; or, First Descriptions of several hundred Fossil Radiata, Articulata, Mollusca, and Pisces, from the Tertiary, Cretaceous, Oolitic, and Palæozoic Strata of Great Britain.**

With numerous Woodcuts. 8vo. cloth, 9s.

\*.\* This forms a complete Series of the Author's Papers from the "Annals of Natural History."

**MASSON.—Essays, Biographical and Critical; chiefly on the English Poets.** By DAVID MASSON, M.A. Professor of English Literature in University College, London. 8vo. cloth, 12s. 6d.

With numerous Woodcuts. 8vo. cloth, 9s.

**MAURICE.—Discourses on the Gospel according to St. John.**

Crown 8vo. cloth. [Nearly ready.]

**MAURICE.—A Photograph Portrait of Rev. F. D. Maurice.**

4to. price 6s.

**MAURICE.—Lectures on the Ecclesiastical History of the First and Second Centuries.** By FREDERICK DENISON MAURICE, M.A. Chaplain of Lincoln's Inn. 8vo. cloth, 10s. 6d.

- MAURICE.**—The Unity of the New Testament, being a Synopsis of, and Commentary, on the first three Gospels, and the Epistles of St. James, St. Jude, St. Peter, and St. Paul. 8vo. cloth, 14s.
- MAURICE.**—Patriarchs and Lawgivers of the Old Testament. **Second Edition.** With new Preface. Crown 8vo. cloth, 6s.
- MAURICE.**—The Prophets and Kings of the Old Testament. Crown 8vo. cloth. **Second Edition.** 10s. 6d.
- MAURICE.**—Theological Essays. **Second Edition,** with a new Preface and other additions. Crown 8vo. cloth, 10s. 6d.
- MAURICE.**—The Doctrine of Sacrifice deduced from the Scriptures. With a Dedicatory Letter to the Young Men's Christian Association. Crown 8vo. cloth, 7s. 6d.
- MAURICE.**—Christmas Day, and other Sermons. 8vo. cloth, 10s. 6d.
- MAURICE.**—The Religions of the World, and their relations to Christianity. **Third Edition.** Fcap. 8vo. cloth, 5s.
- MAURICE.**—The Prayer-Book considered, especially in reference to the Romish System. **Second Edition.** Fcap. 8vo. cloth, 5s. 6d.
- MAURICE.**—The Church a Family. Twelve Sermons on the Occasional Services of the Prayer-Book. Fcap. 8vo. cloth, 4s. 6d.
- MAURICE.**—On the Lord's Prayer. **Third Edition.** Fcap. 8vo. cloth, 2s. 6d.
- MAURICE.**—On the Sabbath Day: the Character of the Warrior; and on the Interpretation of History. Fcap. 8vo. cloth, 2s. 6d.
- MAURICE.**—Learning and Working.—Six Lectures delivered in Willis's Rooms, London, in June and July, 1854. **The Religion of Rome, and its influence on Modern Civilization.**—Four Lectures delivered in the Philosophical Institution of Edinburgh, in December 1854. In One Volume, Crown 8vo. cloth, 5s.
- MAURICE.**—An Essay on Eternal Life and Eternal Death, and the Preface to the new Edition of "Theological Essays." Crown 8vo. sewed, 1s. 6d.  
\* \* \* Published separately for the purchasers of the first edition.
- MAURICE.**—Death and Life. A Sermon Preached in the Chapel of Lincoln's Inn, March 25, 1855. *In Memoriam G. B. M.* 8vo. sewed, 1s.
- MAURICE.**—Plan of a Female College for the Help of the Rich and of the Poor. A Lecture delivered at the Working Men's College, London, to a Class of Ladies. 8vo. 6d.
- MAURICE.**—Administrative Reform. A Lecture delivered at the Working Men's College, London. Crown 8vo. 3d.

- MAURICE.**—The Word "Eternal," and the Punishment of the Wicked. A Letter to the Rev. Dr. Jelf, Principal of King's College. London. **Fifth Thousand.** 8vo. 1s.
- MAURICE.**—The Name "Protestant;" the Seemingly Double Character of the English Church; and the English Bishopric at Jerusalem. Three Letters to the Rev. Wm. Palmer, Fellow and Tutor of Magdalen College, Oxford. **Second Edition.** 8vo. 3s.
- MAURICE.**—On Right and Wrong Methods of Supporting Protestantism. A Letter to Lord Ashley. 8vo. 1s.
- MAURICE.**—Thoughts on the Duty of a Protestant, in the Oxford Election of 1847. 8vo. 1s.
- MAURICE.**—The Case of Queen's College, London. A Letter to the Lord Bishop of London, in reply to the "Quarterly Review." 8vo. 1s. 6d.
- MAURICE.**—Lectures on Modern History and English Literature. [Preparing.]
- MAURICE.**—Law's Remarks on the Fable of the Bees, with an Introduction of Eighty Pages by FREDERICK DENISON MAURICE, M.A. Chaplain of Lincoln's Inn. Fcp. 8vo. cloth, 4s. 6d.  
 "This introduction discusses the Religious, Political, Social, and Ethical Theories of our day, and shows the special worth of Law's method, and how far it is applicable to our circumstances."
- MINUCIUS FELIX.**—The Octavius of Minucius Felix. Translated into English by LORD HAILES. Fcp. 8vo. cloth, 3s. 6d.
- NAPIER.**—Lord Bacon and Sir Walter Raleigh. Critical and Biographical Essays. By MACVEY NAPIER, late Editor of the *Edinburgh Review* and of the *Encyclopædia Britannica*. Post 8vo. cloth, 7s. 6d.
- NIND.**—Sonnets of Cambridge Life. By Rev. W. NIND, M.A. Fellow of St. Peter's College. Post 8vo. boards, 2s.
- NIND.**—The German Lyrist; or, Metrical Versions from the principal German Lyric Poets. By Rev. W. NIND, Fellow of St. Peter's College. Crown 8vo. cloth, 3s.
- NORRIS.**—Ten School-Room Addresses. Edited by J. P. NORRIS, M.A. Fellow of Trinity College, and one of Her Majesty's Inspectors of Schools. 18mo. sewed, 8d.
- PALEY.**—An Analysis of Paley's Evidences of Christianity, in the form of Question and Answer. With the Senate House Papers for the Year 1854. By CHARLES H. CROSSE, M.A. of Gonville and Caius College. 18mo. boards, 3s. 6d.
- PARKINSON.**—A Treatise on Elementary Mechanics. For the Use of the Junior Classes at the University, and the Higher Classes in Schools. With a Collection of Examples. By S. PARKINSON, M.A. Fellow and Assistant Tutor of St. John's College, Cambridge. Crown 8vo. cloth, 9s. 6d.

**PARMINTER.**—Materials for a Grammar of the Modern English Language. Designed as a Text-book of Classical Grammar for the use of Training Colleges, and the Higher Classes of English Schools. By GEORGE HENRY PARMINTER, of Trinity College, Cambridge; Rector of the United Parishes of SS. John and George, Exeter. Fcap. 8vo. cloth, 3s. 6d.

**PAYN.**—Poems.

By JAMES PAYN. Fcp. 8vo. cloth, 5s.

**PEACE IN WAR.** In Memoriam R. R.

Crown 8vo. sewed, 1s.

**PEARSON.**—Elements of the Calculus of Finite Differences, treated on the Method of the Separation of Symbols. By J. PEARSON, M.A. Rector of St. Edmund's, Norwich, Mathematical Master of Norwich Grammar School, and formerly Scholar of Trinity College, Cambridge. **Second Edition**, enlarged. 8vo. 5s.

**PEROWNE.**—"Al-Adjrumiieh."

An Elementary Arabic Grammar, with a Translation. By J. J. S. PEROWNE, M.A. Fellow of Corpus Christi College, Cambridge, and Lecturer in Hebrew in King's College, London. 8vo. cloth, 5s.

**PERRY.**—Five Sermons Preached before the University of Cambridge, in November 1855. By the Right Rev. CHARLES PERRY, Lord Bishop of Melbourne, formerly Fellow and Tutor of Trinity College, Cambridge. Crown 8vo. cloth, 3s.

**PHEAR.**—Elementary Mechanics.

Accompanied by numerous Examples solved Geometrically. By J. B. PHEAR, M.A., Fellow and Mathematical Lecturer of Clare Hall, Cambridge. 8vo. cloth, 10s. 6d.

**PHEAR.**—Elementary Hydrostatics.

Accompanied by numerous Examples. Crown 8vo. cloth, 5s. 6d.

**PLATO.**—The Republic of Plato.

Translated into English, with Notes. By Two Fellows of Trinity College, Cambridge, (J. Ll. Davies M.A., and D. J. Vaughan, M.A.) **Second Edition.** 8vo. cloth. [Preparing.]

**POWELL.**—The Scriptural Doctrine of the Influence of the

Holy Ghost, as Illustrated by the Analogy of Nature. The Burney Prize Essay for the year 1853. 8vo. sewed, 2s. 6d.

**PRATT.**—The Mathematical Principles of Mechanical

Philosophy. By J. H. PRATT, M.A., Fellow of Caius College.

\* \* \* The above work is now out of Print: but the Part on STATICS has been re-edited by Mr. Todhunter, with numerous alterations and additions: the Part on DYNAMICS, by Messrs. Tait and Steele, is just published. The other parts will be published in separate forms, improved and altered as may seem needful.

**PRINCIPLES of ETHICS according to the NEW TESTAMENT.** Crown 8vo. 2s.



**PROCTER.**—**A History of the Book of Common Prayer: with a Rationale of its Offices.** By FRANCIS PROCTER, M.A., Vicar of Witton, Norfolk, and late Fellow of St. Catherine's Hall. **Second Edition,** revised and enlarged. Crown 8vo. cloth, 10s. 6d.

\*.\* This is part of a series of Theological Manuals, now in progress.

**PUCKLE.**—**An Elementary Treatise on Conic Sections and Algebraic Geometry.** With a numerous collection of Easy Examples progressively arranged, especially designed for the use of Schools and Beginners. By G. HALE PUCKLE, M.A., St. John's College, Cambridge; Principal of Windermere College. **Second Edition,** enlarged and improved. Crown 8vo. cloth, 7s. 6d.

**PURTON.**—**The Acts of the Apostles.**

With a Paraphrase and Exegetical Commentary. By JOHN SMYTH PURTON, M.A. Fellow and Tutor of St. Catherine's Hall, Cambridge. 8vo.

[Preparing.]

**RAMSAY.**—**The Catechiser's Manual; or, the Church Catechism Illustrated and explained, for the use of Clergymen, Schoolmasters, and Teachers.** By ARTHUR RAMSAY, M.A. of Trinity College, Cambridge. 18mo. cloth, 3s. 6d.

**REICHEL.**—**The Lord's Prayer and other Sermons.**

By C. P. REICHEL, B.D., Professor of Latin in the Queen's University; Chaplain to his Excellency the Lord Lieutenant of Ireland; and late Donnellan Lecturer in the University of Dublin. Crown 8vo. cloth, 7s. 6d.

**ROBINSON.**—**Autobiography of Matthew Robinson.**

Now first published. With Illustrations. By J. E. B. MAYOR, M.A. Fellow and Assistant Tutor of St. John's College. Fcap. 8vo. uniform with Lives of Ferrar, bound in cloth, 5s. 6d.

[Just ready.]

**ROBINSON.**—**Missions urged upon the State on grounds both of Duty and Policy.** An Essay which obtained the Maitland Prize in the year 1852. By C. K. ROBINSON, M.A., Fellow and Assistant Tutor of St. Catherine's Hall, Cambridge. Fcp. 8vo. cloth, 3s.

**ROSE (Henry John).**—**An Exposition of the Articles of the Church of England.** By HENRY JOHN ROSE, B.D. late Fellow of St. John's College, and Hulsean Lecturer in the University of Cambridge.

[Preparing.]

\*.\* This is part of a Series of Theological Manuals now in progress.

**SALLUST.**—**Sallust.**

The Latin Text, with English Notes. By CHARLES MERIVALE, B.D.; late Fellow and Tutor of St. John's College, Cambridge, &c., Author of the "History of Rome," &c. Crown 8vo. cloth, 5s.

**SEDGWICK AND M'COY'S British Palæozoic Fossils.**

Part I. 4to. sewed, 16s.

Part II. 4to. sewed, 10s.

Part III. completing the

work, 16s. just ready.

\*.\* THE WHOLE BOUND IN TWO VOLS., 4to. cloth, £2 2s.

**SELWYN.**—*The Work of Christ in the World. Four Sermons*, preached before the University of Cambridge, on the four Sundays preceding Advent in the year of our Lord 1854. By the Right Rev. GEORGE AUGUSTUS SELWYN, D.D. Bishop of New Zealand, formerly Fellow of St. John's College. **Third Edition.** Crown 8vo. 2s.

**SELWYN.**—*A Verbal Analysis of the Holy Bible.*

Intended to facilitate the translation of the Holy Scriptures into Foreign Languages. Compiled by THE BISHOP OF NEW ZEALAND, for the use of the Melanesian Mission. Small folio, cloth, 14s.

**SIMPSON.**—*An Epitome of the History of the Christian Church during the first Three Centuries and during the Time of the Reformation*, adapted for the use of Students in the Universities and in Schools. By WILLIAM SIMPSON, M.A. With Examination Questions. **Second Edition**, Improved. Fcp. 8vo. cloth, 5s.

**SMITH.**—*Arithmetic and Algebra, in their Principles and Application*: with numerous systematically arranged Examples, taken from the Cambridge Examination Papers. With especial reference to the ordinary Examination for B.A. Degree. By BARNARD SMITH, M.A., Fellow of St. Peter's College, Cambridge. **Third Edition**, enlarged and revised throughout. Crown 8vo. cloth, 10s. 6d.

**SMITH.**—*Arithmetic for the use of Schools.* By BARNARD SMITH, M.A. Fellow of St. Peter's College. **Fourth Thousand**, with Additions. Crown 8vo. cloth, 4s. 6d.

\* \* \* This has been published in accordance with very numerous requests from Schoolmasters and Inspectors of Schools. It comprises a complete reprint of the Arithmetic from Mr. Smith's larger work, with such alterations as were necessary in separating it from the Algebra, and references throughout to the *Decimal System of Coinage*.

**SMITH.**—*A Key to Mr. Smith's Arithmetic for Schools.*

Crown 8vo. cloth, 8s. 6d.

**SMITH.**—*Mechanics and Hydrostatics, in their Principles and Application*: with numerous systematically arranged Examples, taken from the Cambridge Examination Papers. With a special reference to the Ordinary Examination for B.A. Degree. By BARNARD SMITH, M.A. Fellow of St. Peter's College, Cambridge. [*Preparing.*]

**SNOWBALL.**—*The Elements of Plane and Spherical Trigonometry.* Greatly improved and enlarged. By J. C. SNOWBALL, M.A. Fellow of St. John's College, Cambridge. **Eighth Edition**, with Addition and Improvements. Crown 8vo. cloth, 7s. 6d.

**SWAINSON.**—*A Hand Book to Butler's Analogy.*

With a few Notes. By C. A. SWAINSON, M.A. Principal of the Chichester Theological College, formerly Fellow and Tutor of Christ's College Cambridge, and Preacher at Whitehall. Crown 8vo. 1s. 6d.

**TAIT and STEELE.**—A Treatise on Dynamics, with numerous Examples. By P. G. TAIT, Fellow of St. Peter's College, and Professor of Mathematics in Queen's College, Belfast, and W. J. STEELE, Fellow of St. Peter's College. Crown 8vo. cloth, 10s. 6d. [Just ready.]

This is a new Edition of that part of Pratt's Mechanical Philosophy which treats of Dynamics, with large additions and improvements.

**TAYLOR.**—The Restoration of Belief.

By ISAAC TAYLOR. Crown 8vo. cloth, 6s. 6d.

**THEOCRITUS.**—Theocritus.

The Greek Text, with English notes, Critical and Explanatory, for the use of Colleges and Schools. By E. H. PEROWNE, M.A. Fellow of Corpus Christi College. Crown 8vo. [Byegones.]

**THEOLOGICAL Manuals.**

Just published:—

**CHURCH HISTORY: THE MIDDLE AGES.** By CHARLES HARDWICK. With Four Maps. Crown 8vo. cloth, price 10s. 6d.

**THE COMMON PRAYER: ITS HISTORY AND RATIONALE.** By FRANCIS PROCTER. Second Edition. Crown 8vo. cloth, 10s. 6d.

**A HISTORY OF THE CANON OF THE NEW TESTAMENT.** By B. F. WESTCOTT. Crown 8vo. cloth, 1s. 6d. 4s. 7d. 9c.

**CHURCH HISTORY: THE REFORMATION.** By CHARLES HARDWICK. Crown 8vo. cloth, 10s. 6d. aa

The following will shortly appear:—

**INTRODUCTION TO THE STUDY OF THE OLD TESTAMENT.**

**NOTES ON ISAIAH.**

**INTRODUCTION TO THE STUDY OF THE GOSPELS.**  
EPISTLES.

**NOTES ON THE GOSPELS AND ACTS.**

EPISTLES AND APOCALYPSE.

**CHURCH HISTORY, THE FIRST SIX CENTURIES.**

17th CENTURY TO THE PRESENT TIME.

**THE THREE CREEDS.**

**THE THIRTY-NINE ARTICLES.**

\*.\* Others are in progress, and will be announced in due time.

**THRING.**—A Construing Book.

Compiled by the Rev. EDWARD THRING, M.A. late Fellow of King's College, Cambridge, and Head Master of Uppingham School. Fcap. 8vo. cloth, 2s. 6d.

**THRING.**—The Elements of Grammar taught in English.

By EDWARD THRING, M.A. Head Master of the Royal Grammar School, Uppingham; late Fellow of King's College, Cambridge. Second Edition. 18mo. bound in cloth, 2s.

**THRING.**—The Child's Grammar.

Being the substance of the above, with Examples for Practice. Adapted for Junior Classes. A New Edition. 18mo. limp cloth, 1s.

- THRUPP.**—Psalms and Hymns for Public Worship. Selected and Edited by JOSEPH FRANCIS THRUPP, M.A. Vicar of Barrington, late Fellow of Trinity College. 18mo. cloth, 2s. Second paper in limp cloth, 1s. 4d.
- THRUPP.**—Antient Jerusalem: a New Investigation into the History, Topography, and Plan of the City, Environs, and Temple. Designed principally to illustrate the records and prophecies of Scripture. With Map and Plans. By JOSEPH FRANCIS THRUPP, M.A. Vicar of Barrington, Cambridge, late Fellow of Trinity College. 8vo. cloth, 15s.
- TODHUNTER.**—A Treatise on the Differential Calculus; and the Elements of the Integral Calculus. With numerous Examples. By I. TODHUNTER, M.A., Fellow and Tutor of St. John's College, Cambridge. **Second Edition.** Crown 8vo. cloth, 10s. 6d.
- TODHUNTER.**—A Treatise on the Integral Calculus. With numerous Examples. Crown 8vo. cloth. [*In the Press.*]
- TODHUNTER.**—A Treatise on Analytical Statics, with numerous Examples. Crown 8vo. cloth, 10s. 6d.
- TODHUNTER.**—A Treatise on Conic Sections. With numerous Examples. For the Use of Colleges and Schools. Crown 8vo. cloth, 10s. 6d.
- TODHUNTER.**—A Treatise on Algebra, for the Use of Students in the Universities, and of the Higher Classes in Schools. [*Prep.*]  
*Also by the same Author,*
- An Elementary Work on the same subject, for the use of Beginners.
- TRENCH.**—Synonyms of the New Testament. By RICHARD CHENEVIX TRENCH, B.D., Vicar of Itchenstoke, Hants, Professor of Divinity, King's College, London, and Examining Chaplain to the Bishop of Oxford. **Third Edition,** revised. Fcp. 8vo. cloth, 5s.
- TRENCH.**—Hulsean Lectures for 1845—46.  
CONTENTS. 1.—The Fitness of Holy Scripture for unfolding the Spiritual Life of Man. 2.—Christ the Desire of all Nations; or the Unconscious Prophecies of Heathendom. **Third Edition.** Foolscap 8vo. cloth, 5s.
- VAUGHAN.**—Sermons preached in St. John's Church, Leicester, during the years 1855 and 1856. Crown 8vo. cloth, 5s. 6d.
- WATERS OF COMFORT.**—A Small Volume of Devotional Poetry of a Practical Character. By the Author of "Visiting my Relations." Fcap. 8vo. cloth, 4s.
- WESTCOTT.**—A general View of the History of the Canon of the New Testament during the First Four Centuries. By BROOKE FOSS WESTCOTT, M.A., Assistant Master of Harrow School; late Fellow of Trinity College, Cambridge. Crown 8vo. cloth, 12s. 6d.

**WESTCOTT.**—An Introduction to the Study of the Gospels; including a new and improved Edition of "The Elements of the Gospel Harmony." With a Catena on Inspiration, from the Writings of the Antenicene Fathers. Crown 8vo. [Preparing.]

**WESTCOTT.**—An Introduction to the Study of the Canonical Epistles; including an attempt to determine their separate purposes and mutual relations. By BROOKE FOSS WESTCOTT, M.A. [Preparing.]

\*.\* These three books are part of a series of Theological Manuals now in progress.

**WILSON.**—The Five Gateways of Knowledge.

By GEORGE WILSON, M.D., F.R.S.E., Regius Professor of Technology in the University of Edinburgh. In fcap. 8vo. cloth, 2s. 6d.; elegantly bound in cloth, with richly gilt back and sides, and with gilt leaves, 5s. 6d.

[Just ready.]

**WILSON.**—A Treatise on Dynamics.

By W. P. WILSON, M.A., Fellow of St. John's, Cambridge, and Professor of Mathematics in the University of Melbourne. 8vo. bds. 9s. 6d.

**WRIGHT.**—Hellenica; or, a History of Greece in Greek, beginning with the Invasion of Xerxes; as related by Diodorus and Thucydides. With Explanatory Notes, Critical and Historical, for the use of Schools. By J. WRIGHT, M.A., of Trinity College, Cambridge, and Head-Master of Sutton Coldfield Grammar School. 12mo. cloth, 3s. 6d.

\*.\* This book is in use in Rugby and other Public and Private Schools.

**WRIGHT.**—A Help to Latin Grammar;

or, the Form and Use of Words in Latin. With Progressive Exercises. By J. WRIGHT, M.A. Crown 8vo. cloth, 4s. 6d.

**WRIGHT.**—The Seven Kings of Rome:

An easy Narrative, abridged from the First Book of Livy by the omission of difficult passages, so as to make a First Construing Book for Beginners of Latin. With Grammatical Notes. By J. WRIGHT, M.A. Fcap. 8vo. cloth, 3s.

## THE JOURNAL

OF

## CLASSICAL AND SACRED PHILOLOGY.

No. VIII. for June 1856, 4s.

Vol. I. for 1854, and Vol. II. for 1855, are now ready, cloth lettered, 12s. 6d. each.

CASES CAN BE HAD FOR BINDING VOLS. I. AND II.

\*.\* Three Numbers published annually, at 4s. each.

Cambridge: MACMILLAN & Co.

London: BELL & DALDY, 186, FLEET-STREET.

Edinburgh: EDMONSTON & DOUGLAS. Oxford: J. H. & JAS. PARKER.

Glasgow: WILLIAM ROBERTSON. Glasgow: JAS. MACLEHOSE.

R. CLAY, PRINTER, BREAD STREET HILL.





